

Heterosis and combining ability analysis in quality protein maize inbred lines

Neha Singhal, Sitar Singh Verma, Dinesh Chandra Baskheti* and Anil Kumar

Department of Genetics and Plant Breeding, College of Agriculture,
G.B. Pant University of Agriculture and Technology, Pantnagar-263 145 U.S. Nagar (U.A.) India

(Accepted : March, 2006)

In the present investigation, 42 crosses of quality protein maize (QPM) inbred lines (L1, L2,L14) with three testers (T1, T2 and T3) were evaluated against a QPM check, Shakti-1 in a randomized complete block design with 3 replications at G.B. Pant University of Agriculture and Technology, Pantnagar, India during Kharif, 2004. Observations were recorded on days to 50% tasselling and silking, plant height, ear height, ear length, ear diameter, kernel rows per ear, 100-kernel weight, grain yield and incidence of BLSB and BSDM. Protein, lysine and tryptophan contents were also estimated. The characters under study showed highly significant genotypic differences and wide range of variation in mean values. The crosses, L8 x T1, L1 x T3, L8 x T2, L5 x T2, L5 x T1, L2 x T1, L13 x T3, L3 x T1 and L8 x T3 revealed significant positive heterosis for grain yield. The gca effects of the parents showed that among lines, L8 was good for ear length, ear diameter, kernel rows per ear, 100-kernel weight and grain yield. L1, L12 and L13 were the other good general combiners. Based on the sca effects, L3 x T1 and L3 x T2 were good for ear length, ear diameter, kernel rows per ear, 100-kernel weight and grain yield. L14 x T3, L1 x T3, and L4 x T1 were the other specific cross combinations good for different characters. All the inbred lines and their crosses were moderately resistant to both the foliar diseases, BLSB and BSDM. The cross, L13 x T3 superceded the standard check for protein (9.74%), lysine (3.24%) and tryptophan (0.81%).

Key words : QPM, Top cross, GCA, SCA, Grain yield

INTRODUCTION

MAIZE (*Zea mays* L.), a major cereal crop cultivated all over the world for human food, animal feed as grain and fodder and many other industrial products like starch, oil and glucose etc., has poor quality of protein due to lower contents of two essential amino acids, lysine and tryptophan and an undesirable proportion of leucine and isoleucine.

The opaque-2 gene which is responsible for the enhancement of lysine and tryptophan of maize protein, if incorporated into any population, inbred line or any other germplasm of maize through back cross technique, improves the nutritional status of maize. The modified opaque-2 maize with hard endosperm and vitreous kernels is known as Quality Protein Maize (QPM). Its nutritional superiority over normal maize is well established (Gupta *et al.*, 1979). Concerted research efforts have been made on QPM in India and abroad particularly at CIMMYT, Mexico where Drs. S.K. Vasal and E. Villegas were jointly honoured by 'Millennium World Food Prize' during 2000 for their pioneering work on QPM.

In fact, the future is of hybrid cultivars. In QPM also, hybrids are more successful among the farmers as these are always planted with fresh F1 seed every year and thus remain unaffected from contamination through normal maize cultivars. Since the seed of open pollinated maize varieties can be saved and used for next 3-4 years, it gets contaminated if not grown in isolation. If the presently grown maize hybrids and varieties are replaced with QPM hybrids, the latter will provide a definite promise in upgrading the nutritional security of maize-based population. The QPM hybrids like Shaktiman-1, Shaktiman-2, Shaktiman-3, Shaktiman-4, Shaktiman-5, HQPM-1 and a QPM variety, Shakti-1 are under cultivation and catching the area up in eastern U.P., Bihar and some other states.

In a highly cross pollinated crop like maize, exploitation of heterosis can be accomplished through the development and identification of high *per se* performing vigorous parental inbred lines and their subsequent evaluation for combining ability in cross combinations to identify hybrids with high heterotic effects.

The magnitude of heterosis shown by the hybrids depends largely on the heterotic pattern and genetic divergence between parental lines. The information about the heterotic patterns and combining ability of parents and crosses both facilitate the breeders in the selection and development of single cross hybrids.

The present study was, therefore, undertaken to assess nature and magnitude of heterosis and combining ability effects for yield and other characters in a line x tester system using 14 QPM inbred lines and 3 testers.

MATERIALS AND METHODS

The experiment was conducted at Crop Research Centre, G.B. Pant University of Agriculture and Technology, Pantnagar, Uttaranchal during Kharif, 2004. The experimental material comprised 42 top crosses developed by crossing 14 QPM inbred lines of DMR series with 3 broad based testers abbreviated as L1, L2,L14 and T1, T2 and T3, respectively. The parental inbred lines and top crosses were evaluated against a QPM check, Shakti-1 in randomized complete block design with 3 replications. The net plot area was 7.5 m². Observations were recorded on whole plot basis for days to 50% tasselling and silking and fresh ear weight (later converted at 15% grain moisture for grain yield), whereas, the data on other characters like plant height, ear height, ear length, ear diameter, number of kernel rows per ear, 100-kernel weight were taken on 10 randomly selected competitive plants from every plot. The average value of these plants was calculated and used for analysis of variance, heterosis and combining ability. The incidence of two major foliar diseases namely, BLSB and BSDM was also recorded on 1-5 scale. Protein, lysine and tryptophan analysis was done using standardized laboratory procedures.

RESULTS AND DISCUSSION

QPM may make its greatest contribution in improving the protein quality of the diet of maize-based population. In past few years, many developing countries have released new QPM hybrids and varieties for the farmers and several others including India

* Author for Correspondence