

## The effect of temperature on the respiration of the fresh water bivalves, *Lamellidens marginalis*, from Godavari river, Nanded

S.K. PADEWAR, A.R. JAGTAP\*, A.B. HARKAL AND L.M. MUDKHEDE  
P.G. Department of Zoology, Yeshwant Mahavidyalaya, NANDED (M.S.) INDIA  
E-mail:ashu\_anamica@rediffmail.com

### ABSTRACT

The respiration rate of fresh water mussel, *i.e.* *Lamellidens marginalis*, edible fresh water bivalve mollusk. Mussels are able to move slowly by means of the muscular foot. They feed and breathe by filtering water through extensible tubes called siphons, a large muscle filters. *Lamellidens marginalis* is a species of freshwater bivalve in the Unionidae family. *Lamellidens marginalis* was monitored during the experimental manipulation of temperature. The oxygen consumed for one hour was measured at water temperatures ranging from 10° C to 25° C. The animals were acclimated to different temperatures in the temperature bath. The total oxygen consumption and rate of oxygen consumption per unit body weight along with shell was studied by Winkler's Method. The results showed that the rate of oxygen consumption increased by increasing temperature *i.e.* from 10° C to 25°C.

Padewar, S.K., Jagtap, A.R., Harkal, A.B. and Mudkhede, L.M. (2011). The effect of temperature on the respiration of the fresh water bivalves, *Lamellidens marginalis*, from Godavari river, Nanded. *Asian Sci.*, 6(1 & 2):6-8.

**Key Words :** Temperature effect, Oxygen consumption, Fresh water bivalve, *Lamellidens marginalis*.

### INTRODUCTION

Temperature is considered as a major environmental factor influencing most physiological rates, including those involved in the energy balance, such as feeding, excretion and respiration among bivalves (Bayne, 1976; Gabbott and Bayne, 1997). The temperature affects the metabolic activities of animals. The temperature affects on the dissolved oxygen concentration in the water so the ability to maintain a relatively constant uptake of oxygen under conditions of low oxygen availability is important for organisms that may encounter low dissolved oxygen frequently in their environment (Herried, 1980). In general, life activities occur within a range of about 0°C to 45°C. However, most animals live within much narrower limits. The effects of temperature on the metabolism of poikilotherms are elucidated through studies on oxygen consumption in relation to thermal stress of animals to wide geographical areas occurring at different latitudes. The effects of temperature on physiological processes studied for several species of bivalves (Melody, 1979; Mane, 1975; Vedpathak *et al.*, 1987; Widdows and Bayne *et al.*, 1991).

The economically important molluscan *i.e.* *Lamellidens marginalis* used as food source of man and it fulfill the human need of food to some extent in our country. But since last decade, the natural environment

is being disturbed due to global warming which results in the change in climatic temperature. The increasing temperature in climate affects the rate of oxygen consumption in fresh water bivalves. If the rate of oxygen consumption is not maintained at low levels of dissolved oxygen, anaerobic metabolism, which is inefficient with respect to energy production, may be activated in the tissues (Gade and Grieshaber, 1986). This ability to maintain oxygen consumption in water of reduced dissolved oxygen has been studied by number of investigators in bivalves (Wang and Widdows, 1993; Taylor and Brand, 1975; Sheldon and Walker, 1989). These studies indicate that the extent of metabolic regulation depends upon the environmental temperature.

The freshwater bivalves (*Lamellidens marginalis*) was selected for experimentation. It is abundantly available in Godavari river, Nanded. The present investigations were subjected to effect of temperature on fresh water bivalve. The purpose of this study was to determine the effect of temperature on the oxygen consumption of freshwater bivalves, *Lamellidens marginalis*.

### RESEARCH METHODOLOGY

The freshwater bivalves, *Lamellidens marginalis* used in the present study were collected from the

\* Author for correspondence.