

## Influence of nitrogen on quality parameters of different varieties of chrysanthemum

N.K. PATEL\* AND S.R. CHAUDHARI

ASPEE College of Horticulture and Forestry, Navsari Agricultural University, NAVSARI (GUJARAT) INDIA

### ABSTRACT

Investigation was carried out to evaluate the performance of different varieties under varying levels of nitrogen for quality parameters. Variety Flirt combine with 200 kg N per ha gave superior performance in all quality parameters like fresh weight, dry weight, stalk length and shelf life of flower (*in situ*) as compared to other varieties. While maximum vase life was observed in variety Flirt with 100 kg N per hectare and size of flower was largest in variety IIHR-6 with 200 kg N per hectare.

Patel, N.K. and Chaudhari, S.R. (2011). Influence of nitrogen on quality parameters of different varieties of chrysanthemum *Asian Sci.*, 6(1 & 2):63-65.

**Key Words :** Nitrogen, Chrysanthemum, Fresh and dry weight, Size, Stalk length, Vase and shelf life

### INTRODUCTION

Chrysanthemum is an important flower crop grown commercially in Karnataka, Andhra Pradesh, Maharashtra and Gujarat state. Small flower varieties of chrysanthemum are in great demand for making garland, veni and bouquet as well as in different religious occasions. Also there is great demand of quality flower in national and international flower market. So proper selection of variety and judicious use of nitrogen was quite imperative factor which use to sustained boosting in the quality parameters of chrysanthemum.

### RESEARCH METHODOLOGY

A field trial was conducted at the Floriculture Research Scheme, Regional Horticulture Research Station, Navsari Agricultural University, Navsari during 2008-2009. The soil of experimental field had a pH of (7.7), available nitrogen (160 kg/ha), available phosphorus (40.02 kg/ha), available potash (387.50 kg/ha). The experiment was laid out in Factorial Randomized Block Design with three nitrogen levels *i.e.* 100, 150 and 200 kg/ha and three varieties *viz.*, IIHR-6, Flirt and Shyamal. Basal dose of FYM was applied at rate of 20 tonnes per ha, phosphorus and potash were applied uniformly to each plot at the rate of 100 kg/ha. Nitrogen was applied in two splits; first half dose of nitrogen was applied as basal and second split was applied 30 days after transplanting. The data were collected on fresh and dry

weight of flower, size of flower, stalk length, vase life and shelf life (*in situ*) of flower.

### EXPERIMENTAL FINDINGS

The findings of the present experiment as well as relevant discussions have been presented under following heads :

#### Fresh weight of flower (g):

Fresh weight of flower affected significantly by nitrogen levels and varieties (Table 1). Maximum fresh weight of flower (2.937g) was recorded in the 200 kg N per hectare. Among different varieties, highest fresh weights of flower (2.928g) was obtained in Flirt. Improvement in fresh weight may be due to improvement in vegetative growth of plant under optimum level of nitrogen which caused more storage of carbohydrates and thus improved in fresh weight. Same results were earlier obtained by Jain and Gupta (2004) in marigold.

#### Dry weight of flower (g):

Data in Table 1 clearly reveal that the maximum dry weight of flower was recorded in 200 kg N per ha (0.653 g) where as incase of different varieties Flirt variety showed maximum dry weight (0.648 g). This might be due to the vigorous growth and development of plant under optimum nitrogen, ultimately produced maximum growth of flower and produced more dry matter. These results are in conformity with Vaghasia, (1997).

\* Author for correspondence.