

Studies on heritability and genetic advance in mutant populations of sesame (*Sesamum indicum* L.)

M.B. BORANAYAKA*, S.M. IBRAHIM¹, C.R. ANANDA KUMAR¹, D.S. RAJAVEL², T.V. SHADAKSHARI³ AND G. RAJESHA⁴

Zonal Agricultural Research Station, University of Agricultural Sciences, G.K.V.K., BENGALURU (KARNATAKA) INDIA email : mbboranayak@gmail.com

ABSTRACT

An investigation was undertaken to elucidate information on induced variability for yield and yield components in sesame (*Sesamum indicum* L.). The study consisted of two genotypes treated with physical (gamma rays) and chemical (EMS) mutagens. Two sesame varieties were treated with gamma rays ⁶⁰Co source with doses of 10, 20, 30, 40 and 50 krad followed by Ethyl methane sulphonate with concentrations of 0.8, 1.0, 1.2, 1.4 and 1.6 per cent. The LD₅₀ values based on germination reduction in the M₁ generation were fixed at 30krad and 1.2 per cent for gamma rays and EMS, respectively. High GCV for the traits plant height and number of capsules per plant in both the varieties was observed. Induced genetic variability was more in Cardeboriga than SVPR 1. The high heritability and genetic advance combined with increased genetic variability was realized for the characters viz., number of seeds per capsule and number of capsules per plant. The enhanced genetic variability observed for seed yield and its components in M₂ generation of the present study indicated the scope for effective selection.

Boraanayaka, M.B., Ibrahim, S.M., Kumar, C.R. Anand, Rajavel, D.S., Shadakshar, T.V. and Rajesha, G. (2011). Studies on heritability and genetic advance in mutant populations of sesame (*Sesamum indicum* L.). *Internat. J. agric. Sci.*, 7(2): 447-451.

Key words : Sesame, Gamma rays, EMS, Mutant population, Variability studies

INTRODUCTION

Mutational genetic manipulation of crop plants has been used very successfully to reconstruct crop ideotypes and improve a number of productive characteristics, in effect increasing the yield potential. This has been amply demonstrated in many crop species as exemplified by Xue Bai *et al.* (2000) in soybean, Shadakshari *et al.* (2001) in rice, Sharma (2001) in pea, Singh *et al.* (2001) in blackgram, Muthusamy and Jayabalan (2002) in cotton, Samiullah (2004) in blackgram, Singh (2006) in cowpea and Janila *et al.* (2007) in groundnut. More than 1500 mutant cultivars of crop plants with significantly improved attributes have been released worldwide in the last 30 years through induced mutation.

Sesame is probably the most ancient oilseed known and used by man. Success in any breeding programme depends on the amount of genetic variability present for the different characters in population. The genetic variability offered by mutagenic agents is of extreme importance in plant breeding. The variability in quantitative characters increases considerably by treating the biological

materials with different mutagenic agents. An estimation of the extent of variability induced in M₂ generation will be of great value to provide useful information for carrying out further selection.

MATERIALS AND METHODS

Two promising sesame genotypes namely, SVPR 1 (ruling popular white seeded type) and Cardeboriga (monostem African type) were treated with the two mutagens viz., gamma rays and EMS. Two hundred well filled dry seeds were sealed in butter paper covers and exposed to 10 to 50 krad doses of gamma rays from ⁶⁰Co source at Indira Gandhi Centre for Research, Kalpakkam, Tamil Nadu. Another variety of two hundred seeds of each variety, for each treatment were presoaked in distilled water for four hours then treated with different concentrations of EMS ranging from 0.8 to 1.6 per cent for three hours. After the treatment, the seeds were thoroughly washed with tap water ten times. The normal good looking plants based on base population randomly selected in each treatment in the M₁ generation were

* Author for correspondence.

¹ Department of Plant Breeding and Genetics, Agricultural College and Research Institute, MADURAI (T.N.) INDIA

² Department of Agricultural Entomology, Agricultural College and Research Institute, MADURAI (T.N.) INDIA

³ Department of Plant Breeding and Genetics, Tamil Nadu Agricultural University, COIMBATORE (T.N.) INDIA

⁴ Department of Plant Pathology, Tamil Nadu Agricultural University, COIMBATORE (T.N.) INDIA