



Effect of various agronomical practices in increasing and stabilising the yield of pigeonpea varieties

ASHALATA K. ZOTE, P.K. WAGHMARE AND V.B. SHELKE

ABSTRACT

The present investigation was undertaken to assess the effect of various agronomical practices in increasing and stabilising the yield of pigeonpea newly released disease resistant varieties (*i.e.* BSMR-736 and BSMR-853) by M.A.U. for large scale cultivation in the state. The agronomical practices like different row spacings and different planting patterns will help to enhance the productivity and stabilise the yield of the newly developed pigeonpea varieties. As regards to effects, the variety BSMR-853 proved superior over BSMR-736 in increasing the yield with various agronomical practices.

KEY WORDS : Pigeonpea, Varieties, Stabilising, Yield, Row spacings, Planting patterns

Zote, Ashalata K., Waghmare, P.K. and Shelke, V.B. (2010). Effect of various agronomical practices in increasing and stabilising the yield of pigeonpea varieties, *Internat. J. Forestry and Crop Improv.*, 1 (2) : 149-151.

INTRODUCTION

Since the primary objective of pigeonpea cultivation has been to meet domestic requirement for food and fuel with limited market surplus of grains, as such there was not much increase in production and productivity of pigeonpea. The major pigeonpea growing state in the country is the Maharashtra with an area of 10.41 lakh hectares and productivity of 834 kg/ha. This clearly indicate that there is not much increase in yield and stagnated over a period of time with substantial fluctuations inspite of availability of number of disease resistant varieties. Therefore, the present investigation was undertaken for study.

MATERIALS AND METHODS

A field experiment was conducted at college farm, Department of Agronomy, Marathwada Agricultural University, Parbhani during *Kharif* season of 2001 to study the effect of different row spacings and planting pattern

Correspondence to:

ASHALATA K. ZOTE, Marathwada Agricultural University, PARBHANI (M.S.) INDIA

Authors' affiliations:

P.K. WAGHMARE, Marathwada Agricultural University, PARBHANI (M.S.) INDIA

V.B. SHELKE, Directorate of Research Extension and Education, Marathwada Agricultural University, PARBHANI (M.S.) INDIA

on growth and yield of two newly released pigeonpea varieties. The site of the experimental field was fairly leveled. The residual effect of previous crop was uniform. The soil was clayey in texture, low in total nitrogen, medium in available phosphorus, rich in potassium and slightly alkaline in reaction.

The present experiment was laid out in Split Plot Design (SPD) with three replications. The treatments were of three row spacing (60, 90 and 120 cm), two planting patterns (normal and paired) and two varieties of pigeonpea (BSMR-736 and BSMR-853) constituting total twelve combinations. Six combinations of row spacing and planting pattern in main plots, while varieties of pigeonpea were randomly allotted as subplot treatments within each replication.

RESULTS AND DISCUSSION

The most important findings emerged from the present investigations are here-in. The mean plant height was not influenced significantly due to varieties under trial upto 90 DAS. However, the effect of the varieties on plant height were found significant then onwards *i.e.* the variety BSMR-736 recorded significantly taller plants than BSMR-853 at 120, 150 DAS and at harvest. As regards the number of functional leaves, the significant effect showed at only 90, 120 DAS and at harvest *i.e.* the number of functional leaves was significantly higher with the variety BSMR-736 compared to BSMR-853.