

Role of Isozymes in Rust (*Phakopsora pachyrhizi* Syd.) Resistance in Soybean

R. AMMAJAMMA AND P.V. PATIL

International Journal of Plant Protection, Vol. 2 No. 1 : 1-3 (April to September, 2009)

See end of the article for authors' affiliations

Correspondence to :

P.V. PATIL

Department of Plant Pathology, College of Agriculture, BIJAPUR (KARNATAKA) INDIA

SUMMARY

In the present investigation two known rust resistant (EC241778 and EC241780) and two susceptible (JS335 and DSb1) soybean genotypes have been used in order to know the differences in isozyme pattern. It revealed that at 45 DAS, PO peroxidase and polyphenol oxidase PPO isozymes shown monomorphism in all the genotypes (except in JS335 for peroxidase activity) irrespective of healthy and diseased leaves. At 75 DAS, both PO and PPO isozyme shown polymorphic activities under diseased condition indicating the PO and PPO isozyme synthesis is need based and polymorphic activities appear in response to infection by *Phakopsora pachyrhizi*. However, in resistant genotypes, bands with higher Rm values were found when compared to susceptible genotypes.

Key words :

Peroxidase, Polyphenol oxidase, Soybean, *Phakopsora pachyrhizi*, Electrophoresis.

Soybean [*Glycine max* (L.) Merrill] has been referred as gold due to its high protein content (40-43%) besides high oil content of 21-23 per cent. But the soybean production is largely influenced by repeated outbreak of the rust caused by *Phakopsora pachyrhizi* Syd. which can cause losses from 15 to 80 per cent depending on the locality, season and cultivar (Bromfield and Yang, 1976). The disease is appearing every year in epiphytotic form in rust endemic areas and causes substantial yield losses (Anonymous, 1995). Three triazole fungicides have been found effective in the control of soybean rust (Patil and Anahosur, 1998). However, continuous use of these chemical fungicides may pose the problem of development of resistance to pathogen. Therefore, the development of high yielding rust resistant varieties is of prime importance. In nature, some genotypes exhibit resistance to foliar diseases in any crop. The degree of resistance has been attributed to the genetic and biochemical make up of particular genotype, which needs to be investigated. Therefore, it is utmost important to study and find out the factors of resistance. Isoenzymes occurring within the same organism can be the rapid, sensitive laboratory tool for characterization of resistance in the genotypes. Keeping this in view, electrophoresis analysis of enzyme protein *i.e.* peroxidase (PO) and polyphenol oxidase (PPO) was attempted in four soybean genotypes.

MATERIALS AND METHODS

The investigations were carried out during

khariif 2004 at Main Agricultural Research Station, University of Agricultural Sciences, Dharwad. At the same time, pot experiment was conducted for maintaining the disease free plants in the glasshouse conditions. Two rust susceptible genotypes (JS335 and DSb1) and two rust resistant (EC241778 and EC 241780) soybean genotypes were used in this study. All the four genotypes were sown in the field in three replications in completely randomized block design. Uredospore suspension of *P. pachyrhizi* (10^5 spores/ml) was sprayed at 25 days after sowing. The sampling was done at 45 and 75 DAS and assessed the rust severity using 0 to 9 scale of Mayee and Datar (1986) and further per cent disease index (PDI) was calculated using the formula given by Wheeler (1969). For comparison, leaves of same age from healthy plants maintained in the glass house were used. Acetone powder of fresh composite leaf samples was prepared as per the procedure described by Brynt and Forrest (1979). The isozyme analysis of PO and PPO was done by using vertical slab gel electrophoresis technique with non discontinuous buffer system (Hames, 1990). Peroxidase isozyme was extracted by homogenizing 300mg of acetone powder of each sample in 5ml of chilled extraction buffer, pH 5.0 (Farkas and Stahmann, 1966). Electrophoresis was performed according to Davis (1964). The isozyme bands of peroxidase were localized by incubating the gel in guaiacol (0.25%) for 30 min. followed by incubation in 0.3% hydrogen peroxide for 15 min, which

Accepted :
October, 2008