

RESEARCH NOTE

Management of *Corynespora* blight of sesame through fungicides and antagonists

■ C.S. CHOUDHARY^{1*}, ANJANA ARUN¹ AND S.M. PRASAD²

¹Regional Research Station, Agwanpur, SAHARSA (BIHAR) INDIA

²Department of Plant Pathology, Birsa Agricultural University, RANCHI (JHARKHAND) INDIA

ARTICLE INFO

Received : 04.02.2014

Accepted : 28.03.2014

Key Words :

Fungicides, Fungal antagonists,
Corynespora blight, Sesame

***Corresponding author:**

Email: csrau07@gmail.com

ABSTRACT

Seed treatment with a mixture of Carbendazim 50 WP (0.1%) and Thiram 75 WP (0.15%) recorded minimum PDI of 11.15 per cent, respectively and 9.91 per cent and highest seed yield of 637kg and 646kg/ha during 2002-03 and 2003-04 crop seasons. First spraying of Carbendazim 50 WP (0.05%) and second spraying of *Trichoderma viride* formulation (10⁷ spores/g) proved to be the most economical.

How to view point the article : Choudhary, C.S., Arun, Anjana and Prasad, S.M. (2014). Management of *Corynespora* blight of sesame through fungicides and antagonists. *Internat. J. Plant Protec.*, 7(1) : 267-269.
