

Pigment analysis studies with reference to leaf reddening in Bt cotton

G.A. PAGARE AND D.V. DURGE

Received : July, 2010; Accepted : August, 2010

SUMMARY

Healthy cotton leaves has maximum chlorophyll content as compared to all reddening affected leaves. The reduction of chlorophyll content in reddening affected leaf was maximum in MRC-6301 Bt cotton variety. Anthocyanine content in reddening affected leaves was maximum in all the Bt cotton varieties as compared to healthy leaves of Bt cotton varieties at all the growth stages. Anthocyanine content increased in reddening affected leaves in the range of 4.94 to 5.06% at square formation stage and in general similar position was observed at rest of the growth stages.

Pagare, G.A. and Durge, D.V. (2010). Pigment analysis studies with reference to leaf reddening in Bt cotton. *Internat. J. Plant Sci.*, 6 (1): 42-44.

Key words : Reddening, Chlorophyll, Anthocyanine

In Bt. Cotton (*Gossypium hirsutum*) the reddening of leaves may appear at the square formation stage to boll development stage leading to great loss in yield. The yield reduction are reported by 30-60% depending upon the variety and reddening intensity (Bhatt and Patil, 1976). The information on pigment analysis is meagre. Attempts were made here to estimate what are the changes occur in chlorophyll content and in antocynine content in leaf during leaf reddening.

MATERIALS AND METHODS

Present investigation was undertaken on the field of Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola during the *Kharif* season of 2006 – 2007. Three cotton varieties, which were studied under this investigation, were TCHH-4, MRC-6301 and NCS-145. In these three varieties Bt gene was introduced, therefore, Bt gene introduced varieties were treated as Bt.

The observations were recorded at three stages *i.e.* square formation stage, boll development stage and boll bursting stage. Depending upon the reddening leaves were classified into two categories *viz.*, healthy and reddening affected leaves as suggested by snell and snell (1959). For these observations top fifth leaf was taken.

Correspondence to:

G.A. PAGARE, Department of Agricultural Botany, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, AKOLA (M.S.) INDIA

Authors' affiliations:

D.V. GURGE, Department of Agricultural Botany, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, AKOLA (M.S.) INDIA

The chlorophyll content in leaf were estimated with DMSO (Dimethyl sulphoxide) method by adopting the procedure given by Hiscox and Israelston (1979).

Fresh leaf discs weighting 37.5 mg were placed in 10ml of extractant and held for 2 hr. at 60°C. The supernant was used for estimation of pigments. The optical density of the aliquot was measured on spectronic- 20 at the wavelength of 652nm with red filter. The total chlorophyll content of leaves was calculated as per, Arnon (1949) equations and was expressed in (mg/g).

Anthocyanine content in leaf (ppm):

Fully developed red leaf was selected for the extraction of anthocyanine content of in leaf of Bt and Non-Bt cotton varieties. The extraction of anthocyanine from fully developed red leaf was done with DMSO (Dimethyl sulphoxide) method by adopting the procedure given by Hiscox and Israelstan (1979). Stock solution of pure anthocyanine having the strength of 25 ppm was prepared and by using these 25 ppm stock solutions of different ppm concentrations of 10 ml volume were prepared in the range of 1 ppm, 5 ppm, 10 ppm and 15 ppm. The optical density of these known concentration solutions was measured on Spectronic – 20 and accordingly stander curve was prepared by using these standard curve of unknown solutions.

RESULTS AND DISCUSSION

The results obtained from the present investigation as well as relevant discussion have been presented under following heads :