



RESEARCH
ARTICLE

Histomorphological and histochemical studies on isthmus of oviduct in Japanese quails

■ P. N. Thakur, P. J. Kapadnis¹ and N. M. Karad¹

Members of the Research Forum

Associate Author :

¹Department of Veterinary Anatomy,
College of Veterinary and Animal
Sciences, **Parbhani (M.S.) India**

AUTHOR FOR CORRESPONDENCE :

P. N. Thakur,
Department of Veterinary Anatomy,
College of Veterinary and Animal
Sciences, **Parbhani (M.S.) India**
Email : pravin_thakur75@rediffmail.com

Abstract : The mucosa of each segment presented lamina epithelialis, lamina propria and lamina mucosae. The mucosa was thrown into folds *viz.*, primary and secondary folds. The lamina epithelialis presented pseudostratified columnar ciliated epithelium was observed in isthmus, in both the groups of quail.

Key words : Histomorphological, Histochemical, Isthmus, Oviduct of Japanese quail

How to cite this paper : Thakur, P.N., Kapadnis, P.J. and Karad, N.M. (2019). Histomorphological and histochemical studies on isthmus of oviduct in Japanese quails. *Vet. Sci. Res. J.*, **10**(1) : 7-11, DOI : [10.15740/HAS/VSRJ/10.1/7-11](https://doi.org/10.15740/HAS/VSRJ/10.1/7-11). Copyright@2019: Hind Agri-Horticultural Society.

Paper History : Received : 02.02.2019; Revised : 12.03.2019; Accepted : 19.03.2019