

Seasonal Incidence of Leaf Webber, *Diaphania pulverulentalis* Hamps. on Mulberry in Kerala

TOMY PHILIP AND S.M.H. QADRI

International Journal of Plant Protection, Vol. 2 No. 1 : 24-26 (April to September, 2009)

See end of the article for authors' affiliations

Correspondence to :
TOMY PHILIP
Central Sericultural
Research and Training
Institute, Srirampuram,
MYSORE
(KARNATAKA)
INDIA

SUMMARY

A survey on the incidence of leaf webber, *Diaphania pulverulentalis* in Ernakulam and Trichur, the two major sericultural districts of Kerala was carried out. The survey was conducted at weekly intervals by the fixed plot method for three years (2002-2004). Generally, the infestation increases from June onwards and decreased from January. Maximum infestation was observed during the winter months. There was a gradual increased in the incidence of leaf webber year after year in all the villages in these districts. The average incidence was 2.2, 5.3 and 12.7, respectively during 2002, 2003 and 2004.

Key words :

Incidence, Kerala,
Leaf webber,
Mulberry.

Over 300 insect pests belonging to various Orders such as Lepidoptera, Hemiptera, Coleoptera, Thysanoptera, Orthoptera and Isoptera are known to attack mulberry and cause damage (Kotikal and Devaiah, 1987). They attack mulberry during different seasons, causing severe leaf yield loss and leaf quality. Since, majority of insect pests destroy tender leaves, young age silkworm rearing will be affected which in turn affect the later stage also. Leaf webber (*Diaphania pulverulentalis* Hamps.) has recently become a major problem for mulberry cultivators in South India causing a leaf yield loss of 12.8 per cent with an average incidence of 21.77 per cent (Siddegowda *et al.*, 1995; Rajadurai *et al.*, 1999). The pest incidence has been noticed in Karnataka since 1995 (Geetha Bai *et al.*, 1997) and has spread to neighbouring states of Tamil Nadu, Andhra Pradesh and Kerala. The early stage larvae inhabit the apical part of mulberry shoot and feed on the tender leaves. They bind together the apical leaves by a silky thread. Sometimes a single leaf is also rolled by the web with the larvae inside and hence the name, leaf roller/leaf webber has been given. Since the larvae feed on the tender leaves, the growth of the plant becomes stunted. Systematic survey on the distribution, seasonal incidence, natural enemies as well as qualitative and quantitative loss in different areas on mulberry are essential to formulate long-standing and effective control measures against the pest.

MATERIALS AND METHODS

A survey on the incidence of leaf webber was carried out in Ernakulam and Trichur, the two major sericultural districts in Kerala, at weekly intervals for three years (2002-2004). The survey was conducted by the fixed plot method. In Ernakulam district, Angamaly, Mallussery, Moozhikulam, Puthenvelikkara and Aduvassery villages and in Trichur district, Kottat, Thuruthiparambu, Valiyaparambu, Meladoor and Alathur villages were selected for the survey. In each of the above villages, four mulberry gardens were chosen. In each garden, five micro plots, at five different locations were demarcated. In each micro plot, twenty plants were selected randomly (100 plants in total) for recording the pest infestation. The percentage of incidence was calculated by using the formula—

$$\text{Incidence percentage} = \frac{\text{Total no. of infested plants}}{\text{Total no. of plants observed}} \times 100$$

Meteorological data such as temperature, relative humidity and rainfall during the survey period were also recorded to assess the influence of these factors on the pest incidence. The results are presented in Table 1 and Fig. 1 and 2.

RESULTS AND DISCUSSION

During 2002, maximum incidence of leaf webber was observed in the month of November (6.5%), when the average maximum temperature, minimum temperature, relative

Accepted :
December, 2008