

Research Paper :

Integrated biological management of chickpea wilt caused by *Fusarium axysporum* f. sp. *ciceri*

ASHISH SHRIVASTAVA AND VIJAY AGRAWAL

International Journal of Plant Protection (April, 2010), Vol. 3 No. 1 : 89-90

See end of the article for authors' affiliations

Correspondence to :
VIJAY AGRAWAL
Precision Farming
Development Centre,
CIAE, BHOPAL
(M.P.) INDIA

SUMMARY

Two isolates each of *Trichoderma viride* and *T. harzianum* and one of *Gliocladium virens* and *Bacillus subtilis* were used against *F. axysporum* f.sp. *ciceri*, Bio-agent tested *in vitro* by dual culture method, were found antagonistic to *F. axysporum* f.sp. *ciceri*. A maximum growth of inhibition of *F. axysporum* f.sp. *ciceri*, (44.8%) was reported by *T. varide*-1 (PDBCTv 23) followed by *T. varide*-2 (Gulbarga strain Tv) (39.8%), whereas *G. virens* was found least effective. Wilt incidence was also high in control (20.7, 26.7 and 35.3%) at 30, 45 and 60 days. *T. viride*-1 (PDBCT v 23) recorded only 4.9, 9.2 and 12.0% wilt incidence at 30, 45 and 60 days, respectively and was also found to be superior to all other treatments.

Key words :

Biological control,
Fusarium axysporum f.sp. *ciceri*,
Trichoderma viride, *T. harzianum*,
Chickpea, Wilt

Chickpea (*Cicer arietinum* L.) commonly known as channa or gram is one of the most important pulse crops in the Indian sub-continent. Wilt incited by *Fusarium axysporum* f.sp. *ciceri* is an important constraint in higher production of chickpea in Vindhyan Plateau Zone of Madhya Pradesh. In general resistant varieties, cultural practices and chemical fungicides are advocated for management of the disease but application of chemical fungicides proved short term measures as chemicals were reported to induce new strains of pathogen and also had environmental hazards on the other hands, for disease resistance, would constitute a medium term strategy due to the presence of several races (2, 3 and 5) of the pathogen. Various species of *Trichoderma* and *Gliocladium* had been studied for their bio-control ability against plant disease caused by *Fusarium* spp. (Lewis and Papavizas, 1980; Papavizas, 1985; Selvarajan and Jeyarajan, 1996), therefore, integrated biological management to keep the disease below economic injury level as long term measure. In this context, the present study was planned for the integrated management of chickpea wilt by the use of biotic agent.

MATERIALS AND METHODS

The pathogen was isolated from infected roots of chickpea, collected from 23, locations of Vindhyan Plateau Zone, by routine method on P.D.A. Soil samples collected from

rhizosphere of healthy chickpea plants just adjacent to wilted ones and also from wilt sick plants were used for isolation of micro organisms by using *Trichoderma* selective medium Method (Elad and Chet, 1983).

The antagonistic activity of micro organisms towards pathogen was tested on PDA medium in petriplate under sterilized conditions. Five mm disc of antiagonists were placed at four corners of petriplates in such a way that they were approximately 4.5 cm away from each other. The inoculated plates were placed in an incubator at 27±01°C for seven days after, radial growth of *F. axysporum* f.sp. *ciceri* was measured in(mm).

F. axysporum f.sp. *ciceri* was multiplied on sand maize meal medium (9:1) and fungal biocontrol agent were multiplied on wheat bran sand medium(1:9). *F. axysporum* f.sp. *Ciceri* and antagonists were added in quantities (1:1) in 15 cm plastic pots filled with sterilized soil. Surface sterilized seed of chickpea (JG 62) with 0.1% mercuric chloride for 2 minute, were sown in each pot. Observation on wilt incidence was recorded at 30, 45 and 60 days after sowing.

RESULTS AND DISCUSSION

Antagonistic potential of different biocontrol agent was noted in dual culture (Table 1). Bioassay test with these biocontrol agent revealed maximum zone of inhibition in case of *B. subtilis* followed by *T. viride*-2, *T.*

Accepted :
February, 2010