



Research Note

## Influence of water quality on efficacy of *Bacillus thuringiensis* var. *kurstaki* used as a spray fluid

■ V.S. DESAI<sup>1</sup>\* AND M.N. KAPADIA<sup>2</sup>

<sup>1</sup>Department of Agriculture Entomology, Dr. B.S. Konkan Krishi Vidyapeeth, Dapoli, RATNAGIRI (M.S.) INDIA

<sup>2</sup>Department of Agriculture Entomology, Junagadh Agricultural University, JUNAGADH (GUJARAT) INDIA

### ARTICLE INFO

#### Article Chronicle :

Received : 28.12.2011

Revised : 25.01.2012

Accepted : 14.03.2012

#### Key words :

*Bacillus thuringiensis* var. *kurstaki*,

*Earia vittella*, pH

\*Corresponding author: [vvpp\\_s@yahoo.co.in](mailto:vvpp_s@yahoo.co.in)

### ABSTRACT

The effect of quality of water on *Btk* formulation was studied at Junagadh Agriculture University, Junagadh, Gujarat. The influence of water quality on efficacy of *Btk* used as spray fluid revealed that water pH affected the efficacy of *Btk* formulation. The water with pH 7.0 to 8.5 was the most favourable for the efficacy of *Btk*. whereas, distilled water was more effective than tap water.

**How to view point the article :** Desai, V.S. and Kapadia, M.N. (2012). Influence of water quality on efficacy of *Bacillus thuringiensis* var. *kurstaki* used as a spray fluid. *Internat. J. Plant Protec.*, 5(1) : 187-188.

The *Bacillus thuringiensis* var. *kurstaki* (*Btk.*) is being used all over India for the control of different crop pests. It is now a day's very popular bio pesticide available in market that has been exploited and showed very good effect as compared to chemical insecticides. The water used by the farming community of the country for the preparation of the *Btk.* suspension is whatever available in the vicinity of the field. The available water is of different quality in the different regions as well as in different seasons. It is well known fact that the commercial formulations of *Btk.* available in market are d endotoxin based and the toxin is very well affected by pH. This indicated that there would be some effect of pH of the water that is used to prepare spray fluid on the *Btk.* formulation. The present experiment was conducted to study the influence of quality of water on *Btk.* formulation.

The present investigation was carried out to study the influence of quality of water that is used as a spray fluid on the efficacy of *Btk.* at Department of Agril. Entomology, Junagadh Agriculture University, Junagadh under laboratory conditions.

In order to know the influence of water quality on effectiveness of *Btk.*, different categories of water *viz.*, pH 6.0, 7.0, 7.5, 8.0, 8.5, 9.0, tap water and distilled water were studied. The experiment was conducted in C.R.D with four replications. The different quality of water having different pH was prepared. The *Btk.* formulation @ 1.5 g/l was mixed in

the water that is having different pH. In bioassay, okra fruits were cut into slices with the help of knife. These slices were immersed in a suspension of *Btk.*, having different quality of water for 30 seconds and the excess fluid was removed by uniform jerking. These treated okra slices were air dried for a half hour. The treated okra fruit slices were placed separately inside the Petridishes (15 cm diameter x 2.5 cm height) over moist blotting paper. A single 24-hour starved third instar larva was released on each slice. For each treatment, ten larvae were used. The larvae were allowed to feed on the treated fruit slice for 24 hours and then fresh, untreated okra fruit slices were provided daily to the larvae for feeding. Mortality counts were taken at 1, 2, 3 and 4 days after exposure. The larval mortality was converted into percent larval mortality.

The results pertaining to laboratory experiment on influence of water quality on toxicity of *Btk.* against third instar larvae of *E. vittella* are presented in Table 1.

There was no any larval mortality observed one day after feeding. After two days of feeding, the larval mortality of 40.00 to 45.00 per cent was observed in all the treatments but there was no significant difference among the treatments.

The highest larval mortality (75.00 %) was observed in the treatment with distilled water and was found to be best treatment. This was followed by pH 7.0, 7.5 and 8.0, which recorded 70.00, 67.50 and 67.50 per cent, larval mortality, respectively and was at par with above treatment. The