

RESEARCH NOTE

Survey and surveillance of early blight of tomato caused by *Alternaria solani* in Konkan region

■ V.G. MUNDE, M.P. DIWAKAR, B.B. THOMBRE AND UTPAL DEY

Department of Plant Pathology, College of Agriculture, Dr. B.S. Konkan Krishi Vidyapeeth, Dapoli, RATNAGIRI (M.S.)
INDIA

ARTICLE INFO

Received : 13.08.2012

Accepted : 16.05.2013

Key Words :

Survey, Foliar disease, Tomato, Konkan region

***Corresponding author:**

Email: utpaldey86@gmail.com

ABSTRACT

A roving survey of early blight of tomato disease were carried out during *Rabi* season, 2008-2009 at Thane, Raigad, Ratnagiri and Sindhudurg districts and per cent disease incidence was recorded ranged from 26.33 to 50.77 per cent. The mean incidence of early blight recorded was 40.97 per cent in Konkan region. The maximum per cent disease incidence was observed in Wada Tahsil in Thane district with 50.77 per cent and minimum per cent disease incidence in Goregaon Tahsil of Raigad districts with 26.33 per cent in Konkan.

How to view point the article : Munde, V.G., Diwakar, M.P., Thombre, B.B. and Dey, Utpal (2013). Survey and surveillance of early blight of tomato caused by *Alternaria solani* in Konkan region. *Internat. J. Plant Protec.*, **6**(2) : 476-477.
