

RESEARCH ARTICLE

# *In vitro* evaluation of fungicides against *Fusarium oxysporum* f. sp. *cubense*

■ R. SOMU<sup>1</sup>, N. THAMMAIAH<sup>2\*</sup>, G.S.K. SWAMY<sup>3</sup>, M.S. KULKARNI<sup>4</sup> AND V. DEVAPPA<sup>4</sup>

<sup>1</sup>Department of Plant Pathology, K.R.C. College of Horticulture, ARABHAVI (KARNATAKA) INDIA

<sup>2</sup>K.R.C. College of Horticulture, ARABHAVI (KARNATAKA) INDIA

<sup>3</sup>Department of Fruit Science, K.R.C. College of Horticulture, ARABHAVI (KARNATAKA) INDIA

<sup>4</sup>Department of Horticultural Plant Pathology, K.R.C. College of Horticulture, ARABHAVI (KARNATAKA) INDIA

---

## ARTICLE INFO

**Received** : 31.01.2014

**Revised** : 17.03.2014

**Accepted** : 28.03.2014

## Key Words :

*Fusarium oxysporum* f. sp. *cubense*,  
Fungicides, Banana

**\*Corresponding author:**

Email: nthammaiah@gmail.com

---

## ABSTRACT

Among the six fungicides evaluated in *in vitro* against *Fusarium oxysporum* f.sp. *cubense*, carbendazim, carboxin, propiconazole and benomyl showed total inhibition of the fungal growth at the concentrations of 500, 1000 and 2000 ppm. However, difenconazole showed total inhibition of the fungal growth at 2000 ppm concentration.

**How to view point the article :** Somu, R., Thammaiah, N., Swamy, G.S.K., Kulkarni, M.S. and Devappa, V. (2014). *In vitro* evaluation of fungicides against *Fusarium oxysporum* f. sp. *cubense*. *Internat. J. Plant Protec.*, 7(1) : 221-224.