

RESEARCH NOTE

Residual toxicity of *Nomuraea rileyi* (Farlow) Samson against *Helicoverpa armigera* (Hubner) on pigeonpea

■ A.H. BARAD, M.N. KAPADIA AND D.M. JETHVA*

Department of Entomology, Junagadh Agricultural University, JUNAGADH (GUJARAT) INDIA

ARTICLE INFO

Received : 01.11.2013

Accepted : 20.03.2014

Key Words :

Persistent effect, *Nomuraea rileyi*,
Pigeonpea

***Corresponding author:**

Email: dr_dharmraj@yahoo.co.in

ABSTRACT

A field – cum – laboratory trial was conducted to determine the persistent effect of *Nomuraea rileyi* (Farlow) Samson against *Helicoverpa armigera* (Hubner) on pigeonpea at Junagadh Agricultural University Campus, Junagadh during *Kharif* season of 2009-10. The results revealed that *Nomuraea rileyi* persisted up to 11 days of application with expressing the maximal persistency up to 5 days of its application on pigeonpea pods. The larval mortality became half within seven days.

How to view point the article : Barad, A.H., Kapadia, M.N. and Jethva, D.M. (2014). Residual toxicity of *Nomuraea rileyi* (Farlow) Samson against *Helicoverpa armigera* (Hubner) on pigeonpea. *Internat. J. Plant Protec.*, 7(1) : 246-247.
