

## Biology of *Tetranychus urticae* Koch (Acarina: Tetranychidae) on carnation under laboratory conditions

■ DINESH L. PATIL\*<sup>1</sup>, K.A. PATEL<sup>2</sup>, N.R. TOKE<sup>1</sup> AND ARCHANA T. AMBULE<sup>1</sup>

<sup>1</sup>Department of Entomology, N.M. College of Agriculture, Navsari Agricultural University, NAVSARI (GUJARAT) INDIA

<sup>2</sup>Main Sorghum Research Station (N.A.U.), SURAT (GUJARAT) INDIA

### ARTICLE INFO

**Received** : 31.03.2014  
**Revised** : 26.07.2014  
**Accepted** : 12.08.2014

### KEY WORDS :

*Tetranychus urticae*, Biology,  
Carnation

### ABSTRACT

Biology of *Tetranychus urticae* Koch was studied under laboratory at  $27.79 \pm 4.40^\circ\text{C}$  temperature and  $79 \pm 5.75$  per cent RH during Sept. 2008. The finding showed that life cycle of *T. urticae* consisted of five stages viz., eggs, larva, protonymph, deutonymph and adult and one short inactive stage called quiescent stage. The eggs incubation period of *T. urticae* was  $3.29 \pm 0.62$  days. The larval body measured  $0.14 \pm 0.01$  mm in length and  $0.10 \pm 0.01$  mm in width and lived for  $1.79 \pm 0.64$  days in male and  $2.71 \pm 0.57$  days in female. The mature larva entered in a quiescent stage (nymphochrysalis). The total developmental period was longer in female ( $10.61 \pm 1.21$  days) as compared to male ( $8.44 \pm 1.75$  days). Mated female lived for longer time and laid higher number of eggs than unmated female. Mated female produced progeny consisting of both sexes, whereas, unmated female invariably gave rise to male progeny.

**How to view point the article** : Patil, Dinesh L., Patel, K.A., Toke, N.R. and Ambule, Archana T. (2014). Biology of *Tetranychus urticae* Koch (Acarina: Tetranychidae) on carnation under laboratory conditions. *Internat. J. Plant Protec.*, 7(2) : 334-338.

\*Corresponding author:

Email: [patildinesh7901@gmail.com](mailto:patildinesh7901@gmail.com)