

Pharmacognostical and Phytochemical studies on the leaves of *Lantana camara* Linn

E. EDWIN*, E. SHEEJA AND P. JAIN

Department of Pharmacognosy & Phytochemistry, B.R. Nahata College Pharmacy, Mhow-Neemuch Road, Mandsaur - 458001, Madhya Pradesh.

(Accepted : December, 2005)

SUMMARY

According to the literatures, the plant *Lantana camara* (Linn.) belonging to the family Verbenaceae, have a great medicinal value. So far proper pharmacognostical and phytochemical studies are not been reported for the leaves. In this direction our efforts were devoted to determine the pharmacognostical and preliminary phytochemical properties of this plants. Pharmacognostical evaluation included examination of morphological and microscopical characters, determination of leaf constants (stomatal number, index, veinlet and vein termination nos), determination of physicochemical constants (ash values). Phytochemical screening included, qualitative chemical examinations, determination of % yield and extractive values. The studied characters would be an useful tool to identify this plant by its Pharmacognostical characters.

Key words: *Lantana camara*, Verbenaceae, rheumatism

India is one of the richest floristic regions of the world and has been a source of plants and their products since antiquity and man uses them in different ways according to his needs, particularly as food and medicine. Among the entire flora, 35,000 to 70,000 species have been used for medicinal purposes Ponnu *et al.*, (2003). The plant *Lantana camara* Linn (Verbenaceae) commonly known as Lantana is used for fever, cold, rheumatism, high blood pressure, etc E. L. Ghisalberti, (2000). The reported constituents in the plant are essential oils, lantadenes A-D, lantanilic acid, icterogenin, farnesene, caryophyllene, flavonoids, etc The wealth of India, (2000). The main part used in these plants is leaves. So far proper pharmacognostical and phytochemical studies are not been reported for the leaves. Therefore our efforts were devoted in this direction to study the pharmacognostical and phytochemical aspects of *Lantana camara*.

MATERIALS AND METHODS

The plant materials (leaves) of *Lantana camara*. Linn were collected in Mandsaur and was positively identified and confirmed by taxonomist in the KNK College of Horticulture, Mandsaur.

Pharmacognostical Studies:

Morphological studies were done using simple microscope. The shape, apex, base, margin, taste and odor of leaves were determined. Microscopical studies were done by preparing a thin section of midrib and lamina region of *Lantana camara*. The section was cleared with chloral hydrate solution and then stained with phloroglucinol and hydrochloric acid, mounted in Glycerin. A separate section was prepared and stained with iodine solution for the

*Author for correspondence

identification of starch grains. The powders of the dried leaves were used for the observation of powder microscopical characters. The powdered drug was separately treated with Phloroglucinol - HCl solution and iodine solution to determine the presence of lignified cells and starch grains C.K. Kokate, (1994).

As a part of quantitative microscopy, stomatal number, stomatal index, vein islet, vein termination number were determined by using fresh leaves of the plant. Ash values are used to determine the quality and purity of the crude drug. Total ash and acid insoluble ash were determined. Alcohol and water-soluble extractive values were determined to find out the amount of water and alcohol soluble components C. K. Kokate, (1994); Indian Pharmacopoeia, (1985).

Phytochemical Studies:

The dried powder material was initially defatted with petroleum ether (60-80°C) in a soxhlet apparatus for 72 hrs and successively extracted with ethanol and water for 72 hrs. The extracts were filtered while hot and solvent removed by distillation under reduced pressure and the percentage yields of the extracts were calculated.

The concentrated Pet. ether, (60-80°C), ethanol and water extracts of the leaves were subjected to chemical test for the identification of the various active constituents J. B. Harborne, (1988); A. Mohammed, (1994); O. P. Agarwal, (2000).

RESULTS AND DISCUSSION

The morphological studies revealed the shape of leaves as ovate, deep green on the upper surface and light at the lower, with serrate margin, symmetrical base,