

Effect of harvesting stages on seed quality parameters of groundnut (*Arachis hypogaea* L.)

■ A.P. GAIKWAD AND R.W. BHARUD

SUMMARY

The studies revealed that, in six varieties of groundnut seed quality parameters viz., number of well developed pods per plant increased after physiological maturity. 100 seed weight (g) and seed moisture content (%) decreased after physiological maturity.

Key Words : Groundnut, Pods, 100 seed weight, Moisture

How to cite this article : Gaikwad, A.P. and Bharud, R.W. (2014). Effect of harvesting stages on seed quality parameters of groundnut (*Arachis hypogaea* L.). *Internat. J. Plant Sci.*, 9 (1): 196-198.

Article chronicle : Received : 08.10.2013; Revised : 10.11.2013; Accepted : 19.11.2013

MEMBERS OF THE RESEARCH FORUM

Author to be contacted :

A.P. GAIKWAD, Department of Agricultural Botany, Mahatma Phule Krishi Vidyapeeth, Rahuri, AHMEDNAGAR (M.S.) INDIA

Address of the Co-authors:

R.W. BHARUD, Department of Agricultural Botany, Mahatma Phule Krishi Vidyapeeth, Rahuri, AHMEDNAGAR (M.S.) INDIA
