



RESEARCH PAPER

DOI: 10.15740/HAS/TAJH/9.2/400-403

Article history :

Received : 07.04.2014

Revised : 29.10.2014

Accepted : 10.11.2014

Effect of different levels of row spacing and nitrogen on growth characters in roselle (*Hibiscus sabdariffa* var. *sabdariffa*)

■ **A.K. RAJA¹ AND E. ARIVAZHAGAN**

Members of the Research Forum

Associated Authors:

¹Department of Horticulture, Faculty of Agriculture, Annamalai University, ANNAMALAINAGAR (T.N.) INDIA

Author for correspondence :

E. ARIVAZHAGAN

Department of Horticulture, Faculty of Agriculture, Annamalai University, ANNAMALAINAGAR (T.N.) INDIA
Email : gskspice@gmail.com

ABSTRACT : The present investigation was carried out to study the effect of different levels of nitrogen and spacing on the growth characters in roselle. A field trial was conducted in a Factorial Randomized Block Design with five levels of nitrogen (50, 100, 150, 200 and 250 kg ha⁻¹) and three levels of row spacing (60, 90 and 120 cm). The results of the present study revealed that the spacing of 120 × 75 cm and nitrogen application of 250 kg ha⁻¹ was found to be superior in improving the growth characters like plant height, number of branches plant⁻¹, stem diameter, plant spread and leaf area.

KEY WORDS : Nitrogen, Spacing, Roselle, Growth

HOW TO CITE THIS ARTICLE : Raja, A.K. and Arivazhagan, E. (2014). Effect of different levels of row spacing and nitrogen on growth characters in roselle (*Hibiscus sabdariffa* var. *sabdariffa*). *Asian J. Hort.*, 9(2) : 400-403.