

RESEARCH PAPER

Length weight relationship of *Laubuka laubuca*, *Salmophasia phulo* and *Esomus danricus* (Hamilton, 1822) from lower Brahmaputra drainage of Assam, North-East India

Chandana Kalita and Dandadhar Sarma

Department of Zoology, Gauhati University, Guwahati (Assam) India

Email : sarma_dandadhar@yahoo.com; ckalita30@gmail.com

Article Info : Received : 18.08.2018; Revised : 10.09.2018; Accepted : 21.09.2018

Length-weight relationships (LWRs) for three cyprinids (*Laubuka laubuca*, *Salmostoma phulo* and *Esomus danricus*) collected seasonally from May 2016 to January 2017 along the lower Brahmaputra drainage in Assam (Northeast India), using fishing gears [cast nets: 2.5 m, 10–15 mm mesh size; gillnets: 30 × 0.9 m, 18–20 mm mesh size], were estimated. The 'b' values in the LWRs were 3.46 for *L. laubuca*, 2.98 for *S. phulo* and 3.13 for *E. danricus*, respectively.

Key words : Length weight, *L. laubuca*, *S. phulo*, *E. danricus*

How to cite this paper : Kalita, Chandana and Sarma, Dandadhar (2018). Length weight relationship of *Laubuka laubuca*, *Salmophasia phulo* and *Esomus danricus* (Hamilton, 1822) from lower Brahmaputra Drainage of Assam, North-East India. *Asian J. Bio. Sci.*, **13** (2) : 62-64. DOI : 10.15740/HAS/AJBS/13.2/62-64. Copyright © 2018: Hind Agri-Horticultural Society.