

## Acute toxicity of Abamectin to fishes, *Cyprinus carpio* Linnaeus and *Tilapia mosambica* (Peters)

R. SHEEBA JASMINE\*, S. KUTTALAM AND J. STANLEY

Department of Agricultural Entomology, Tamil Nadu Agricultural University, COIMBATORE, (T.N.)INDIA

(Accepted : February, 2008)

Laboratory studies conducted to determine the acute toxicity of abamectin, a broad spectrum pesticide to fresh water common carp, *Cyprinus carpio* (Linnaeus) and *Tilapia mosambica* (Peters) revealed that the fish are highly susceptible to abamectin. The median lethal concentration ( $LC_{50}$ ) of abamectin for *C. carpio* and *T. mosambica* were 0.475 and 6.928 ppb, respectively. Likewise, the median lethal time ( $LT_{50}$ ) were 3.119 and 1.804 h for abamectin at 0.50 and 0.55 ppb concentration to *C. carpio* and 19.039, 13.933 h for abamectin at 8.0 and 9.0 ppb concentration to *T. mosambica*, respectively.

Key words: Abamectin, *Cyprinus carpio*, *Tilapia mosambica*,  $LC_{50}$ ,  $LT_{50}$

### INTRODUCTION

Extensive use of broad spectrum synthetic chemicals results in the destruction of non target organisms where they are transported. These synthetic insecticides have a long life period, which has resulted in the process of bioaccumulation and biomagnification in the environment and in the living organisms (Sahai, 1992). The toxicity of the insecticides has a greater impact on the aquatic environment as these insecticides enter into the hydrosphere via many pathways like, direct application for pest and disease vectors, surface runoff from non-point sources including agricultural soil, aerosol and particulate deposition, rainfall and absorption from the vapour phase at the air-water inter phase etc. The wide spread usage of synthetic insecticides ultimately pollute the aquatic environment there by affecting the aquatic fauna mainly fishes which constitute the major economy of the country and a valuable source of protein. Moreover, fish is one of the components of integrated farming in certain parts of the country. It is observed that the structure and stability of natural fish populations would be adversely affected if the water where their development takes place is contaminated even at sublethal concentrations.

### MATERIALS AND METHODS

The acute toxicity of abamectin to common carp was assessed as per Sprague (1969 and 1970). The aquarium tanks of dimensions 15 cm x 30 cm x 45 cm with 80 l capacity were provided with artificial aeration facilities. Abamectin 1.9 EC at different concentrations viz., 7, 9,

10, 11 and 12  $\mu$ l 20 l<sup>-1</sup> of water were taken as different treatments after range finding test and control tanks were maintained without abamectin. Each experiment was replicated four times. In each tank, a school of twenty fish of same size was released. Observations on mortality were made on 3, 6, 12, and 24 h and per cent mortality was worked out for each concentration and sufficient replicates were used to construct a reliable regression line known as log concentration probit mortality (lcpm) line/ curve. The mortality data obtained were corrected for the mortality in control by Abbott's formula (Abbott, 1925) and the data were subjected to probit analysis (Finney, 1971) and confirmed by EPA probit analysis program used for calculating LC/ EC values Version 1.5. Another experiment to assess the acute toxicity of abamectin to *T. mosambica* was conducted in the same method. The fishes were exposed to slightly higher concentrations of 100, 120, 140, 160 and 180  $\mu$ l of abamectin in 20 l<sup>-1</sup> of water after range finding test and the mortality observed and statistically analysed.

### RESULTS AND DISCUSSION

The  $LC_{50}$  values of abamectin to fresh water common carp *C. carpio* and a common fish, *T. mosambica* were 0.475 and 6.928 ppb, respectively (Table 1) (Fig.1 & 2). The median lethal time ( $LT_{50}$ ) were 3.119 and 1.804 h for abamectin at 0.50 and 0.55 ppb concentration to *C. carpio* and 19.039 and 13.933 h for abamectin at 8.0 and 9.0 ppm concentration to *T. mosambica*, respectively (Table 2). Toxicity of abamectin for the two fish tested

\* Author for Correspondence