

Effect of fungicides on soft rot of Turmeric

SHAIKH RUHINA AND B.M. KAREAPPA*

Department of Biotechnology, D.S.M., College, PARBHANI (M.S.) INDIA

(Accepted : June, 2008)

Turmeric is infected by *Phytophthora myriophyllum* Drch. causing soft rot. In order to control soft rot, the different fungicides used were Mancozeb, Thiram and Carbendazim. The concentration used for the treatment was from 0.25 to 2.0% by using slice method. Among these fungicides, thiram shows highest sensitivity by inhibiting the maximum growth of the pathogen. Carbendazim shows intermediate inhibition and mancozeb least inhibition of the pathogen. This clearly indicates that these fungicides can be used to control soft rot of turmeric caused by *Phytophthora myriophyllum* Drch.

Key words : Turmeric, Soft rot, Fungicides.

INTRODUCTION

Haldi or turmeric (*Curcuma longa* L.) is one of the most important condiment and colouring agents of the world. Turmeric is also a commodity of auspicious religious ceremonies in India. It is being used essentially in curry powder and is a common food preservative. Turmeric is grown in the states of Tamil Nadu, Kerala, Assam, Andhra Pradesh, West Bengal and Maharashtra. 'India produced six lakh metric tones of turmeric rhizomes on 1.4 lakh hectares in 1997 - 98 (Anonymous, 1999). Maharashtra produced 38,202 tonnes of turmeric in 6,191 ha. with the productivity of 6.17 tonnes per ha. (Anonymous, 2000).

Turmeric plays a very important role in agro-based industries. Domestically it is an important condiment, colouring and flavouring agent used in prepared dishes. It is important meat dressing materials and used in salad. Additionally, it has pharmaceutical value too. It's colour properties are due to crystalline pigment "curcumin". The dried rhizome has 0.5 - 0.6% curcumin and 5 - 6% volatile oil (Mogle, 1999).

In the Marathwada region, a step rise in the coverage under turmeric crop is evident and with the enhanced irrigation facilities due to Puma and Jayakwadi Project more area is likely to be occupied by this cash crop.

Therefore, the present investigation has been carried out on the fungal rots of turmeric caused by *Phytophthora myriophyllum Dreschi* and their suitable control measures which possibly would ensure enhanced yield and also improve planting values of the mother sets or finger sets.

MATERIALS AND METHODS

In vivo evaluation of fungicides :

For evaluation of different fungicides, cut rhizome method was followed. Each fungicides was tested at different levels of concentrations *i.e.* 0.25, 0.50, 0.75, 1.0, 1.25, 1.50, 1.75 and 2.0%. The cut rhizome of size 2 cm in diameter and placed on blotter paper. A 5mm inoculum disc of the pathogen (*Phytophthora myriophyllum*) was inoculated in the centre of each cut rhizomes. The treatment was given in three replications. The observations were recorded after 7 days of incubation at room temperature ($28^{\circ}\text{C} \pm 1^{\circ}\text{C}$).

RESULTS AND DISCUSSION

To ascertain the sensitive conc. of fungicides in controlling rot rhizomes of turmeric were subjected to the different conc. of the fungicides.

From the data presented in Table 1, showed that as the conc. of fungicide increased there was decline in the growth of pathogen. All the fungicidal conc. reduced

Table 1 : Effect of various conc. of mancozeb on rhizome rot of turmeric caused by *Phytophthora myriophyllum*

Sr. No.	Conc. of fungicides (%)	Linear growth (mm)
1.	0.25	13
2.	0.50	12.5
3.	0.75	12.1
4.	1.0	11
5.	1.25	9
6.	1.50	8.7
7.	1.75	8.3
8.	2	8
9.	Control	18

* Author for correspondence.