

Solubilization of insoluble zinc compounds by *Gluconacetobacter diazotrophicus* and its influence on maize

C. SARATHAMBAL, M. THANGARAJU AND M.GOMATHY

Department of Agricultural Microbiology, Tamil Nadu Agricultural University, COIMBATORE (T.N.) INDIA

(Accepted : March, 2009)

Zinc deficiency was created in maize seedlings in the sand culture and deficiency was rectified with the addition of insoluble zinc sources and *G. diazotrophicus*. The basic strategy behind this approach is decreasing the pH and making zinc soluble and as a consequence, the available zinc will get increased in the soil system. Since lowering of soil pH to 5 or below increases the Zn solubility. The plant growth, chlorophyll content and plant zinc content were found to be enhanced due to inoculation of *G. diazotrophicus* with insoluble zinc compounds.

Key words : *G. diazotrophicus*, Zinc solubilization, Chlorophyll content.

INTRODUCTION

Zinc plays amazing number of critical roles in all organisms. It required in small but critical concentration to allow several key plant physiological pathways to function normally. These pathways have important roles in photosynthesis, sugar formation, protein synthesis, fertility and seed production, growth regulation and defence against disease. Unlike other micronutrients deficiency, Zn deficiency is ubiquitous (Guerinot and Eide, 1999). Where zinc is deficient, morphological functions will be impaired and the health and productivity of the plants will be adversely affected, resulting in lower yields (or even crop failure) and frequently in poor quality crop products.

The whole world is shifting to an organic based sustainable agriculture and is also our duty to preserve the nature's wealth without any pollution for the future generation. So the need of the hour is to devise some technology or strategies to alleviate zinc deficiency. Heterotrophic bacterial zinc solubilization was reported in literatures (Di Simine *et al.*, 1998; Coles *et al.*, 2001; Whiting *et al.*, 2001 and Fasim *et al.*, 2002) focusing mainly on the remediation aspects of zinc pollution. Recently a bacterial based approach was devised to solve these micronutrient deficiency problems (Saravanan, 2004 and Anthoni Raj, 2002).

MATERIALS AND METHODS

A pot culture experiment was conducted to study the zinc solubilization potential of *G. diazotrophicus*. Acid washed sand was used as medium for the experiment. About 2.5 kg of acid washed sand was taken in plastic pot (size:

15cm height, 15cm diameter) and sterilized at 20 lbs pressure for 2hrs each for two consecutive days. Maize seeds (Col) obtained from the Department of Millets, Tamil Nadu Agricultural University, Coimbatore were used. Watering was done with sterile distilled water. Fahraeus N-free nutrient solution without zinc source was used as the nutrient solution for maize. Initially 30 ml of *G. diazotrophicus* culture was added for each pot and every 10 days another 20ml was added. One gram fresh leaf sample taken from each replication was analyzed for total chlorophyll content following the method of Hiscox and Isrealstam (1979). Total zinc content of maize seedlings was analyzed according to Jackson, 1973. The shoot and root length of maize plant was measured. According to treatment details Zn is added in the form of insoluble sources (zinc oxide, zinc carbonate and zinc phosphate).

The treatment details of this experiment are as :

T₁ - Uninoculated control+ without Zn, T₂ - ZnCO₃ alone, T₃ - ZnCO₃ + *G. diazotrophicus*, T₄ - ZnO alone, T₅ - ZnO + *G. diazotrophicus*, T₆ - Zn₃(PO₄)₂ alone, T₇ - Zn₃(PO₄)₂+ *G. diazotrophicus*, T₈ - ZnSO₄

RESULTS AND DISCUSSION

The maize seedlings in pot culture experiments developed Zn deficiency after two months of sowing. This deficiency was corrected with the application of insoluble forms of Zn viz., ZnCO₃, ZnO and Zn₃(PO₄)₂ along with inoculation of *G. diazotrophicus*. The inoculated organism might have released the gluconic acid during its growth which in turn solubilize the insoluble zinc compounds there by making available the zinc for plants