
RESEARCH PAPER

Effect of *Rhizobium japonicum* in relation to nodulation and chlorophyll content of soybean

D.C. KAMTHANE

Department of Microbiology, Shri Guru Buddhiswami College, Purna, PARBHANI (M.S.) INDIA

Email : daiwa.kamthana@rediffmail.com

The soybean (*Glycine max*), commonly called 'Cindrella crop' or 'king of legumes' *R. japonicum* is the symbiotic microorganism which forms root nodules on the roots of soybean to fix atmospheric nitrogen. The seeds of soybean were coated with *R. japonicum* to prepare legume inoculants. These seeds were referred as treated seeds. The treated seeds and untreated seeds were sown in the field. After 15 days of sowing of treated seeds and untreated seeds, the nodulation was checked. It was noted that treated seeds shown more nodules as compared to untreated seeds. The investigation was again carried out to check the effect of *R. japonicum* in relation to chlorophyll content. The plants arrived from treated seeds shown high chlorophyll content as compared to the plants arrived from untreated seeds.

Key words : *Glycine max*, *Rhizobium japonicum*, Legume inoculants, Nodulation, Chlorophyll content.

How to cite this paper : Kamthane, D.C. (2012). Effect of *Rhizobium japonicum* in relation to nodulation and chlorophyll content of soybean. *Asian J. Bio. Sci.*, 7 (1) : 59 - 61.

INTRODUCTION

The *Rhizobium* culture strains are antigenically very selective and require particular host for nodulation. The surface antigen on the rhizobial cells recognizes the binding sites (specific root exudates) on the roots of the leguminous plants. This characteristic makes them host-specific. Specific rhizobial cell can penetrate the roots of the specific leguminous plants only and form nodules. They multiply within the nodule using the carbon source from the plant and in turn fix part of the atmospheric nitrogen to the plant.

Leguminous plants such as soybean (*Glycine max*) form root nodules that contains population of *Rhizobium* a nitrogen fixing bacterium. The symbiotic association between the plant and the *Rhizobium* is initiated when bacteria in the soil attach to root hairs. This highly specific attachment process is mediated by plant proteins, the lectins, that bind the bacteria to the surface of root hairs; that is then penetrated by the microbes. The bacteria from the root hairs travel to inside and infect the root cells. The infected root cells divide and form a nitrogen fixing root nodule. The nodule provides the anaerobic environment necessary for nitrogen fixation (Aneja, 1996).

There is increase in the shoot length, root length, shoot dry weight, root dry weight, nodule number, chlorophyll-a,

chlorophyll-b and total chlorophyll content of soybean. plants arrived due to *R. japonicum* treated seeds as compared to untreated seeds. The *R. japonicum* strains could significantly improve the plant growth. This is due to that *R. japonicum* produces various growth promoting substances, fix atmospheric nitrogen symbiotically with the help of soybean plant. The *Rhizobium* spp. are being used as biofertilizers to reduce the need for expensive chemical fertilizers. About 10 to 15 per cent increase of crop yield can be achieved with the use of this culture.

RESEARCH METHODOLOGY

Isolation of *Rhizobium japonicum*:

R. japonicum was isolated from fresh, healthy, unbroken and pink nodules from soybean (*Glycine max*) roots on sterile yeast extract mannitol agar (YEMA) medium using pour plate technique. The incubation was at 26°C for four days. After incubation, large gummy colonies appear on sterile YEMA plates (Kamthane, 2010).

After isolation, the seeds were inoculated with *R. japonicum*.

Seeds treatment with *R. japonicum*:

This was done by mixing isolated rhizobia culture in 10