



RESEARCH PAPER

Antifungal activity of brown, red and green alga seaweed extracts against *Macrophomina phaseolina* (Tassi) Goid., in pigeonpea var. CO (Rg) 7

S. AMBIKA* AND K. SUJATHA

Department of Seed Science and Technology, Agricultural College and Research Institute, MADURAI (T.N.) INDIA

(Email : ambikasingaram@gmail.com)

Abstract : *In vitro* studies was conducted to evaluate the effect of seaweed extracts of *Caulerpa racemosa* (green alga), *Sargassum myricocystum* (brown alga) and *Gracilaria edulis* (red alga) against the mycelial growth of *Macrophomina phaseolina* at different concentrations of 10, 15, 20, 25 and 30 per cent along with control by poison food technique. The result revealed that extract of *S. myricocystum* showed significant antifungal activity against pathogen followed by *G. edulis* and *C. racemosa*. *S. myricocystum* (30%) extract recorded the lowest mycelial growth (45.2, 50.6, 58.4 and 61.5 mm) at 24, 48, 72 and 96 hrs after incubation. Among the antagonists tested against *Macrophomina phaseolina*, the fungal antagonists *Trichoderma viride* was found to be most effective in reducing the mycelial growth than the bacterial antagonist *Pseudomonas fluorescens*. Both the antagonistic of fungi and bacteria has compatability with seaweed extracts in all the concentrations.

Key Words : Seaweeds, Soil borne pathogen, Red gram, *Macrophomina phaseolina*

View Point Article : Ambika, S. and Sujatha, K. (2015). Antifungal activity of brown, red and green alga seaweed extracts against *Macrophomina phaseolina* (Tassi) Goid., in pigeonpea var. CO (Rg) 7. *Internat. J. agric. Sci.*, **11** (2) : 210-216.

Article History : Received : 01.01.2015; Revised : 01.05.2015; Accepted : 15.05.2015