



Effect of bio-regulators on the yield of greengram

K. ANANTHI AND M. GOMATHY

ABSTRACT

An experiment was conducted at Millet Breeding Station, TNAU, Coimbatore during *Kharif* (July, 2006 - October, 2006) in CO 6 greengram with nine different treatments. The yield and yield components were significantly increased by the application of chemicals and bioregulators. Various yield components such as days to 50 per cent flowering, days to maturity, number of clusters per plant, number of flowers per plant, number of pods per cluster, number of pods per plant, fertility co-efficient, number of seeds per pod, pod weight per plant, seed weight per plant, seed - pod ratio, pod length, hundred seed weight, grain yield per plot, grain yield per hectare and harvest index were favourably enhanced by the foliar spray of 0.1% humic acid with 0.1 ppm brassinosteroid treatment.

KEY WORDS : Bio regulators, Greengram, Yield

Ananthi, K. and Gomathy, M. (2011). Effect of bio-regulators on the yield of greengram, *Internat. J. Forestry & Crop Improv.*, 2 (1) : 12-15.

INTRODUCTION

Pulses occupy a strategic position in intensive as well as subsistence agriculture, as they are excellent source of dietary protein for millions of people, nutritious feed for livestock and a mini nitrogen plant having profound ameliorative effect on the soil. Green gram is extensively grown in all types of soil under varying climatic conditions. Greengram is considered as the most whole some among pulses, free from heaviness and flatulence. The green pods are used as vegetable and haulms are used as fodder. On an average, the biomass has 1.5 % N on fresh weight basis, serving as a good source of animal feed. Flower and pod shedding is a common problem in greengram. There is also a possibility to overcome these constraints by foliar application of bio-regulators at the pre-flowering stage, which is one of the latest trends in agriculture. These bioregulators includes humic acid (HA), naphthalene acetic acid (NAA), benzyl adenine (BA), salicylic acid (SA), brassinosteroid (BR) etc. These plant growth regulators (PGRs) in general, help to increase the number of flowers

on the plant when applied at the time of flowering. The flower and pod drop may be reduced to some extent by spraying various growth regulators on foliage (Ramesh and Thirumuguran, 2001). The foliar application of PGRs and urea significantly increased seed yield per plant (Patil *et al.*, 2005). The HA increased the yield of plants more especially, with insufficient nutrient supply than with optimum nutrient supply (Pagel, 1960). The NAA is one of the important synthetic, stable and also widely used form of auxins. Auxins had been very effective in promoting growth and development in pulse crops. Upadhaya (1994) reported that foliar spray of NAA at bud initiation and pod formation stage of chick pea increased plant height, number of branches, number of flower bud, number of flowers and yield. The main role of NAA rests with the efficient transport of sugars from the photosynthesizing parts of the plant (source) to the developing grains (sinks) and also facilitating nitrogen accumulation that probably result in higher total dry matter production. Foliar spray of 25 ppm NAA recorded significantly higher seed yield by 21 to 22 per cent than control through increased flower production, clusters per plant, pod setting percentage and pods per plant in mungbean (Patil *et al.*, 2005). Cytokinins are adenine derivatives characterized by an ability to induce cell division in the presence of auxins. Cytokinin improves crop growth and yield through redirecting the metabolic balance of growth, delayed leaf senescence, increased photosynthetic CO₂ uptake and partitioning of assimilates (Brault and

Correspondence to:

K. ANANTHI, Department of Crop Physiology, Tamil Nadu Agricultural University, COIMBATORE (T.N.) INDIA
Email : ananthipad@yahoo.com

Authors' affiliations:

M. GOMATHY, Department of Crop Physiology, Tamil Nadu Agricultural University, COIMBATORE (T.N.) INDIA