

Research Paper :

Effect of *Pseudomonas fluorescens* Pf1 formulation application on rhizosphere and phyllosphere population in groundnut



B. MEENA

International Journal of Plant Protection, Vol. 4 No. 1 (April, 2011) : 92-94

SUMMARY

Late leaf spot caused by *Cercosporidium personatum* is the destructive foliar disease in groundnut. The effect of seed treatment and foliar application of *Pseudomonas fluorescens* Pf1 formulation on groundnut rhizosphere and phyllosphere population was studied. The results revealed that when Pf1 formulation treated seeds were sown, the rhizosphere population of *P. fluorescens* was increased. The rhizosphere population of *P. fluorescens* increased with increase in age of the crop. *P. fluorescens* population was also found to be increased in phyllosphere due to foliar application of Pf1 formulation.

Correspondence to :
B. MEENA
Sugarcane Research
Station (T.N.A.U.),
Sirugamani, TRICHY
(T.N.) INDIA

Meena, B. (2011). Effect of *Pseudomonas fluorescens* Pf1 formulation application on rhizosphere and phyllosphere population in groundnut. *Internat. J. Pl. Protec.*, 4(1): 92-94.

Key words :
Pseudomonas
fluorescens,
Groundnut,
Rhizosphere,
Phyllosphere

The late leaf spot disease of groundnut, caused by the fungus, *Cercosporidium personatum* (Berk. and Curt.) Deighton is almost co-existent with the crop and contributes to significant loss in yield throughout the world (Wells *et al.*, 1994). Leaf spots damage the plant by reducing available photosynthetic area by lesion formation and by stimulating leaflet abscission. Smith *et al.* (1992) reported pod loss of 10 to 50 per cent by late leaf spot disease. Fluorescent *Pseudomonads* have emerged as the largest and potentially most promising group of plant growth promoting rhizobacteria (PGPR) for biocontrol of plant diseases (Kloepper and Schroth, 1978). Among the various biocontrol agents, fluorescent *Pseudomonads* are known to survive both in rhizosphere (Parks *et al.*, 1991) and phyllosphere (Wilson *et al.*, 1992). The present study was undertaken to find out the effect of seed treatment and foliar application of *P. fluorescens* Pf1 formulation on rhizosphere and phyllosphere population in groundnut.

MATERIALS AND METHODS

Groundnut seeds cv. (TMV 7) were treated with Pf1 formulation at different dosages viz., 10, 20, 50, 100 and 125 g kg⁻¹ of

seed and sown in pots containing field soil at the rate of 5 seeds per pot and 5 pots for each replication. As a check, seeds were treated with carbendazim at the rate of 2 g kg⁻¹ of seeds. At 45 days after sowing, the plants were inoculated with the pathogen. The trial was laid out in a randomized block design with six replications. The rhizosphere population of *P. fluorescens* was enumerated at 15 days interval by the method of Papavizas and Davey (1961).

Pf1 talc-based formulation at different concentrations viz., 0.5, 1.0, 1.25, 1.5 and 2.0 kg ha⁻¹ was dissolved in water, allowed to settle for 2 h, filtered through muslin cloth and the filtrate was sprayed on to groundnut plants on 30, 45, 60, 75 and 90 DAS. As a check, the plants were sprayed with carbendazim at the rate of 500 g ha⁻¹ and mancozeb 1 kg ha⁻¹. The plants were inoculated with pathogen at 45 DAS. The population of *P. fluorescens* was assessed from 1 g of leaf samples. The leaf samples without surface sterilization were transferred to a test tube containing 10 ml of sterile water and shaken well for 30 minutes. Then serial dilutions were made and the population of *P. fluorescens* was estimated using KB medium.

Received :
October, 2010
Accepted :
December, 2010