

## Botanical management of *Rhizoctonia solani* in soybean

■ S.P. PATOLE\*, R.S. SALUNKHE AND A.D. PHAPALE

Department of Plant Pathology, Mahatma Phule Krishi Vidyapeeth, Rahuri, AHMEDNAGAR (M.S.) INDIA

### ARTICLE INFO

**Received** : 24.07.2016

**Revised** : 07.09.2016

**Accepted** : 21.09.2016

### KEY WORDS :

Soybean, *Rhizoctonia solani*, Botanical

### ABSTRACT

Soybean is one of the important most pulse and oilseed crops grown all over the world. The root rot disease of soybean caused by *Rhizoctonia solani* is generally found in the soybean growing area in Maharashtra state. The present investigation was undertaken to study effects of botanical extract on root rot of soybean under glass house condition from pot culture experiment in glass house, it was found that the minimum PDI was observed in seed treatment with botanical extracts viz., *Zingiber officinale* L. (31.48) followed by *Allium sativum* L. (35.18) and *Azadirachta indica* (40.18)

**How to view point the article** : Patole, S.P., Salunkhe, R.S. and Phapale, A.D. (2016). Botanical management of *Rhizoctonia solani* in soybean. *Internat. J. Plant Protec.*, 9(2) : 561-565, DOI : 10.15740/HAS/IJPP/9.2/561-565.

\*Corresponding author:

Email : [santoshpatole57@rediffmail.com](mailto:santoshpatole57@rediffmail.com)