

## Standardisation of propagation methods for JACK (*Artocarpus heterophyllus*, Lam)

A. BASKARAN, S. SARASWATHY AND S. PARTHIBAN

Accepted : October, 2008

See end of the article for authors' affiliations

Correspondence to:

**A. BASKARAN**

Department of Fruit Science  
Horticulture College and  
Research Institute,  
PERIYAKULAM (E.) (T.N.)  
INDIA

### ABSTRACT

Clonal propagation is the common and fast method employing in fruit crops for production of elite grafts by mass multiplication. An experiment was conducted to standardize propagation methods for jack (*Artocarpus heterophyllus*) under 50 per cent shade (shade net) condition at Horticulture College and Research Institute, Periyakulam. Among the different types of propagation methods viz., inarching, epicotyl and soft wood grafting tested, soft wood grafting performed at different age groups of rootstock (*i.e.*, two, three and four months old seedlings). The studies revealed that in general softwood grafting performed well compared to other methods of propagation. In particular soft wood grafting on two months old rootstock recorded least number of days taken to graft union and the highest percentage of sprouting and graft take, length of shoots, girth of rootstock, girth of scion and number of leaves followed by soft wood grafting on three and four months age old rootstocks.

**Key words :** Jack, Clonal propagation, Root stock, Scion, Inarch grafting, Epicotyl grafting, Soft wood grafting.

Jack (*Artocarpus heterophyllus*, Lam) is an indigenous fruit, belonging to the family Moraceae is very popular among the rural areas of India. It is an ever green tree growing well in areas of receiving annual rainfall more than 1,200 mm (Mitra, 1998) and is widely grown in eastern and southern regions of India. The agro-climatic conditions prevailing in low hills and plain areas of India as a whole is highly propitious and offer splendid scope for successful cultivation of jack. In south India, the jack is a popular food ranking next to the mango and banana in total annual production.

Among the various constraints for expanding the jack fruit cultivation, lack of availability of suitable clonal planting materials is one of the impediments to expand the area of cultivation. Though, seed propagation is the most common method of propagation, which will not produce true-to-type progenies, besides having longer gestation period. Hence, standardization of suitable vegetative propagation technique is a prerequisite for successful cultivation. Though, inarching and air layering is in practice, these methods have some limitations like cumbersome and multiplication rate is low.

### MATERIALS AND METHODS

A study was conducted on standardization of propagation methods for jack (*Artocarpus heterophyllus*, Lam) at Horticulture College and Research Institute, Department of Fruit Crops, Periyakulam. The location of experiment is situated at 10° N latitude and 77.8° E longitude with an elevation of about 300 m above

MSL. The average rainfall is 750 mm per year.

Healthy jack seeds were collected from local and were utilized for raising seedling rootstocks. Soil, sand and well decomposed FYM in the composition of 2:1:1 ratio was used as the rooting medium. The rooting medium was thoroughly mixed. It was filled in polythene bags of 200 gauge thickness, 20 cm length and 10 cm width. The polythene bags were provided with number of vents for proper drainage. Soft and well matured terminal shoots of 10 cm length were selected with same thickness of the seedling rootstock. The experiment was carried out by Randomized Block Design (RBD) with four replications.

The experiment comprised of five treatments performed with different methods of propagation – T<sub>1</sub> - Inarch grafting, T<sub>2</sub> - Epicotyl grafting (20-25 days old), T<sub>3</sub> - Softwood grafting (2 months old), T<sub>4</sub> - Softwood grafting (3 months old), T<sub>5</sub> - Softwood grafting (4 months old).

The grafted plants were maintained under shade net (50% shade).

### RESULTS AND DISCUSSION

Days taken for the union of stock and scion were found to be significantly influenced by the age of the rootstock and the maturity of the scion. Among the different ages of the rootstock, two months old rootstock of softwood grafting recorded the least number of days (20.6) for graft union followed by three and four months old rootstocks where as inarch grafting took the maximum number of days for union.