

## Studies in the biomass production of papaya varieties

P.N. KATIYAR, J.P. SINGH, P.C. SINGH AND KUVAR SINGH

Accepted : November, 2008

See end of the article for authors' affiliations

Correspondence to:

**P.N. KATIYAR**

Department of  
Horticulture, Chandra  
Shekhar Azad University  
of Agriculture and  
Technology, KANPUR  
(U.P.) INDIA

### ABSTRACT

Papaya, being dioecious in nature, produces plants having male and female flowers separately on different plants. Cross pollination always occur in the crop. As a result of this phenomenon, there is wide diversity in varietal behaviour of papaya cultivars. With a view to identify the suitability of such standard variety of papaya for cultivation under the agroclimatic conditions of the Gangetic plains of the country, a stage wise study on screening of suitable dwarf varieties of papaya was undertaken in Horticultural garden of C.S.A. Univ. of Agric. and Tech., Kanpur during 2006-07. Seven varieties of papaya namely Surya, HG-05, Pant-1, Pusa Delicious, Selection S-1, Honey Dew and Pusa Nanha, were included in the studies. The maximum germination and survival per cent of papaya seedlings in the field was recorded in Pusa Nanha, while HG-05 variety proved most vigorous and produced the maximum biomass. The other varieties, included in the present study, remained intermediary with regard to biomass production.

**Key words :** Papaya, Varieties, Biomass production.

The Papaya (*Carica papaya* L.) is an important fruit of tropical and subtropical region of the world. It is native of tropical America and was introduced in India in 16<sup>th</sup> Century. In India, though it is grown all over the country but important papaya growing states are Karnataka, West Bangal, Assam, Kerla, Andhra Pradesh, Tamilnadu, Bihar, U.P and Maharashtra. In northern India, the occurrence of low temperature and frost restricts its cultivation. Papaya is an evergreen plant and bears flowers and fruits almost in all part of the year and hence, it requires a soil of high fertility and good drainage. Papaya being dioecious in nature produces plants having male and female flowers separately on different plants. Cross pollination always occur in the crop. As a result of these practices, there is wide diversity in varietal behaviour of papaya cultivars. However, some good varieties of papaya have been evolved and the purity of seed of these varieties have been maintained under ideal conditions of production.

With a view to identify the suitability of such standard variety of papaya for cultivation under the agro climatic conditions of central U.P., a stage wise experiment on screening of suitable dwarf varieties of papaya was undertaken.

### MATERIALS AND METHODS

The present investigation aimed for suitable dwarf varieties of papaya (*Carica papaya* L.) was undertaken mostly dealing with vegetative growth under the title "Studies in the biomass production in papaya cultivars". Experiment was conducted in the experimental block of

garden of the Department of Horticulture, Chandra Shaker Azad University of Agriculture and Technology Kanpur, U.P. during the year 2006-07.

The Investigation was undertaken with seven varieties of Papaya namely Surya, HG-05, Pusa Delicious, Pant-1, Honey Dew, Selection-1 (S-1) and Pusa Nanha under randomized block design. Observations were recorded on germination and survival percentage of papaya seedlings and growth parameters viz. height of plant (cm), girth of stem (cm), number of leaves per plant, length of leaves (cm), width of leaves (cm), plant spread North- South and East- West (cm), and biomass production of plant (green weight) in kg at monthly interval. Final observations were recorded 7 months after transplanting.

### RESULTS AND DISCUSSION

Germination of papaya seed in the present investigation varied significantly with the varieties, Pusa Nanha gave the maximum of 63.33% germination followed by Honey Dew and Pant-1 varieties. Similarly significantly highest percentage of survival was noted under Pusa Nanha (88.00) closely followed by Pusa Delicious (87.33) and Pant-1 (86.66) as compared to all others. Surya variety, however, showed the poorest germination of 34% followed by HG-05 exhibiting 41.00% germination (Table 1). The variation expressed by papaya varieties may be attributed to the varietal behaviour. Similar results were reported by Naik (1949) in papaya.

The vegetative growth like height which is one of the most important parameters of judging growth of plant