

Research Note

THE ASIAN JOURNAL OF HORTICULTURE

Vol. 6 | Issue 2 | December, 2011 | 544-545



Article history:

Received : 01.07.2011

Accepted : 12.11.2011

Effect of nitrogen and vermicompost on floral and yield parameters on marigold

■ P.R. PATEL¹, N.K. PATEL¹, R.Z. VALIA¹ AND S.R. CHAUDHARI

Associated Authors:

¹ASPEE College of Horticulture and Forestry, Navsari Agricultural University, NAVSARI (GUJARAT) INDIA

Author for correspondence :

S.R. CHAUDHARY

ASPEE College of Horticulture and Forestry, Navsari Agricultural University, NAVSARI (GUJARAT) INDIA

Abstract : An experiment was conducted on marigold cv. SIERRA YELLOW to evaluate the effect of nitrogen along with vermicompost on floral and yield parameters. The various levels of nitrogen (0, 120, 160, 200, 240 kg/ ha) and vermicompost (0, 5, 10, 15 t/ha) studied under FRBD with three replications. Application of 160 kg N per ha with 10 t/ha vermicompost minimize days to 50 % flowering (64.25), increase the number of flowers (58.38), flower yield (203.42 g/plant and 8793.60 kg/ha), flower diameter (7.60 cm), and keeping quality (9.28 days) of marigold cv. SIERRA YELLOW.

Key words : Broccoli, Date of planting, Spacing, Yield

How to cite this article : Patel, P.R., Patel, N.K., Valia, R.Z. and Chaudhari, S.R. (2011). Effect of nitrogen and vermicompost on floral and yield parameters on marigold, *Asian J. Hort.*, 6 (2) : 544-545.

African marigold (*Tagetes erecta* Linn.) is commonly used for the loose flower in India because of easy cultivation, adaptability to varying soil and climatic conditions, long duration of flowering and keeping quality. It has been established that nutrition play an important role in improvement of flower and yield in marigold. Nitrogen is absorbed by plants in huge amount and it is most limiting factor for crop production. With application of vermicompost it applies nitrogen phosphorus and other micro nutrient in trace quantity. Under such conditions balanced nitrogen and other nutrients are highly imperative to obtain higher yield of marigold.

Present investigation was carried out at college farm, N.M. College of Agriculture, Navsari Agricultural University Navsari during winter 2005-06 to standardize the dose of nitrogen along with vermicompost in African marigold cv. SIERRA YELLOW. Total 20 treatments combinations comprising five levels of nitrogen (0, 120, 160, 200, 240 kg/ ha) and four levels of vermicompost (0, 5, 10, 15 t/ha) were tried in Factorial Randomized Block Design with three replications in black soil. In the initial stage soil has available nitrogen (160.00 kg/ha), available Phosphorus (40.02 kg/ha), available potash (384.50 kg/ha) with pH (7.7). One month old seedlings of uniform

growth were transplanted at the spacing of 40x30 cm. The observations taken in the experiment were days taken to 50 % flowering, flower yield per plant and hectare basis, no of flower per plant, flower diameter, keeping quality etc.

The results obtained from the present investigation are summarized below :

Days taken for 50 % flowering:

Data present in Table 1 clearly reveals that significantly minimum (64.25) days taken for 50 % after transplanting were recorded in N₂ (160 kg N/ ha). Among vermicompost treatment V₂ has shows minimum (66.20) days for 50 % flowering. This might be due to vigorous vegetative growth of plant, which resulted in better food assimilation so increase in carbohydrates levels. This was translocated rapidly to flower bud initiation resulting early development. This was in accordance with the result obtained by Dahiya *et al.* (1998) in marigold.

Number of flower per plant:

Data in Table 1 clearly indicated that the significantly maximum number of flower per plant (58.38) in N₂ (160 kg/ha) treatment. Among vermicompost doses 10 t/ha