



Canopy temperature (CT), stress degree days (SDD) as influenced by treatments and varieties in soybean

S.R. PATIL*, N.R. SATPUTE¹ AND M.G. JADHAV

Department of Agrometeorology, Marathwada Krishi Vidyapeeth, PARBHANI (M.S.) INDIA

Abstract : The experiment was laid out in Split Plot Design with three replications and two factors viz., date of sowing D₁ (MW-27), D₂ (MW-28), D₃ (MW-29) and D₄ (MW-30) and cultivars V₁ (MAUS-47), V₂ (MAUS-71), V₃ (MAUS-81), V₄ (MAUS-158), V₅ (JS-9305) and V₆ (JS-335) to find out the optimum sowing time for soybean genotypes. Experiment was carried out at research farm of Department of Agricultural Meteorology, Parbhani The canopy temperature designates the plant water stress. If the canopy temperature of soybean crop is greater, then soil moisture stress occurred in the field. Canopy temperature is one of the most reliable indicators of the crop water stress due to its direct relation with the plant water status. The highest mean canopy temperature (32.0°C) and (32.1°C) were observed in D₄ (MW-30) date of sowing and genotype V₁ (MAUS-47), respectively whereas stage P₁₀ (maturity stage) indicated the highest mean canopy temperature 32.4°C. The lowest mean canopy temperature (30.9°C) and (30.7°C) were recorded in D₁ (MW-27) date of sowing and genotype V₄ (MAUS-158), respectively. Whereas stage P₁ (emergence stage) indicated the lowest mean canopy temperature i.e. 30.30°C. The variety growth characters like emergence and final plant count, plant height, number of functional leaves, number of branches, number of pods, mean leaf area, leaf area index, dry matter, weight of pods per plant, weight of grain per plant, 1000 seed weight (test weight), grain yield, straw yield and biological yield were observed maximum in D₁ (MW-27) date of sowing and in cultivar V₄ (MAUS-158). Whereas, minimum observed in D₄ (MW-30) date of sowing and cultivar V₁ (MAUS-47).

Key Words : Canopy temperature, Stress degree days, Varieties, Soybean

View Point Article : Patil, S.R., Satpute, N.R. and Jadhav, M.G. (2014). Canopy temperature (CT), stress degree days (SDD) as influenced by treatments and varieties in soybean. *Internat. J. agric. Sci.*, **10** (2): 587-591.

Article History : Received : 17.10.2013; Revised : 03.04.2014; Accepted : 18.04.2014

INTRODUCTION

In Maharashtra mainly the cultivation of soybean is as rainfed crop. It is the common experience of farmer and scientist that the rainfall at the maturity leads to the loss of seeds besides deterioration of seed quality. Thus, it is observed that harvesting period of soybean is coupled with post monsoon rains resulting in the loss of seed and its quality. In the past years, it has been observed that germination of seeds was considerably reduced to 11 to 20 per cent (Anonymous, 2000). This has alarmed for finding proper time for sowing so as to skip the crops from rains at

harvest. Performance of the crop has been reported to be highly governed by timely sowing and spatial arrangement (Sharma *et al.*, 1984). Delayed sowing of soybean not only resulted in yield reduction (Karmarkar and Bhatnagar, 1995) but also deteriorating the quality in respect to oil and protein content (Billare *et al.*, 2000). The temperature is an important meteorological variables that affect plant growth and development (Londe and Woodward, 1988). Day light or bright sunshine hours play an important role in growth and development of soybean crop. Same varieties flower in less than 30 days after emergence if exposed to day light less than twelve hours (Beard and Knowles, 1973).

* Author for correspondence

¹Zonal Agriculture Research Station, Krishak Bhavan, SOLAPUR (M.S.) INDIA

MATERIAL AND METHODS

The present investigation was carried out by laying out experiment on soybean with objective to study the performance of soybean (*Glycine max* L.) genotypes under varied weather conditions at Parbhani, Maharashtra. The experiment was conducted during *Kharif* season of 2010-2011, with Split Plot Design, Three replications and 24 treatments [Main treatment (sowing dates 4) × Sub treatment (Cultivar 6)].

RESULTS AND DISCUSSION

The data collected during the investigation were analyzed by using appropriate statistical methods.

Mean leaf area per plant (dm²) :

The data on mean leaf area (dm²) per plant as influenced by different treatments at 15 days interval are presented in Table 1.

Date of sowing :

The data on mean leaf area (dm²) per plant was influenced significantly by different date of sowing, at all stages of crop growth. Mean leaf area was observed significantly more in D₁ (MW-27) than other treatments.

Cultivars :

The mean leaf area was influenced significantly by different cultivars at all stages of crop growth. Mean leaf area was observed significantly more in D₁ (MW-27) than other treatments.

Interaction (D × V):

The interaction effect between date of sowing and different cultivars was found to be non-significant at all stages.

Dry matter per plant (g/plant) :

The data on mean dry matter per plant (g/plant) as influenced by different treatments at 15 days of interval are presented in Table 1. It was observed that mean dry matter per plant (g/plant) increased continuously up to 75 DAS of crop but later it decreased.

Date of sowing :

The data presented in Table 1 indicate that the production was influenced significantly by different dates of sowing at all stages of crop growth. Dry matter was observed significantly more in D₁ (MW-27) than the other dates of sowing.

Cultivars :

The mean dry matter production was influenced

Table 1 : Mean leaf area (dm²) per plant, Mean dry matter (g/plant) and Mean leaf area index (LAI) of soybean as influenced by different treatments

Treatments	Mean leaf area (dm ²) per plant					Mean dry matter (g/plant)					Mean leaf area index (LAI)				
						Days after sowing									
	30	45	60	75	At harvest	30	45	60	75	At harvest	30	45	60	75	At harvest
D ₁ (MW-27)	12.15	16.72	23.24	32.45	20.84	7.66	13.15	18.90	27.14	21.71	0.54	0.74	1.03	1.44	0.92
D ₂ (MW-28)	10.84	14.22	21.66	28.58	19.30	6.93	12.21	17.15	26.56	20.12	0.48	0.63	0.96	1.27	0.85
D ₃ (MW-29)	11.62	15.71	22.40	30.27	19.98	7.56	12.88	17.69	27.11	20.40	0.51	0.69	0.99	1.34	0.88
D ₄ (MW-30)	9.85	13.42	21.02	27.33	19.21	5.71	11.30	16.60	25.50	18.89	0.43	0.59	0.93	1.21	0.85
S.E.±	0.05	0.03	0.07	0.04	0.11	0.06	0.04	0.07	0.08	0.10	--	--	--	--	--
C.D. (P=0.05)	0.16	0.09	0.19	0.12	0.33	0.17	0.13	0.21	0.23	0.30	--	--	--	--	--
Cultivars															
V ₁ (MAUS-47)	9.10	12.94	18.15	26.37	17.55	5.41	10.04	15.48	24.49	18.41	0.40	0.59	0.80	1.17	0.78
V ₂ (MAUS-71)	12.22	16.10	24.30	31.42	21.27	7.80	13.51	18.62	27.91	21.38	0.54	0.71	1.08	1.39	0.94
V ₃ (MAUS-81)	11.16	14.81	21.13	29.30	19.25	6.72	12.10	17.31	25.93	19.98	0.49	0.65	0.93	1.30	0.85
V ₄ (MAUS-158)	12.64	16.80	25.30	32.13	21.91	8.54	14.05	19.21	28.86	22.19	0.56	0.74	1.12	1.42	0.97
V ₅ (JS-93-05)	9.89	13.95	20.18	28.05	18.56	6.15	11.61	16.59	25.17	19.08	0.43	0.62	0.89	1.24	0.82
V ₆ (JS-335)	11.83	15.50	23.51	30.53	20.44	7.16	13.08	18.15	27.10	20.64	0.52	0.68	1.04	1.35	0.90
S.E.±	0.07	0.06	0.10	0.14	0.12	0.08	0.08	0.06	0.11	0.08	--	--	--	--	--
C.D. (P=0.05)	0.20	0.19	0.29	0.40	0.36	0.24	0.23	0.18	0.33	0.24	--	--	--	--	--
Interaction															
S.E.±	0.15	0.13	0.21	0.28	0.24	0.17	0.16	0.13	0.22	0.17	--	--	--	--	--
C.D. (P=0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	--	--	--	--	--
G. Mean	11.15	15.01	22.08	29.63	19.83	6.96	12.39	17.56	26.57	20.28	--	--	--	--	--

NS=Non-significant

significantly by different cultivars at all stages of crop growth. The cultivar V₄ (MAUS-158) produced higher dry matters than other cultivars.

Interaction (D × V):

The interaction effect between date of sowing and different cultivars was found to be non-significant at all stages.

Growth analysis :

Mean leaf area index :

The data on mean leaf area index (LAI) per plant as influenced by different treatments at 15 days interval are presented in Table 1.

Date of sowing :

The data on mean leaf area index (LAI) per plant was influenced significantly by different dates of sowing at all stages of crop growth. Mean leaf area was significantly more in D₁ (MW-27) than other treatments.

Cultivars :

The mean leaf area index was influenced significantly by different cultivars at all stages of crop growth. V₄ (MAUS-158) produced more leaf area than other cultivars.

Grain yield (kg/ha) :

The data regarding grain yield are presented in Table 2.

Date of sowing :

The data on grain yield Table 2 indicated that the crop sown in D₁ MW-27 (02-08 July) recorded higher grain yield (2876 kg/ha) and found significantly superior over other treatments whereas the lowest yield was recorded in treatment D₄ (23-29 July). The crop sown in second week of July recorded low seed yield, dry spell resulted in low germination of crop. Over all, this year the crop recorded highest yield due to ample soil moisture during crop growing period.

Cultivars :

Statistical analysis of soybean cultivars showed significant results. During this year, variety MAUS-158 (V₄) produced higher grain yield (2579 kg/ha) and was found significantly superior over remaining treatments. Whereas, the variety V₁ (MAUS-47) produced lowest grain yield (1870 kg/ha) (Table 2).

Interaction :

The interaction effect between date of sowing and different cultivars was found to be non-significant at all stages and the results to that effect are presented in Table 2.

Table 2 : Mean grain yield (kg/ha), straw yield (kg/ha), biological yield (kg/ha) and mean soil moisture (%) as influenced by different treatments

Treatments	Grain yield (kg/ha)	Straw yield (kg/ha)	Biological yield (kg/ha)	Mean soil moisture (%)					
				Days after sowing					
				15	30	45	60	75	At harvest
Date of sowing									
D ₁ (MW-27)	2876	4057	6933	32.55	33.37	37.96	35.15	32.25	31.51
D ₂ (MW-28)	2035	3167	5202	26.86	31.60	34.10	30.18	27.47	23.42
D ₃ (MW-29)	2304	3342	5648	30.32	32.45	35.92	33.97	30.18	27.47
D ₄ (MW-30)	1780	2853	4632	28.13	31.20	32.80	28.34	25.90	22.42
S.E. ±	31.50	7.22	32.51	0.81	0.03	0.018	0.04	0.01	0.03
C.D. (P=0.05)	94.12	20.10	96.98	2.41	0.10	0.05	0.12	0.03	0.08
Cultivars									
V ₁ (MAUS-47)	1870	2934	4802	27.42	31.95	34.27	30.90	29.05	25.72
V ₂ (MAUS-71)	2451	3579	6030	29.95	32.16	35.78	32.38	29.45	26.10
V ₃ (MAUS-81)	2182	3320	5502	29.80	32.15	34.80	32.06	29.20	25.95
V ₄ (MAUS-158)	2579	3697	6276	30.05	32.20	35.79	32.48	29.54	26.35
V ₅ (JS-93-05)	2051	3191	5243	29.72	32.14	34.68	31.83	29.16	25.81
V ₆ (JS-335)	2363	3406	5770	29.85	32.15	35.76	32.29	29.32	26.08
S.E. ±	42.30	9.18	44.20	0.02	0.03	0.03	0.04	0.03	0.03
C.D. (P=0.05)	126.42	27.51	131.95	0.05	0.10	0.10	0.12	0.09	0.08
Interaction (D × V)									
S.E. ±	84.60	18.16	88.47	--	--	--	--	--	--
C.D. (P=0.05)	NS	NS	NS	--	--	--	--	--	--
G. Mean	2249	3355	5604	29.41	32.12	35.34	31.99	29.28	26.00

NS=Non-significant

Soil moisture studies :

The data on soil moisture at different sowing dates of soybean crop from sowing to maturity are presented in Table 2. The data on soil moisture revealed that the soil moisture in D₁ (MW-27) sowing dates treatment was on an average more than rest of the treatments, while in D₂ (MW-28) and D₄ (MW-30) sowing date treatments, the soil moisture stress at early growth stages was noticed, due to the yield of soybean crop was affected. The differences in soil moisture of the genotypes were significant at all stages. The soil moisture was significant

at all stages. The soil moisture content showed consistent increasing from 15 DAS to 45 DAS and then continuously decreased upto harvest of crop. At 45 DAS, the mean soil moisture was 35.34 per cent while at harvest the mean soil moisture was 26.00 per cent (Table 2).

Canopy temperature (°C) :

Canopy temperature for soybean crop under different sowing dates and varieties from sowing to maturity are presented in Table 3.

Table 3 : Mean canopy temperature (°C) at different phenophases of soybean crop

Treatments	Phenophases										Mean
	P ₁	P ₂	P ₃	P ₄	P ₅	P ₆	P ₇	P ₈	P ₉	P ₁₀	
D ₁ (MW-27)	29.9	30.2	30.4	30.5	30.6	31.2	31.3	31.4	31.8	31.4	30.9
D ₂ (MW-28)	30.5	30.7	31.1	31.2	31.4	31.7	32.0	32.1	32.7	32.8	31.6
D ₃ (MW-29)	30.2	30.6	30.8	30.9	31.3	31.4	31.5	31.9	32.1	32.2	31.3
D ₄ (MW-30)	30.8	31.0	31.4	31.5	32.1	32.3	32.4	32.4	33.0	33.2	32.0
Cultivars											
V ₁ (MAUS-47)	31.1	31.3	31.6	31.8	31.9	32.4	32.4	32.7	33.0	33.1	32.10
V ₂ (MAUS-71)	29.9	30.2	30.4	30.5	31.0	31.3	31.4	31.5	32.1	32.2	31.1
V ₃ (MAUS-81)	30.5	30.8	31.2	31.3	31.7	31.9	32.1	32.2	32.5	32.6	31.7
V ₄ (MAUS-158)	29.6	29.9	30.3	30.1	30.8	30.7	30.9	31.2	31.7	32.0	30.7
V ₅ (JS-93-05)	30.8	31.1	31.4	31.6	31.6	32.1	32.3	32.4	32.8	33.0	31.9
V ₆ (JS-335)	30.2	30.5	30.7	30.6	31.3	31.6	31.7	31.9	32.2	32.3	31.3
Mean	30.3	30.6	30.9	31.0	31.4	31.6	31.8	31.9	32.4	32.4	31.4
P ₁ – Sowing to emergence			P ₂ – Emergence to seedling				P ₃ – Seedling to branching				
P ₄ – Branching to flowering			P ₅ – Flowering to pod formation				P ₆ – Pod formation to grain formation				
P ₇ – Grain formation to pod development			P ₈ – Pod development to pod containing full size				P ₉ – Pod containing full size to dough stage				
P ₁₀ – Dough stage to maturity											

Table 4 : Mean canopy air temperature difference (Tc-Ta) at different phenophases of soybean crop

Treatments	Growth stages										Mean
	P ₁	P ₂	P ₃	P ₄	P ₅	P ₆	P ₇	P ₈	P ₉	P ₁₀	
D ₁ (MW-27)	-3.3	-3.1	-2.8	-1.7	-1.3	-1.1	-1.2	-1.2	-0.8	-0.4	-1.7
D ₂ (MW-28)	-2.8	-2.7	-1.2	-1.1	-0.5	-0.8	-1.0	-1.1	-0.5	0.2	-1.2
D ₃ (MW-29)	-3.0	-2.9	-2.2	-1.5	-0.7	-1.0	-1.1	-1.2	-0.6	0.1	-1.4
D ₄ (MW-30)	-2.6	-2.5	-1.0	-0.9	-0.5	-0.5	-0.8	-1.0	-0.2	0.4	-1.0
Cultivars											
V ₁ (MAUS-47)	-2.2	-2.1	-1.0	-0.6	-0.3	-0.4	-0.5	-0.7	-0.1	0.5	-0.7
V ₂ (MAUS-71)	-3.4	-3.2	-2.3	-1.7	-1.0	-1.1	-1.4	-1.4	-0.9	-0.2	-1.7
V ₃ (MAUS-81)	-2.7	-2.6	-1.5	-1.1	-0.6	-0.8	-0.9	-1.0	-0.5	0.30	-1.2
V ₄ (MAUS-158)	-3.9	-3.7	-2.8	-2.1	-1.3	-1.2	-1.6	-1.7	-1.0	-0.4	-2.0
V ₅ (JS-93-05)	-2.5	-2.4	-1.2	-0.9	-0.4	-0.6	-0.6	-0.7	-0.2	0.4	-0.9
V ₆ (JS-335)	-3.0	-2.8	-2.0	-1.5	-0.8	-1.0	-1.1	-1.2	-0.7	-0.1	-1.4
Mean	-2.9	-2.8	-1.8	-1.3	-0.7	-0.8	-1.0	-1.1	-0.5	0.1	-1.30
P ₁ – Sowing to emergence			P ₂ – Emergence to seedling				P ₃ – Seedling to branching				
P ₄ – Branching to flowering			P ₅ – Flowering to pod formation				P ₆ – Pod formation to grain formation				
P ₇ – Grain formation to pod development			P ₈ – Pod development to pod containing full size				P ₉ – Pod containing full size to dough stage				
P ₁₀ – Dough stage to maturity											

Canopy temperature ($^{\circ}\text{C}$) at different phenophases :

Data of Table 3 indicate that there were significant differences in canopy temperature at each phenophase of different dates of sowing and different cultivars. The canopy temperature was higher under stressed conditions as compared to unstressed conditions throughout the crop growth period. As per the date of sowing and different cultivars, the highest mean canopy temperature (32.0°C) and (32.1°C) were observed in D_4 date of sowing and V_1 (MAUS-47) genotype, respectively. Whereas stage P_{10} (maturity stage) indicated the highest mean canopy temperature *i.e.* 32.4°C .

The canopy temperature designates the plant water stress. If the canopy temperature of soybean crop is greater, then soil moisture stress occurred in the field. Canopy temperature is one of the most reliable indicators of the crop water stress due to its direct relation with the plant water status. As per the date of sowing and variety, the mean lowest canopy temperature (30.9°C) and (30.7°C) recorded in D_1 date of sowing and genotype V_4 (MAUS-158), respectively (Table 3). Whereas P_1 (emergence stage) indicated the lowest mean canopy temperature *i.e.* 30.3°C . The data presented in Table 3 revealed that the average canopy temperature ranged from 30.3°C to 32.4°C in P_1 to P_{10} stage (emergence to maturity). The canopy temperature was less than air temperature because of the occurrence of rainfall in all stages except only P_{10} stage (maturity stage). While in maturity stage, canopy temperature was more than air temperature so there was moisture stress observed. Similar results were reported by Singh and Kanemasu (1983), Idso (1982) and Zhang Wen-Zhang *et al.* (2007).

Canopy-air temperature (T_c-T_a) difference during phenophases :

The data pertaining to canopy-air temperature differential (T_c-T_a) during crop growth period in all dates of sowing and different cultivars are presented in Table 4. The T_c-T_a values were recorded in similar fashion as that of canopy temperature in all dates of sowing and genotypes. The average T_c-T_a values ranged from -2.9°C to 0.1°C in P_1 to P_{10} stage.

In maturity stage (P_{10}) there was significant difference observed in all dates of sowing and genotypes. The P_{10} stage showed higher positive ($T_c - T_a$) difference (0.1°C), which showed the soil moisture stress in P_{10} stage. While in other stages, rainfall was occurred, so that T_c-T_a were negative which showed no any moisture stress in remaining stage except P_{10} stage (maturity stage). The highest (T_c-T_a)

difference was observed in P_{10} stage in all dates of sowing and genotypes. Whereas, the lowest ($T_c - T_a$) difference was observed in P_1 (emergence stage) in all dates of sowing and genotypes. The similar results were reported by Ajayi and Pandey (1983) and Idso (1982).

Conclusion :

It was found that the highest canopy temperature was observed in D_4 (MW-30) *i.e.* 32.0°C and cultivar V_1 (MAUS-47) *i.e.* 32.1°C due to bright sunshine and clear weather. Lowest canopy temperature was recorded in D_1 (MW-27) *i.e.* 30.9°C and cultivar V_4 (MAUS-158) *i.e.* 30.7°C . So, no any moisture stress observed in all phenophases stage except P_{10} (maturity stage). In P_{10} stage canopy temperature was more than air temperature so moisture stress was observed there.

REFERENCES

- Ajayi, A.E. and Pandey, E. (1983). Use of canopy temperature as an indicator for water use efficiency and yield productivity in wheat. *Saudi. J. Bio. Sci.*, **5**(1) : 57-70.
- Anonymous (2000). Epitome of agriculture in Maharashtra state published by Commission rate. Government of Maharashtra Dept. of Agricultrure.
- Beard, B.H. and Knowles, P.F. (1973). Soybean research in California Calif, *Agril. Exp. Sat. Bull.*, 862.
- Billare, S.D., Joshi, C.P. and Ramesh, A. (2000). Performance of soybean (*Glycin max L.*) genotypes on different sowing dates and row spacing in vertisols. *Indian J. Agric. Sci.*, **70**(9) : 577-580.
- Idso, S.B. (1982). Non-water stressed baseline : A key to measuring and interpreting plant water stress. *Agric. Forest Met.*, **27** : 59-70.
- Karmarkar, P.G. and Bhatnagar, P.S. (1995). Performance of soybean (*Glycine max L.*) at different dates of sowing in Mallwa plateau of Madhya Pradesh. *Indian J. Agric. Sci.*, **65**(2) : 138-139.
- Londe, S.P. and Woodward, F.I. (1988). Plants and temperature symposium of the society for experimental biology, No. 42.
- Sharma, S.C., Khalita, M.M. and Kakati, N.N. (1984). Effect of dates of planting of five soybean varieties. *Soybean Genetics, Newsletter*, **11** : 34-37.
- Singh, P. and Kanemasu, E.T. (1983). Leaf and canopy temperatures of pearl millet genotypes under irrigated and non-irrigated conditions. *Agron. J.*, **75** : 497-501.
- Zhang, Wen-Zhong, Han, Ya-Dong, Du and Hong Juan (2007). Relationship between canopy temperature at flowering stage and soil water content, yield components in rice. *Sci.*, **14**(1): 67-70.

10th
Year
★★★★★ of Excellence ★★★★★