

# Pests status of coconut in managed and unmanaged garden

■ S.M. Wankhede\*, V.V. Shinde and S.L. Ghavale

AICRP on Palms, Regional Coconut Research Station (D.B.B.S.K.V.V.), Bhatye, **Ratnagiri (M.S.) India**

## ARTICLE INFO

**Received** : 11.05.2019  
**Revised** : 25.08.2019  
**Accepted** : 11.09.2019

## KEY WORDS :

Black headed caterpillar coconut,  
Eriophyid mite, Rhinoceros beetle, Red  
palm weevil

## ABSTRACT

The major pests incidence and intensity on coconut was recorded during fixed plot survey. It was carried out at bimonthly interval from April 2018 to February 2019 at Regional Research Station, Bhatye, Ratnagiri jurisdiction. Two plots were selected for observations, one was well managed (good sanitation, use of fertilizes etc.) and another unmanaged (poor sanitation, no use of fertilizers etc.). The major pest status in managed garden was found minimum which recorded rhinoceros beetle (25.7%), eriophyid mite (41.7%) and mite grade index 0.71. Whereas, maximum infestation of rhinoceros beetle (*Oryctes rhinoceros* L.), red palm weevil (*Rhynchophorus ferrugineus* O.) and eriophyid mite (*Aceria guerreronis* Keifer) were observed 30.3, 0.3, 71.7 per cent, respectively and mite grade index was recorded 1.57 (moderate) in unmanaged garden. The average data of two fixed plots revealed that the maximum infestation (48.0%) was observed in June 2018 and minimum incidence was observed in February 2019 which recorded 7.64 per cent. The infestation of red palm weevil and black headed caterpillar (*Opisina arenosella* W.) were recorded 1.11 and 0.64 per cent only in February 2019. The highest infestation of eriophyid mite was recorded 64.9 per cent in April, 2018 and least incidence (41.6%) was observed in October, 2018. The mite damage grade index 1.39 (Moderate) was recorded in February 19. However, lowest (0.67) was observed in October, 2018.

**How to view point the article** : Wankhede, S.M. Shinde, V.V. and Ghavale, S.L. (2019). Pests status of coconut in managed and unmanaged garden. *Internat. J. Plant Protec.*, **12**(2) : 119-122, DOI : 10.15740/HAS/IJPP/12.2/119-122, Copyright@ 2019: Hind Agri-Horticultural Society.

\*Corresponding author:

Email : [agr\\_agronomist@yahoo.co.in](mailto:agr_agronomist@yahoo.co.in)

## INTRODUCTION

Coconut (*Cocos nucifera* Linn) is the most important versatile tree crop cultivated in the tropics providing livelihood and employment securities to the rural agrarian mass in the region. It is an important plantation crop which is being mainly cultivated by farmers of southern state and states like Goa, Maharashtra, Gujarat, West Bengal, Odisha, Asam, Andaman and Nicobar

Islands, Lakshdweep and Minicoy islands are also having considerable area under coconut. Coconut is essentially a crop of marginal and small farmers in India. The low size of the land holding by coconut growers is a serious challenge to both profitability and sustainability of the crop. The coconut palm is attacked by a variety of pests. The majority attack the leaves, while others attack the trunks (stems), inflorescences, young nuts and roots. Some are major pests which cause considerable damage

and yield loss, while some are less damaging. Most of these insects remain as minor pests, either because the environmental conditions are not favorable for their rapid multiplication or because they are adequately controlled by natural enemies. Coconut eriophyid mite *Aceria guerreronis* Keifer was first reported as a serious pest in Kerala during 1997-98. Subsequently, the devastating effects of these mites were noticed in Coimbatore and Theni districts of Tamil Nadu and Bangalore in Karnataka (Sathiamma *et al.*, 1998 and Mohanasundaram *et al.*, 1999). Mathen and Kurian (1962) reported that the red palm weevil, *Rhynchophorus ferrugineus* Oliv, (RPW) (Curculionidae, Coleoptera) is a serious pest attacking different species of palm trees (e.g. date palm, coconut palm and royal palm). Chalapathi Rao *et al.* (2018) found that per cent of leaf damage due to rhinoceros beetles ranged from 12.5 to 35.5 and spindle damage ranged from 33.3 per cent to 45 per cent. By considering economic importance of pest and yield loss of palm, the present investigation was carried out for assessment of major pests in fixed plots during survey were undertaken with view to know the exact status of coconut pests in Ratnagiri district of Maharashtra.

## MATERIAL AND METHODS

A field survey was carried out as Entomological Experiment under 'Pest surveillance of coconut at (AICRP (Palms), RCRS, Bhatye Dist. Ratnagiri (M.S.) during 2018-19 for knowing of exact pests status on coconut. Two different plots at different location were selected with well managed (Good sanitation, use of fertilizers etc.) and unmanaged (poor sanitation, No use of fertilizers). One plot has need based management treatment for major pests and other plot with no management aspect (natural control). Six observations per year at 2 months interval *viz.*, April, June, Aug., Oct.,

Dec. and February. Major pests like rhinoceros beetle, red palm weevil, black headed caterpillar and eriophyid mite were recorded by using following methods:

### Rhinoceros beetle:

Per cent of palm infested (Out of 100 palms/garden)- The top 10 fronds in each palm with single/ multiple cuts was observed. Per cent leaf damage (25 palms at random/garden) (infested/total number of leaves x 100).

### Eriophyid mite:

Per cent nut infested (Mite infested nuts/total nuts) (25 palms at random/garden). Damage grade (mature bunch) (intensity 0-4 scale) (out of minimum 100 nuts/garden) using CPCRI scales were followed.

### Red palm weevil:

Per cent of palms infested (out of total palm (>100)/garden with typical symptom of RPW).

### Black headed caterpillar:

Per cent leaf damage (infested leaf/total leaf) (50 palms/ garden) were used.

Per cent damage on nut surface	Scale	Grade index	Intensity
Nuts with no mite damage	0	0	Nil
< 25%	1	0.1-1.0	Mild
25-50%	2	1.1-2.0	Moderate
50-75%	3	2.1-3.0	High
>75%	4	3.1-4.0	Severe

## RESULTS AND DISCUSSION

The data depicted in Table 1 indicated that minimum per cent incidence of major pest status *viz.*, rhinoceros

Table 1 : Extent of infestation by different pests in fixed plot (Managed) during survey in Maharashtra (Plot No. 1)

Month	Rhinoceros beetle			Red palm weevil incidence (%)	Black headed caterpillar incidence (%)	Eriophyid mite	
	Incidence (%)	Leaf damage	Spindle damage			Infestation (%)	MGI
April,18	24.0	2.8	0.0	0.0	0.0	56.5	1.2
June,18	52.0	7.0	0.0	0.0	0.0	53.1	0.9
Aug.,18	36.0	4.0	8.0	0.0	0.0	37.8	0.5
Oct., 18	28.0	3.6	0.0	0.0	0.0	35.8	0.6
Dec., 18	08.0	1.6	0.0	0.0	0.0	30.8	0.5
Feb., 19	06.4	2.0	1.2	0.0	1.2	36.2	0.5
Mean ± S.E.	25.7 ± 7.7	3.5 ± 0.8	1.5 ± 1.4	0.0 ± 0.0	0.2 ± 0.0	41.7 ± 4.6	0.7 ± 0.1

Table 2 : Extent of infestation by different pests in fixed plot (Unmanaged) during survey in Maharashtra (Plot No. 2)							
Month	Rhinoceros beetle			Red palm weevil incidence (%)	Black headed caterpillar incidence (%)	Eriophyid mite	
	Incidence (%)	Leaf damage	Spindle damage			Infestation (%)	MGI
April,18	25.0	6.0	0.0	0.0	0.0		
June,18	44.0	7.2	0.0	0.0	0.0	76.0	1.7
Aug.,18	48.0	6.0	4.0	0.0	0.0	72.0	1.3
Oct., 18	24.0	4.4	4.0	0.0	0.0	47.4	0.7
Dec., 18	32.0	5.2	0.0	0.0	0.0	78.8	1.7
Feb., 19	8.8	3.2	3.3	2.2	0.0	82.4	2.2
Mean $\pm$ S.E.	30.3 $\pm$ 6.4	5.3 $\pm$ 0.6	1.8 $\pm$ 0.9	0.3 $\pm$ 0.4	0.0 $\pm$ 0.0	71.7 $\pm$ 5.5	1.5 $\pm$ 0.2

Table 3: Average per cent infestation by different pests in fixed plot survey in Maharashtra							
Month	Rhinoceros beetle			Red palm weevil incidence (%)	Black headed caterpillar incidence (%)	Eriophyid mite	
	Incidence (%)	Leaf damage	Spindle damage			Infestation (%)	MGI
April,18	24.5	4.4	0.0	0.0	0.0	64.9	1.3
June,18	48.0	7.1	0.0	0.0	0.0	64.6	1.3
Aug.,18	42.0	5.0	6.0	0.0	0.0	54.9	0.9
Oct.,18	26.0	4.0	2.0	0.0	0.0	41.6	0.6
Dec., 18	20.0	3.4	0.0	0.0	0.0	54.8	1.1
Feb., 19	7.6	2.6	2.3	1.1	0.6	59.3	1.3
Mean $\pm$ S.E.	28.02 $\pm$ 5.44	4.41 $\pm$ 0.63	1.71 $\pm$ 1.16	0.18 $\pm$ 0.00	0.1 $\pm$ 0.0	56.7 $\pm$ 4.2	1.1 $\pm$ 0.1

beetle (25.73%), eriophyid mite (41.74%) and mite grade index 0.71 was noticed in managed garden. Whereas, maximum infestation of rhinoceros beetle, red palm weevil and eriophyid mite were observed 30.31, 0.37, 71.70 per cent, respectively with mite grade index 1.57 (moderate) in unmanaged garden (Table 2).

The average data of two fixed plots are presented in Table 3 revealed that the incidence was in the range of 20.0 to 48.0 per cent and maximum infestation of rhinoceros beetle was observed in the month of June 2018 which recorded 48.00 per cent and minimum incidence (20.0%) was observed in December, 2018. Chalapathi Rao *et al.* (2018) found that the per cent of leaf damage due to rhinoceros beetles ranged from 12.5 to 35.5 and spindle damage ranged from 33.3 per cent to 45 per cent. The infestation of red palm weevil and black headed caterpillar were recorded 1.1 and 0.6 per cent only in February 2019. The infestation of eriophyid mite was in the range of 41.6 to 64.9 per cent. Similar results on eriophyid mite infestation was also noticed by Alagar *et al.* (2019). Maximum infestation (64.9 %) was noticed in the month of April, 2018 and least incidence (41.6%) was observed in October, 2018. The mite damage grade index 1.37 (Moderate) was recorded in April, 18.

However, lowest (0.67) was observed in October, 2018. Present data correlated with Levin and Mammooty (2003) indicated that most of the infested nuts were in the damage category of two and three and the percentage of mite damage was only 25.4 per cent. Desai *et al.* (2009) indicated that the eriophyid mite infestation was higher in Thane district followed by Sindhudurg district. Gurav *et al.* (2018) observed that pest incidence and pest intensity trend showed that peak incidence was recorded during summer months that are from April to May while it decreases from September onwards and reaches to minimum in November and December. Muyengi *et al.* (2015) showed that about 46.7 per cent of the farmers experienced the problem of *Rhinoceros* beetle (*Oryctes monoceros*) in their farms and about 4.7 per cent problems with coconut mites (*Aceria guerreronis*). Vanderplank (1959); Bedford (1975); Paul (1985) and Seguni (2010) also indicated the same results.

#### Acknowledgement:

The authors are thankful to ICAR-AICRP (Palms) and ICAR-CPCRI, Kasaragod for providing funds, technical help and timely guidance for conduct of experiment.

## REFERENCES

- Alagar, M., Srinivasan, T. and Maheshwarappa, H.P. (2019).** Integrated management of coconut eriophyid mite, *Aceria guerreronis* Keifer (Eriophyidae: Acari). 23rd Plantation Crops Symposium, 6-8 March, 2019, Book Abstracts, ABS P6, 105.
- Bedford, O. (1975).** Trap catches of the coconut rhinoceros beetle, *Oryctes monoceros* (L.) (Coleoptera, Scarabaeidae, Dynastinae) in New Britain. *Bull. Entomological Research*, 65: 443-451.
- Chalapathi Rao, N.B.V., Snehalatharani, A., Nischala, A., Ramanandam, G. and Maheswarappa, H.P. (2018).** Management of rhinoceros beetle (*Oryctes rhinoceros* L.) by biological suppression with *Oryctes baculovirus* in Andhra Pradesh. *J. Plantation Crops*, 46(2): 124-127.
- Desai, V.S., Desai, S.D., Mayekar, A.J. and More, V.G. (2009).** Infestation of coconut eriophyid mite, *Aceria guerreronis* Keifer in Konkan region of Maharashtra. *J. Plant Protec. Sci.*, 1(1): 76-79.
- Gurav, S.S., Radadia, G.G. and Shinde, C.U. (2018).** Status of coconut black headed caterpillar, *Opisina arenosella* (Walker) and its associated larval parasitoid, *Goniozus nephantidis* (Muesebeck) under south Gujarat conditions. *Internat. J. Chemical Studies*, 6(2): 2713-2718.
- Levin, Lily and Mammooty, K.P. (2003).** Incidence of coconut eriophyid mite *aceria guerreronis* keifer (eriophyidae: acari) in different coconut cultivars and hybrids. *J. Tropical Agric.*, 41 : 59-62.
- Mathen, K. and Kurian, C. (1962).** Comparative efficacy of different insecticides on *Rhynchophorus ferrugineus* F. in proceedings of the First Conference of Coconut Research Workers in India 1959 - Indian Central Coconut Committee, Emakuum; P. 464.
- Mohanasundaram, M., Kalyanasundaram, S.K., Somasundaram, O.V.R. and Mahaendran, R. (1999).** Management and control measure for the coconut eriophyid mite *Aceria guerreronis* Keifer (Eriophyidae: Acari) in Tamil Nadu. *Indian Cocon. J.*, 29 (9): 8-10.
- Muyengi, Z.E., Msuya, E. and Lazaro, E. (2015).** Assessment of factors affecting coconut production in Tanzania. *J. Agric. Econ. & Development*, 4(6) : 083-094.
- Paul, W.D. (1985).** Study on integrated pest management (Col.: Scarab. Dynastinae). Ph.D. Thesis, FU Berlin.
- Sathiamma, B., Nair, C.P.R and Koshy, P.K. (1998).** Out break of nut infesting eriophyid mite *Eriophyes guerreronis* (K) in coconut plantation in India. *Indian Cocon. J.*, 29 (8): 1-3.
- Seguni, Z., Nyange, V., Materu, C., Mwingira, G. and Mruma, B. (2010).** Pest control research. In twenty five years of coconut research for development in Tanzania. pp. 78-96.
- Vanderplank, F.L. (1959).** Studies on the coconut pest *Pseudotheraptus wayi* Brown (Coroidae) in Zanzibar Some data on the yield of coconut in relation to damage caused by the insect. *Bull. Ent. Res.*, 50 : 135-149.

★ ★ ★ ★ ★ 12<sup>th</sup> Year of Excellence ★ ★ ★ ★ ★