

RESEARCH ARTICLE : **Development of non-spiny brinjal from spiny brinjal through conventional breeding**

■ **B.K. SAVITHA, M. PANDIYAN, P. VEERAMANI AND T. BALAJI**

ARTICLE CHRONICLE :

Received :

11.07.2017;

Accepted :

26.07.2017

SUMMARY : The Spiny brinjal (*S.melongena*) var. VRM-1 is a familiar and promising variety in Vellore district and also main dish to make a Ennai Kathirikai for briyaniside dish purpose. The spines in the stalk is a major constraints for picking and handling of brinjal fruits. A breeding initiative was undertaken at Agricultural Research Station and Krishi Vigyan Kendra, Virinjipuram to mask the spiny traits through crossing programme. The crosses were attempted to develop non-spiny brinjal with spiny quality. The both parents are pure for respective spiny and non-spiny. A total of 18 crosses were made with spiny brinjal and non-spiny brinjal progenies. The true F1 was tagged in both crosses combination. In F1 plant population of Spiny x Non-spiny, 100 per cent of the plants expressed spiny nature. In F1 plant population of non spiny x spiny reduced percentage of partial spiny plants were observed. From this study in both cases spiny plants are observed in F1 generation which indicates spiny is expressed as dominant characters. Further generation studies were made with continuous selfing programme and the results revealed that 100 per cent non-spiny plants were obtained with high yield from cross derivative of non-spiny x spiny 3rd cross and 5th cross during fourth generation. Comparative trials were carried out during *Kharif* 2015 and *Rabi* 2015-16 with spiny brinjal VRM-1. Mean performance showed that genotype non-spiny x spiny 3rd cross (1.43 kg/plant) registered the highest fruit yield per plant followed by non-spiny x spiny 5th cross (1.56 kg/plant). These genotypes could be used for further breeding programme for the improvement of fruit yield and non-spiny trait.

KEY WORDS :

Non-spiny brinjal,
Conventional
breeding

How to cite this article : Savitha, B.K., Pandiyan, M., Veeramani, P. and Balaji, T. (2017). Development of non-spiny brinjal from spiny brinjal through conventional breeding. *Agric. Update*, **12**(TECHSEAR-1) : 206-209; DOI: 10.15740/HAS/AU/12.TECHSEAR(1)2017/206-209.

Author for correspondence :

B.K. SAVITHA

Agricultural Research
Station and Krishi
Vigyan Kendra
(T.N.A.U.),
VIRINJIPURAM (T.N.)
INDIA
Email: savi_horti@
yahoo.co.in

See end of the article for
authors' affiliations

BACKGROUND AND OBJECTIVES

Brinjal (*Solanum melongena* L.) is an important and widely consumed Solanaceous vegetable of India grown round the year. The crop is extremely variable in India substantiated by the presence of wild relatives of *Solanum melongena* which are perennial herb or shrub with bitter fruits (Bhaduri, 1951). Though the principal method used for

improvement of this crop in India is selection from indigenous germplasm, yet comprehensive characterization of this crop has not been done (Choudhury, 1976). The spines in the stalk is a major constraints for picking and handling of brinjal fruits. Spiny brinjal (VRM-1 Mullukathiri) is a pure line selection from Elavambadi village of Vellore district. Spines are present in the leaf, stem

and calyx of the fruit. VRM-1 Mullukathiri is high yielding (30-35 t/ha⁻¹) and most suitable for North-Eastern Zone of Tamil Nadu. Selection for non-spiny plants from the spiny brinjal population was made and Crossing work was attempted to develop non-spiny fruits. The fruit set was ranged from 80-90 per cent. An attempt has been made in the present study to develop non-spiny brinjal with spiny quality for 18 genotypes. Two crosses (*i.e.*, Spiny x Non-spiny and Non-Spiny x Spiny) were attempted to study the inheritance of non-spiny character from Spiny brinjal. In both the crosses Non-Spiny character was obtained. Cross C1 (*i.e.*, Spiny x Non-spiny) expressed more number of non-spiny plants. However, spiny nature was observed in both the crosses indicating the partial dominance of spiny nature in both crosses.

RESOURCES AND METHODS

The experiment was laid out in a Randomized Block Design (RBD) with two replications. Ten genotypes (nine non-spiny genotypes and one spiny genotypes) were used for crossing programme. The crosses were attempted between spiny x non-spiny and non-spiny x spiny accessions. Nine non-spiny local accessions were crossed with VRM-1 spiny brinjal and reciprocal crosses were also made. Totally eighteen crosses were obtained. The generations were also forwarded from F1 to F5. The genotypes were evaluated and characterized for spyness and yield. Each plot consisted of 50 plants spaced at 60 x 75 cm. Five random plants per replication were selected to record observations on these characters.

OBSERVATIONS AND ANALYSIS

The results obtained from the present study as well as discussions have been summarized under following heads:

Performance of Non-spiny character in F3 generation :

Two crosses *i.e.*, Spiny x Non-spiny and Non-Spiny x Spiny was studied for the inheritance of non-spiny leaf, stem and fruits in brinjal (Table 1) in five accessions. In both the crosses Non-Spiny character was obtained. Cross C2 (*i.e.*, Non-Spiny x Spiny) expressed more number of non-spiny plants in F2 generation.

Performance of non-spiny character in F4 generation :

The F3 was found to be non-spiny in the cross Non-Spiny x Spiny, followed by Spiny x Non-spiny but only upto a certain extent. Eventhough, Spiny nature was observed in both the crosses (*i.e.*, Spiny x Non-spiny and Non-Spiny x Spiny) indicating the partial dominance inheritance of spiny. But, two cross from Non-Spiny x Spiny (C2 and C5) was found to be 100 per cent non-spiny nature indicating the dominance of non-spiny (Table 2).

Performance of non-spiny character in F5 generation with yield characters :

In this generation study, the 100 per cent non-spiny plants were obtained from cross derivative of non-spiny x spiny 3rd cross and 5th cross during fifth generation. These two Crosses were compared with spiny brinjal VRM-1 as yield trials in RBD during *Kharif* 2015 and *Rabi* 2015-16.

Out of 18 entries tested only two entries were selected based on single plant yield with non-spiny character. Among these two cross combination, the cross C2-5 recorded best yield performance (34.6 t/ha) at par with VRM-1 Spiny Brinjal (34.2 t/ha) (Table 3).

The comparative trial was repeated again during *Rabi* 2015-16 to check the non-spiny nature in *Rabi* 2015-16. The selected entries revealed 100% non-spiny nature

Table 1 : Inheritance of Non-spiny character in parents and crosses of VRM-1 Mullukathiri in F3 generation

	Non-spiny nature (%)					Mean
	Acc.1	Acc.2	Acc.3	Acc.4	Acc.5	
Parents						
Spiny Brinjal	Nil	Nil	Nil	Nil	Nil	Nil
Non-Spiny Brinjal	100.00	100.00	100.00	100.00	100.00	100.00
Crosses						
C1 (Spiny x Non-Spiny)	95.00	95.83	99.16	95.83	95.83	96.33
C2 (Non-Spiny x Spiny)	88.33	79.16	75.83	86.66	85.83	83.15

¹ Average of three replications containing 50 plants

Table 2 : Inheritance of Non-spiny character in parents and crosses of VRM-1 Mullukathiri in F4 generation

Sr. No.	Crosses	Non-Spiny plants (%)	No. of spines in leaf midrib	No. of spines in leaf veins
Crosses				
C1 (Spiny x Non-Spiny)				
1.	C1-1	0	3.0	2.0
2.	C1-2	0	4.0	2.0
3.	C1-3	0	2.5	3.0
4.	C1-5	0	3.0	1.0
5.	C1-6	0	3.5	1.5
6.	C1-7	7.69	4.0	4.0
7.	C1-8	0	3.0	3.5
8.	C1-9	0	4.2	3.0
	Mean	0.961	3.275	2.5
C2 (Non-Spiny x Spiny)				
9.	C2-1	81.42	1.0	0.5
10.	C2-2	85.20	1.5	1.2
11.	C2-3	100	0	0
12.	C2-4	88.24	1.3	1.5
13.	C2-5	100	0	0
14.	C2-6	87.15	1.4	1.7
15.	C2-7	80.46	1.4	2.1
16.	C2-8	80.53	1.5	2.5
17.	C2-9	85.27	1.1	1.7
18.	C2-10	88.34	1.7	2.2
	Mean	87.66	1.09	1.34
Parents				
1.	Spiny Brinjal	0	4.5	3.5
2.	Non-Spiny Brinjal	100	0	0

Table 3 : Yield of selected two entries in Kharif 2015

Sr. No.	Entries	Non-spiny nature (%)	Fruit yield/plant (kg)				Fruit yield (t/ha)	% of yield increase over check
			RI	RII	RIII	Mean		
1	C2-5	100	1.52	1.63	1.54	1.56	34.6	1.15
2	C2-3	100	1.72	1.65	1.52	1.43	31.7	-7.88
Check								
1	VRM-1 Spiny Brinjal	0	1.57	1.65	1.42	1.54	34.2	-

Table 4 : Yield of selected two entries in Rabi 2015-16

Sr. No.	Entries	Non-spiny nature (%)	Fruit yield/plant (kg)				Estimated fruit yield (t/ha)	% of yield increase over check
			RI	RII	RIII	Mean		
1	C2-5	100	1.61	1.51	1.36	1.49	33.1	4.22
2	C2-3	100	1.23	1.34	1.41	1.32	29.3	-8.19
Check								
1	VRM-1 Spiny Brinjal	0	1.44	1.52	1.34	1.43	31.7	-

and the cross C2-5 recorded best yield performance (33.1 t/ha) (Table 4).

Non-Spiny was observed in the present study was similar to that obtained by dominance inheritance of

spiny. Similar pattern of inheritance in brinjal for spiny leaf was reported earlier by Nimbalkar and More (1980) and More *et al.* (1982).

Authors' affiliations :

M. PANDIYAN, P. VEERAMANI AND T. BALAJI, Agricultural Research Station, and Krishi Vigyan Kendra (T.N.A.U.), VIRINJIPURAM (T.N.) INDIA

REFERENCES

Bhaduri, P.N. (1951). Inter-relationship of non-tuberiferous species of *solanum* with some consideration on the origin of brinjal. *Indian J. Genet.*, **2** : 75-86.

Choudhury, B. (1976). Brinjal (*Solanum melongena*). In: Evolution of crop plants, Simmonds NW (ed.) New York, pp

278-279.

More, D.C., Patil, S.B. and Nimbalkar, V.S. (1982). Inheritance of some characters in brinjal cross SM 2 x Nimbalkar Green Round. *J. Maharashtra agric. Univ.*, **7** : 243.

Nimbalkar, V.S. and More, D.C. (1980). Genetic studies in brinjal (*S. melongena* L.). *J. Maharashtra agric. Univ.*, **5**: 208.

Ramesh Kumar, S. and Arumugam, T. (2013). Phenotypic evaluation of indigenous Brinjal types suitable for rainfed conditions of South India (Tamil Nadu). *African J. Biotechnol.*, **12** (27) : 4338-4342.

★ ★ ★ ★ ★ **12th** Year of Excellence ★ ★ ★ ★ ★