

RESEARCH ARTICLE

# Study of different grains on spawn development of *P. sajor caju* (Fr.) Singer

■ NAZIMA KHATOON AND PRASHANT SHARMA

## SUMMARY

The important priority of profession is to maximize the production of mushroom by using various techniques. Mushroom seed (Spawn) production involves sophisticated technology with high investment requiring laboratory and equipments with accessories. The major constraint in mushroom production is non-availability of spawn. The technology of spawn production is given by Eco-care and Aware. We tested three grains viz., wheat, rice and gram for production of *Pleurotus sajor caju* spawn (Fr.) Singer. *Plurotus sajor caju* spawn has been recognized as a highly potential converter of cheap cellulosic material in to valuble protein at a very nominal cost. The results obtained during the present investigation are rice grains (9.00 days) were found to be the best grains for speedy development of spawn of *Pleurotus sajor caju* (Fr.) Singer.

**Key Words :** *Pleurotus sajor caju*, Different grains, Grain, Spawn

**How to cite this article :** Khatoon, Nazima and Sharma, Prashant (2017). Study of different grains on spawn development of *P. sajor caju* (Fr.) Singer. *Internat. J. Plant Sci.*, 12 (2): 220-222, DOI: 10.15740/HAS/IJPS/12.2/220-222.

**Article chronicle :** Received : 15.04.2017; Revised : 23.05.2017; Accepted : 14.06.2017

The cultivation of edible fungi has been accepted as the easiest on farm biotechnology for profitable removal of various lignocellulosic agricultural, industrial and forestry by-products, specially in the developing countries, although about 200 species of edible fungi have been reported from India, only three viz., *Pleurotus* spp., *Agraricus bisporus* and *Volvariella*

species are preferred for commercial cultivation and *Pleurotus* spp. alone constitutes about 65 per cent of the total mushrooms production in the country (Munjal, 1982). The most frequently used substrate for spawn production is wheat and there is need to use the more widely available maize grain as substrate for grain mother spawn production of oyster mushrooms (Mbogoh *et al.*, 2010).

Spawn is pure culture of mycelium growing on a solid substrate such as grain. Mushroom spawn has a fundamental role in global agricultural productivity. Its production has not been adopted by many farmers due to insufficient knowledge of scientific techniques needed for spawn production and mushroom cultivation. Just as

MEMBERS OF THE RESEARCH FORUM

**Author to be contacted :**  
PRASHANT SHARMA, Biotech Lab Training and Demonstration Centre, AMBIKAPUR (C.G.) INDIA

**Address of the Co-authors:**  
NAZIMA KHATOON, Biotech Lab Training and Demonstration Centre, AMBIKAPUR (C.G.) INDIA  
Email : [khan.nazi786@gmail.com](mailto:khan.nazi786@gmail.com)

seed quality is important to crop production, so is spawn quality to mushroom production. Oyster mushroom *Pleurotus sajor caju* (Fr.) Singer have great potential as an integral part of a sustainable agricultural system. Many types of organic wastes generated in agriculture or by the food processing industry are added to the waste stream annually. These materials could support oyster mushroom production, if systematically collected and processed. The major problem associated with the transfer of technology for spawn production and mushroom cultivation is the lack of technical know-how for its cultivation.

Present communication, different grains viz., wheat, rice and gram were screened to determine the most suitable grain spawn for better yield, biological efficiency and minimum spawn run time of *Pleurotus sajor caju* (Fr.) Singer. The aim of used various different grains are to get maximum production of fruiting bodies but it is an intricate problem. Moreover, the knowledge, shell help to plan and prepare for the crop and equip one with a physiological tool to control the quality, quantity, timing and other characters of the *Pleurotus sajor caju* (Fr.) Singer mushroom cultivation. Improved spawn production technology is necessary to increase the production of mushroom. Its production technology has not been standardized in Chhattisgarh. Therefore, an attempt has been made for find out the best grain for spawn production.

## MATERIAL AND METHODS

This work was carried out in Biotech Lab Training and Demonstration Center, Ambikapur, Chhattisgarh during April 2014 to July 2014. Parental strains of *Pleurotus species* (Fr.) Singer were provided by Mushroom Biotechnology Lab, Indira Gandhi Agriculture University, Raipur, Chhattisgarh. The cultures were maintained on potato dextrose agar medium (PDA) with regular sub-cultures at monthly intervals (Aneja, 2001). Mushroom spawn was prepared on wheat grain following standard procedure (Sivaprakasam, 1980). 15 days old spawn was used for experimentation. Three grains viz.,

wheat, rice and gram were evaluated as spawn substrates.

Clean and healthy grains were used for the preparation of mushroom spawn. All three grains were washed and soaked overnight in fresh water then boiled with two liter water per kg seed for 15-20 minutes till they become soft but remained firm. Water was drained off and boiled grains were spread over blotting paper to remove the excess water for 20-30 minutes. The grains were impregnated with 10 per cent calcium carbonate ( $\text{CaCO}_3$ ) and 10 per cent calcium sulphate ( $\text{CaSO}_4$ ) on the dry weight basis. Two hundred and fifty grams of coated grains was filled in 500 ml of narrow mouth bottles.

The bottles plugged by non-absorbent cotton. There were three replicates for each grain. These bottles were sterilized at 15 lbs pressure (PSI) and temperature  $121^\circ\text{C}$  for 30 minutes. The sterilized bottles were left for 12 hours and shaken for re-absorption of the condensed water droplets. Sterilized bottles containing grains were inoculated with 15 days old mother spawn @10% (w/w) of boiled grain under aseptic condition and the bottles were incubated at  $25^\circ\text{C}$  for 15 days.

## RESULTS AND DISCUSSION

The present study was undertaken with an aim to find out the best grains for preparation of early and best grain spawn. In the present work, three different grains were evaluated for spawn production of *Pleurotus sajor caju* (Fr.) Singer, an oyster mushroom. The parameter assessed in the study, deals with the number of days taken for spawn production from the day of inoculation to the growth and development of mycelial net on the different grains. The spawn development was determined by the white mat of mycelia of *P. sajor caju* (Fr.) Singer on each grain.

The results presented in Table 1 showed the time required for spawn development of *Pleurotus sajor-caju* (Fr.) Singer. The wheat and gram grain took same period (14.00 days) for spawn development of *Pleurotus sajor-caju* (Fr.) Singer. Rice grain recorded minimum

**Table 1 : Effect of different grains on spawn development of *Pleurotus species* (Fr.) Singer**

Sr. No.	Grains	<i>Pleurotus sajor caju</i> (Fr.) Singer spawn development period (Days)			
		Replication-I	Replication-II	Replication-III	Mean of three replication
1.	Wheat	12.0	14.0	16.0	14.00
2.	Rice	09.0	09.0	09.0	09.00
3.	Gram	13.0	13.0	14.0	13.33

(9.00 days) period for spawn development and recommended for spawn production of cultivation on *Pleurotus sajor-caju* (Fr.) Singer. Jiskani *et al.* (2000) conducted experiments on the effect of different grain on spawn growth of oyster mushroom, *Pleurotus florida* and reported that the sorghum grains were found to be best medium for spawn growth followed by maize, wheat and pearl millet grain, respectively.

Nwanze *et al.* (2006) examined the effect of spawn grains such as wheat, millet and corn on the culture of *lentinus squarrosulus*. The results showed that corn spawn induced highest yield and dry weight of fruiting as compared to wheat and millet spawn. These observations are in agreement with the result of Kotwaliwale *et al.* (1991) who found maize grains were suitable for spawn development of *Pleurotus* species and Sharma (2003) also found that kutki, jowar and maize grain took minimum period for spawn run of *Pleurotus* species.

It is clearly evident from the results that all the three grains are suitable for spawn production purpose. Hence, it is concluded that rice grain is best the grain spawn for production of oyster mushrooms *Pleurotus sajor-caju* (Fr.) Singer.

## REFERENCES

- Aneja, K.R. (2001). Experiments in microbiology, plant pathology, tissue culture and mushroom production technology, 3<sup>rd</sup> Ed. New Age International Publishers, 498-499.
- Jiskani, M.M., Wagan, K.H., Pathan, M.A and Jamro, G.H. (2000). Spawn growth of oyster mushroom as affected by different temperatures and grain media. *The Farm Scientist*, **6** (10): 33-35.
- Kotwaliwale, P., Kumar, S.M. and Ali, S.S. (1991). In Indian Mushrooms, Proceedings of National Symposium on Mushroom, organized by Kerala Agricultural University, Thiruvanthapuram, 72-73.
- Mbogoh, J.M., Anjichi, V.E., Rotich, F. and Ahoya, N.K. (2010). Substrate effects of grain spawn production on mycelium growth of Oyster Mushroom ISHS Acta Horticulturae **911**: I All Africa Horticultural Congress.
- Munjal, R.L. (1982). Prospects of mushroom cultivation in India. *Indian Phyto.*, **35** : 1-13.
- Nwanze, P.I., Khan, A.U., Ameh, A.U. and Umoh, V.J. (2005). The effect of the interaction of various oil types with different culture medium on biomass production of *Psathyrella atroumbonata* Pegler. *African J. Biotechnol.*, **4** : 1285-1289.
- Nwanze, P.I., Khan, A.U., Ameh, A.U. and Umoh, V.J. (2006). The effect of the interaction of various spawn grains with different culture medium on diameters of *Lentinus squarrosulus* (Mon) Singer. *African J. Biotechnol.*, **4** : 615-619.
- Sharma, B.B. (2003). Effects of different substrates (Grain straws) on Spawn growth and yield of pink oyster mushroom, *Pleurotus djamor* (Fr.) Boedijn. *J. Mycol. & Plant Pathol.*, **33** (2): 265-268.
- Sivaprakasam, K. (1980). Studies on oyster mushroom [*Pleurotus sajor caju* (Fr.) Singer]. P.h.D. Thesis, Tamil Nadu Agricultural University, Coimbatore, T.N. (INDIA). 115 pp.
- Yadav, M.C., Dhar, B.L., Verma, R.N. and Kamal, S. (1998). *International Symposium: Science and Cultivation of Mushroom*, Nanjing, October-1988, pp- 12-15.

12<sup>th</sup>  
Year  
★★★★★ of Excellence ★★★★★