

**RESEARCH PAPER**

Standardization of *Azospirillum* concentration and duration of biopriming for rice seed vigour improvement

M. KOKILA* AND M. BHASKARAN¹

Department of Seed Science and Technology, Tamil Nadu Agricultural University, COIMBATORE (T.N.) INDIA

(Email : dr.kokilam@gmail.com)

Abstract : *Azospirillum lipoferum* is nitrogen fixing biofertilizer which can be used in seed treatment for improving seedling vigour. In order to standardize the optimum concentration of liquid *Azospirillum* for rice biopriming seed treatment and duration of biopriming, a laboratory experiment was conducted in Department of Seed Science and Technology, Tamil Nadu Agricultural University, Coimbatore in Completely Randomized Design with four replicates. Rice hybrid CORH 4 and its parental lines COMS 23A and CB 174R seeds were bioprimed separately with 10, 15 and 20 per cent of *Azospirillum* for 6, 12, 18 and 24h of durations. Unprimed and hydroprimed seeds were used as control. The result revealed that rice seeds bioprimed with 20 per cent *Azospirillum* for 12 h was found to be the best seed biopriming treatment for both CORH 4 rice hybrid and its parental lines to enhance the germination rate, total germination percentage, seedling growth and vigour. In rice hybrid CORH 4, biopriming with 20 per cent liquid *Azospirillum* for 12h expressed an increase of 14 per cent of vigour index over the non-primed control seeds.

Key Words : *Azospirillum lipoferum*, Biopriming, Seed germination, Seedling vigour

View Point Article : Kokila, M. and Bhaskaran, M. (2016). Standardization of *Azospirillum* concentration and duration of biopriming for rice seed vigour improvement. *Internat. J. agric. Sci.*, **12** (2) : 283-287, DOI:10.15740/HAS/IJAS/12.2/283-287.

Article History : Received : 06.02.2016; Revised : 11.03.2016; Accepted : 05.05.2016

INTRODUCTION

Seed priming is now a widely used commercial process that accelerates the germination rate and improves seedling uniformity in many crops (Taylor and Harman, 1990). In priming, seeds are exposed to restricted water availability under controlled conditions which allows some of the physiological processes of germination to occur and before germination is completed, the seeds are usually re-dried for short term storage before sowing. Inoculation of seeds with biological agents in combination with priming has, in several cases, been

reported to enhance and stabilize the efficacy of biological agents (Callan *et al.*, 1991).

Biopriming is the process of biological seed treatment that refers to combination of seed hydration and inoculation with beneficial organism to enhance the seed germination and other character. Liquid bio-fertilizers has always more user friendly and eco-friendly effect than the inorganic fertilizers (Gomathy *et al.*, 2009). Many studies are available on the beneficial effects of inoculating biofertilizer on crop productivity. But, the studies on the effect of seed treatment with liquid biofertilizers on the germination and seedling vigour are

* Author for correspondence:

¹Seed Centre, Tamil Nadu Agricultural University, COIMBATORE (T.N.) INDIA

very scarce. The biofertilizer *Azospirillum lipoferum* was found to have not only the ability to fix nitrogen but also the ability to release phytohormones similar to gibberellic acid and indole acetic acid, which could stimulate plant growth, absorption of nutrients, and photosynthesis (Fayez *et al.*, 1985). Hence, the study was undertaken to standardize an optimum concentration and duration of soaking for seed bioprimering using liquid *Azospirillum* in rice hybrid CORH 4 and its parental lines (COMS 23A, CB 174R).

MATERIAL AND METHODS

Genetically pure seeds of rice hybrid CORH 4 and its parental lines (COMS 23A, CB 174R) were obtained from Department of Rice, Tamil Nadu Agricultural University, Coimbatore were bioprimered with liquid biofertilizer *Azospirillum lipoferum* obtained from the Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Coimbatore.

The laboratory experiment was conducted to standardize an optimum concentration of liquid biofertilizer and priming duration was conducted and analyzed with four replications in factorial completely randomized design (FCRD) at the Department of Seed Science and Technology, Tamil Nadu Agricultural University, Coimbatore.

For this study, the seeds of paddy hybrid CORH 4 and its parental lines were soaked in double the volume of water and 10, 15 and 20 per cent of liquid *Azospirillum* (*A. lipoferum*) solutions for 6h, 12h, 18h and 24h, separately. Unprimed seed served as control. Treated seeds along with control were assessed for the physiological seed quality parameters as follows.

The laboratory germination test was carried out in quadruplicate using 100 seeds each with 4 sub replicates of 25 seeds in the paper medium (ISTA, 2009). The test conditions of $25 \pm 2^\circ\text{C}$ temperature and 95 ± 3 per cent relative humidity were maintained in a germination room. Germination was counted at 24 hour intervals and continued until no further germination occurred. At the end of 14 days, the number of normal seedlings was counted and the mean was expressed as a percentage. Speed of germination was calculated based on the following formula of Maguire (1962):

$$\text{Speed of germination} = \frac{X_1}{Y_1} + \frac{X_2 - X_1}{Y_2} + \dots + \frac{X_n - X_{n-1}}{Y_n}$$

where; X_1 , X_2 and X_n are number of seeds

germinated on first, second and n^{th} day, respectively and Y_1 , Y_2 and Y_n are number of days from sowing to first, second and n^{th} count, respectively.

Root length and shoot length were measured at the time of germination count from ten normal seedlings selected at random from each replication and the mean was expressed in centimeter and those seedlings used for growth measurement were placed in a paper cover and dried in shade for 24h and then they were kept in an oven maintained at $85 \pm 2^\circ\text{C}$ for 48h. The dried seedlings were weighed to estimate the dry matter production and the mean values were expressed in g per 10 seedlings. The vigour index was computed as described by Abdul-Baki and Anderson (1973) as follows and expressed in whole numbers.

$$VI = GP \times (RL + SL)$$

where, VI is vigour index, GP is germination percentage, RL is root length (cm), SL is shoot length (cm).

The data obtained were analysed by the 'F' test of significance following the methods described by Rangaswamy (2002). The per cent values were transformed to arc-sine values and used for analysis. The critical differences (CD) were calculated at 1 and 5 per cent probability level. The data were tested for statistical significance by two way ANOVA.

RESULTS AND DISCUSSION

Biofertilizers have several beneficial roles in seed germination and seedling establishment. Beneficial bioagents contained the properties of plant growth promotion. Generally, the simplest method of beneficial microbial inoculation in crop production programme is the application of liquid suspension either directly to the soil or to the seeds. Carrier based biofertilizers had the problem of short shelf life, poor quality, high contamination and unpredictable field performances. However, liquid inoculant formulations showed good field performance with the use of low cost materials and easily attainable by small producers could overcome many problems associated with solid carriers (Singleton *et al.*, 2002).

In the present investigation, the rice seeds bioprimered with 20 per cent *Azospirillum* for 12 h quickened the germination with high percentage (Table 1 and 2) and recorded longest root and shoot (Table 3 and 4), maximum dry matter production (Table 5) and vigour index (Fig.1).

Table 1 : Standardization of seed biopriming with *Azospirillum* on speed of germination of rice hybrid CORH 4 and its parental lines COMS 23A and CB 174R

| Soaking duration (D) \ Biopriming treatment (T) | COMS 23A | | | | | CB 174R | | | | | CORH 4 | | | | |
|---|----------|-------|-------|-----------|------|---------|-------|-------|-----------|------|--------|-------|-------|-----------|------|
| | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean |
| Hydropriming | 5.9 | 6.8 | 6.3 | 6.1 | 6.3 | 5.9 | 6.3 | 6.3 | 6.1 | 6.2 | 5.7 | 7.3 | 7.1 | 6.1 | 6.6 |
| <i>Azospirillum</i> 10 % | 6.5 | 7.1 | 6.8 | 6.3 | 6.7 | 5.9 | 6.0 | 6.1 | 6.3 | 6.1 | 5.6 | 7.4 | 7.2 | 6.3 | 6.6 |
| <i>Azospirillum</i> 15 % | 5.9 | 7.1 | 7.1 | 6.5 | 6.7 | 6.1 | 6.6 | 6.8 | 6.3 | 6.5 | 6.3 | 7.5 | 7.3 | 6.4 | 6.9 |
| <i>Azospirillum</i> 20% | 6.4 | 7.6 | 7.3 | 6.8 | 7.0 | 6.1 | 7.8 | 7.5 | 6.8 | 6.9 | 6.3 | 7.6 | 7.5 | 7.2 | 7.2 |
| Mean | 6.2 | 7.2 | 6.9 | 6.4 | 6.7 | 6.0 | 6.7 | 6.4 | 6.4 | 6.4 | 6.0 | 7.4 | 7.3 | 6.5 | 6.8 |
| Nonprimed seed (C) | | | 5.8 | | | | | 5.6 | | | | | 5.8 | | |
| | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | |
| S.E. ± | 0.052 | 0.052 | 0.104 | 0.076 | | 0.056 | 0.056 | 0.113 | 0.082 | | 0.181 | 0.181 | 0.362 | 0.264 | |
| C.D. (P=0.05) | 0.104 | 0.104 | 0.208 | 0.152 | | 0.113 | 0.113 | 0.227 | 0.165 | | 0.364 | 0.364 | 0.728 | 0.531 | |

Table 2 : Standardization of seed biopriming with *Azospirillum* on seed germination (%) of rice hybrid CORH 4 and its parental lines COMS 23A and CB 174R

| Soaking duration (D) \ Biopriming treatment (T) | COMS 23A | | | | | CB 174R | | | | | CORH 4 | | | | |
|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean |
| Hydropriming | 89 (70.63) | 91 (72.54) | 91 (72.54) | 90 (71.57) | 90 (71.57) | 90 (71.57) | 92 (73.57) | 92 (73.57) | 91 (72.54) | 91 (72.54) | 88 (69.73) | 90 (71.57) | 89 (70.63) | 89 (70.63) | 89 (70.63) |
| <i>Azospirillum</i> 10 % | 89 (70.63) | 91 (72.54) | 92 (73.57) | 90 (71.57) | 91 (72.54) | 90 (71.57) | 92 (73.57) | 92 (73.57) | 91 (72.54) | 91 (72.54) | 88 (69.73) | 90 (71.57) | 89 (70.63) | 89 (70.63) | 89 (70.63) |
| <i>Azospirillum</i> 15 % | 90 (71.57) | 93 (74.66) | 92 (73.57) | 91 (72.54) | 92 (73.57) | 91 (72.54) | 94 (75.82) | 94 (75.82) | 92 (73.57) | 93 (74.66) | 88 (69.73) | 90 (71.57) | 90 (71.57) | 89 (70.63) | 89 (70.63) |
| <i>Azospirillum</i> 20 % | 91 (72.54) | 94 (75.82) | 93 (74.66) | 92 (73.57) | 92 (73.57) | 93 (74.66) | 95 (77.08) | 94 (75.82) | 93 (74.66) | 94 (75.82) | 89 (70.63) | 93 (74.66) | 92 (73.57) | 89 (70.63) | 91 (72.54) |
| Mean | 90 (71.57) | 92 (73.57) | 92 (73.57) | 91 (72.54) | 91 (72.54) | 91 (72.54) | 93 (74.66) | 93 (74.66) | 92 (73.57) | 92 (73.57) | 88 (69.73) | 91 (72.54) | 90 (71.57) | 89 (70.63) | 90 (71.57) |
| Nonprimed seed (C) | | | 89 (70.63) | | | | | 90 (71.57) | | | | | 88 (69.73) | | |
| | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | |
| S.E. ± | 0.643 | 0.643 | 1.286 | 0.937 | | 0.678 | 0.678 | 1.356 | 0.989 | | 0.427 | 0.427 | 0.854 | 0.623 | |
| C.D. (P=0.05) | 1.292 | 1.292 | NS | 1.883 | | 1.362 | 1.362 | NS | 1.986 | | 0.858 | 0.858 | NS | 1.251 | |

Figures in parentheses indicate arcsine values

NS=Non-significant

Table 3 : Standardization of seed biopriming with *Azospirillum* on root length (cm) of rice hybrid CORH 4 and its parental lines COMS 23A and CB 174R

| Soaking duration (D) \ Biopriming treatment (T) | COMS 23A | | | | | CB 174R | | | | | CORH 4 | | | | |
|---|----------|-------|-------|-----------|------|---------|-------|-------|-----------|------|--------|-------|-------|-----------|------|
| | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean |
| Hydropriming | 21.2 | 23.5 | 22.2 | 22.0 | 22.2 | 18.8 | 19.3 | 19.2 | 19.0 | 19.1 | 21.1 | 23.0 | 22.6 | 22.2 | 22.2 |
| <i>Azospirillum</i> 10 % | 21.4 | 23.6 | 23.0 | 22.0 | 22.5 | 19.0 | 19.5 | 19.4 | 19.4 | 19.3 | 21.2 | 23.0 | 22.8 | 22.2 | 22.3 |
| <i>Azospirillum</i> 15 % | 22.0 | 24.0 | 23.1 | 22.8 | 23.0 | 19.5 | 21.0 | 21.0 | 19.8 | 20.3 | 21.3 | 23.0 | 23.0 | 22.9 | 22.5 |
| <i>Azospirillum</i> 20 % | 22.8 | 25.0 | 24.8 | 23.8 | 24.1 | 20.0 | 22.0 | 21.6 | 21.2 | 21.2 | 21.3 | 23.8 | 23.1 | 23.1 | 22.8 |
| Mean | 22.0 | 24.0 | 23.3 | 22.7 | 23.0 | 19.3 | 20.5 | 20.3 | 19.9 | 20.0 | 21.2 | 23.2 | 22.8 | 22.6 | 22.5 |
| Nonprimed seed (C) | | | 21.0 | | | | | 18.5 | | | | | 20.5 | | |
| | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | |
| S.E. ± | 0.141 | 0.141 | 0.283 | 0.206 | | 0.113 | 0.113 | 0.226 | 0.165 | | 0.172 | 0.172 | 0.344 | 0.251 | |
| C.D. (P=0.05) | 0.284 | 0.284 | 0.568 | 0.414 | | 0.227 | 0.227 | 0.454 | 0.331 | | 0.345 | 0.345 | 0.691 | 0.504 | |

Table 4 : Standardization of seed biopriming with *Azospirillum* on shoot length (cm) of rice hybrid CORH 4 and its parental lines COMS 23A and CB 174R

| Soaking duration (D) / Biopriming treatment (T) | COMS 23A | | | | | CB 174R | | | | | CORH 4 | | | | |
|---|----------|-------|-------|-----------|------|---------|-------|-------|-----------|------|--------|-------|-------|-----------|------|
| | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean |
| Hydropriming | 11.2 | 12.0 | 11.8 | 11.8 | 11.7 | 10.3 | 11.0 | 10.5 | 10.5 | 10.6 | 11.5 | 12.4 | 12.3 | 12.2 | 12.1 |
| <i>Azospirillum</i> 10 % | 11.4 | 12.0 | 12.1 | 12.0 | 11.9 | 10.5 | 11.3 | 11.2 | 10.8 | 11.0 | 11.7 | 12.4 | 12.3 | 12.3 | 12.2 |
| <i>Azospirillum</i> 15 % | 11.6 | 12.3 | 12.3 | 12.2 | 12.1 | 11.0 | 11.6 | 11.5 | 11.2 | 11.3 | 11.8 | 12.6 | 12.4 | 12.3 | 12.3 |
| <i>Azospirillum</i> 20 % | 11.8 | 12.7 | 12.5 | 12.3 | 12.3 | 11.2 | 12.2 | 11.8 | 11.6 | 11.7 | 12.3 | 12.8 | 12.6 | 12.5 | 12.6 |
| Mean | 11.6 | 12.3 | 12.2 | 12.1 | 12.0 | 10.8 | 11.5 | 11.3 | 11.0 | 11.1 | 11.8 | 12.6 | 12.4 | 12.3 | 12.3 |
| Nonprimed seed (C) | 11.1 | | | | | 10.2 | | | | | 11.3 | | | | |
| | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | |
| S.E. ± | 0.083 | 0.083 | 0.167 | 0.122 | | 0.059 | 0.059 | 0.119 | 0.087 | | 0.103 | 0.103 | 0.207 | 0.151 | |
| C.D. (P=0.05) | 0.168 | 0.168 | 0.335 | 0.244 | | 0.119 | 0.119 | 0.239 | 0.174 | | 0.208 | 0.208 | 0.415 | 0.303 | |

Table 5 : Standardization of seed biopriming with *Azospirillum* on drymatter production (g / 10 seedlings) of rice hybrid CORH 4 and its parental lines COMS 23A and CB 174R

| Soaking duration (D) / Biopriming treatment (T) | COMS 23A | | | | | CB 174R | | | | | CORH 4 | | | | |
|---|----------|-------|-------|-----------|-------|---------|-------|-------|-----------|-------|--------|-------|-------|-----------|-------|
| | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean | 6h | 12h | 18h | 24h | Mean |
| Hydropriming | 0.108 | 0.125 | 0.123 | 0.118 | 0.118 | 0.085 | 0.090 | 0.090 | 0.087 | 0.088 | 0.117 | 0.123 | 0.120 | 0.115 | 0.119 |
| <i>Azospirillum</i> 10 % | 0.110 | 0.125 | 0.123 | 0.120 | 0.120 | 0.088 | 0.092 | 0.093 | 0.089 | 0.091 | 0.118 | 0.126 | 0.127 | 0.120 | 0.123 |
| <i>Azospirillum</i> 15 % | 0.110 | 0.126 | 0.126 | 0.120 | 0.121 | 0.09 | 0.096 | 0.094 | 0.092 | 0.093 | 0.119 | 0.125 | 0.125 | 0.124 | 0.123 |
| <i>Azospirillum</i> 20 % | 0.115 | 0.128 | 0.128 | 0.125 | 0.124 | 0.090 | 0.110 | 0.098 | 0.094 | 0.098 | 0.120 | 0.134 | 0.130 | 0.125 | 0.127 |
| Mean | 0.111 | 0.126 | 0.125 | 0.122 | 0.121 | 0.088 | 0.097 | 0.094 | 0.090 | 0.092 | 0.118 | 0.127 | 0.125 | 0.121 | 0.123 |
| Nonprimed seed (C) | 0.105 | | | | | 0.083 | | | | | 0.111 | | | | |
| | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | | T | D | T×D | C vs Rest | |
| S.E. ± | 0.001 | 0.001 | 0.002 | 0.001 | | 0.001 | 0.001 | 0.001 | 0.001 | | 0.001 | 0.001 | 0.003 | 0.002 | |
| C.D. (P=0.05) | 0.002 | 0.002 | 0.004 | 0.003 | | 0.001 | 0.001 | 0.003 | 0.002 | | 0.003 | 0.003 | 0.005 | 0.004 | |




Fig. 1 : Seed biopriming with *Azospirillum* on vigour index of rice hybrid CORH 4 and itsparental lines COMS 23A and CB 174R

The increase of vigour index in seeds bioprimered with 20 per cent *Azospirillum* was 10, 15 and 4 per cent over hydroprimed seeds and 18, 21 and 14 per cent over nonprimed seeds of COMS 23A, CB 174R and CORH 4, respectively. The increase in physiological attributes of *Azospirillum* bioprimered seeds might be due to the production of biologically active substance such as auxins, gibberellins, cytokinins, amino acids and vitamins (Afifi *et al.*, 2003). The phytohormones released by *Azospirillum lipoferum* could stimulate plant growth, absorption of nutrients and photosynthesis (Fayez *et al.*, 1985). Biofortification of rice seeds with *Azospirillum* enhanced seedling vigour, speed of germination, seedling length and dry weight of both high and low vigour seed lots (Ramamoorthy *et al.*, 2000).

Azospirillum sp. are non-specific plant growth-promoting rhizobacterias (PGPRs) providing many contributions to the enhancement of growth in many agricultural crop species (Bharathi *et al.*, 2004). Similar result was observed by Kapulnik *et al.* (1981) in rice; Raju *et al.* (1999) in sorghum; Niranjana *et al.* (2004) in pearl millet.

REFERENCES

- Abdul-Baki, A.A. and Anderson, J.D. (1973).** Vigour deterioration of soybean seeds by multiple criteria. *Crop Sci.*, **13** : 630-633.
- Afifi, M.H., Manal, F.M. and Goma, A.M. (2003).** Efficiency of applying biofertilizers to maize crop under different levels of mineral fertilizers. *Ann. Agric. Sci. Moshtohor.*, **41**(4) : 1411-1420.
- Bharathi, R., Vivekananthan, R., Harish, S., Ramanathan, A. and Samiyappan, R. (2004).** Rhizobacteria-based bioformulations for the management of fruit rot infection in chillies. *Crop Protec.*, **23** : 835-843.
- Callan, N.W., Mathre, D.E. and Miller, J.B. (1991).** Field performance of sweet corn seed bioprimered and coated with *Pseudomonas fluorescens* ab254. *Hort. Sci.*, **26** : 1163-1165.
- Fayez, M., Emam, N.F. and Makhoul, H.F. (1985).** The possible use of nitrogen fixing *Azospirillum* as biofertilizer for wheat plants. *Egypt. J. Microbiol.*, **20**(2) : 199-206.
- Gomathy, M., Thangaraju, M. and Sarathambal, C. (2009).** Optimization of liquid phosphobacteria required for maize seeds. *Asian J. Bio. Sci.*, **4**(1) : 53-55.
- ISTA (2009). International rules for seed testing. International Seed Testing Association, Bassersdorf, Switzerland.
- Kapulnik, Y., Kigel, Y., Okon, Y., Nur, I. and Henis, Y. (1981).** Effect of *Azospirillum* inoculation on some growth parameters and N content of wheat sorghum and panicum. *Pl. Soil.*, **61**: 65-70.
- Maguire, J.D. (1962).** Speed of germination - Aid in selection and evaluation of seedling emergence and vigour. *Crop Sci.*, **2** : 176-177.
- Niranjana, S.R., Shetty, N.P. and Shetty, H.S. (2004).** Seed bio-priming with *Pseudomonas fluorescens* isolates enhances growth of pearl millet plants and induces resistance against downy mildew. *J. Pest. Mgmt.*, **50**(1) : 41-48.
- Raju, N.S., Niranjana, S.R., Janardhana, G.R., Prakash, H.S., Shetty, H.S. and Mathur, S.B. (1999).** Improvement of seed quality and field emergence of *Fusarium moniliforme* infected sorghum seeds using biological agents. *J. Sci. Food. Agric.*, **79**: 206-212.
- Ramamoorthy, K., Natarajan, N. and Lakshmanan, A. (2000).** Seed biofortification with *Azospirillum* spp. for improvement of seedling vigour and productivity in rice (*Oryza sativa* L.). *Seed Sci. & Technol.*, **28** : 809-815.
- Rangaswamy, R. (2002).** *A text book of agricultural statistics.* New Age International Pvt., Ltd. NEW DELHI, (INDIA).
- Singleton, P.W., Keyser, H.H. and Sande, E.S. (2002).** Development and evaluation of liquid inoculants. In: *Inoculants and nitrogen fixation of legumes in Vietnam.* (ed.) D. Herridge, ACIAR Proceedings, pp: 52-66.
- Taylor, A.G. and Harman, G.E. (1990).** Concepts and technologies of selected seed treatments. *Phytopathol.*, **28**: 321-339.

12th
Year
★★★★★ of Excellence ★★★★★