

## RESEARCH ARTICLE :

# Seasonal incidence and correlation of abiotic factors against okra shoot and fruit borer (*Earias vittella* Fab.) during *Kharif* season

■ AGURLA RAJU, C. NARENDRA REDDY, D. ANITHA KUMARI AND D. SRINIVASA CHARY

## ARTICLE CHRONICLE :

## Received :

12.07.2017;

## Accepted :

25.07.2017

## KEY WORDS :

Okra shoot, Fruit borer, Standard week, Correlation, Seasonal incidence, Weather parameters

**SUMMARY :** Field experiment was carried out with a view to study the fluctuations in the population of okra shoot and fruit borer against prevailing weather conditions at student farm, college of Agriculture, Hyderabad, PJTSAU during *Kharif*-2015. The results recorded that the incidence of okra shoot and fruit borer was commenced in the 35<sup>th</sup> standard week (0.5 per plant), peak incidence were recorded in terms of shoot infestation in 40<sup>th</sup> standard week (30.63%), fruit infestation on number basis in 42<sup>nd</sup> standard week (58.32%) and fruit infestation on weight basis recorded in 45<sup>th</sup> standard week (33.42%). Maximum larval population was noticed in 40<sup>th</sup> standard week (6.4 per 10 plants). Larval population, fruit damage on number and weight basis were significantly negatively correlated with maximum temperature and non-significantly negatively correlated with minimum temperature, relative humidity and rain fall. Incidence of shoot damage was significantly correlated with maximum temperature and minimum temperature and non-significantly negatively correlated with rain fall and relative humidity.

**How to cite this article :** Raju, Agurla, Reddy, C. Narendra, Kumari, D. Anitha and Chary, D. Srinivasa (2017). Seasonal incidence and correlation of abiotic factors against okra shoot and fruit borer (*Earias vittella* Fab.) during *Kharif* season. *Agric. Update*, 12(TECHSEAR-2) : 554-558; DOI: 10.15740/HAS/AU/12.TECHSEAR(2)2017/554-558.

## Author for correspondence :

## AGURLA RAJU

Department of Entomology, College of Agriculture, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, HYDERABAD (TELANGANA) INDIA  
Email: [agurla25@gmail.com](mailto:agurla25@gmail.com)

See end of the article for authors' affiliations

## BACKGROUND AND OBJECTIVES

Okra [*Abelmoschus esculentus* (L.) Moench] is an important vegetable crop of family Malvaceae. It plays an important role in human diet and has good nutritional value. The crop is attacked by more than 45 insect pests which infests the crop from seedling to harvest stage (Nair, 1984), among which shoot and fruit borer, *Earias vittella* (Fab) are the most important and serious pest causing direct damage to marketable fruits. The caterpillar

passes through 6 stages, becoming full-grown in 10-16 days and the affected fruits become unfit for consumption (Kedar *et al.*, 2014). Eggs and larvae were noticed on five weeks old crop during all the cropping seasons (Siddhartha *et al.*, 2017). The fruit borer alone to reported cause damage to the extent of 3.5 to 90 per cent to okra in different parts of the country (Mandal *et al.*, 2006). Incidence of the fruit borer commenced in 29<sup>th</sup> standard week and maximum number of larva (7.5

larvae per 10 plants) were recorded in 42<sup>nd</sup> standard week (Sharma *et al.*, 2010). The maximum damage was caused to fruit in 41<sup>st</sup> standard week (Nath *et al.*, 2011). Dangi (2004) observed that the incidence of okra shoot and fruit borer commenced from the 4<sup>th</sup> week of August (6<sup>th</sup> after sowing). Yadav *et al.* (2007) reported that maximum fruit damage was observed in the 3<sup>rd</sup> week of September. Yadvendu (2001) reported that maximum damage 86.5 and 72.5 per cent during first and fourth week of September. Pareek *et al.* (2001) reported damage was started in first week of September. Gautham *et al.* (2014); Dhabi *et al.* (2013) and Aziz *et al.* (2011) reported that weather parameters were significantly correlated with population of okra shoot and fruit borer. The incidence and spread of okra shoot and fruit borer largely controlled by various weather factors *viz.*, temperature, relative humidity and rain fall. Information on interaction of weather parameters and insect development can provide vital role in pest surveillance, forecasting, monitoring and management of pest population by timely decision making. Hence, an attempt has, therefore, been made to study the population dynamics of okra shoot and fruit borer in relation to prevailing weather parameters, so that the information can effectively be utilized in formulating pest management programme.

## RESOURCES AND METHODS

The field experiment was conducted at Prof. Jayashankar Telangana State Agricultural University, Hyderabad, Telangana during *Kharif*-2015. The experimental field was located at Southern Telangana Zone with longitude '78.415503', latitude '17.316171' and Mean Sea Level 546 m. A popular okra variety "Arka Anamika" was selected to conduct the experiment. Sowing was taken up on 15<sup>th</sup> July 2015 at 60 × 30 cm spacing. Seed rate followed was 4 kilograms per acre. The crop was raised as per the package and practices of PJTSAU. Ten plants from each plot were randomly selected and tagged. Weekly observations on was taken on entire tagged plants thought the season. Incidence of okra shoot and fruit borer was recorded in terms of per cent infested plants, no of larva per ten plants, per cent fruit damage on number and weight basis. Incidence of okra shoot and fruit borer in terms of percentage of damaged fruits on number and weight was recorded by counting and weighing healthy and damaged fruits at each

picking separately using formula:

$$\text{Per cent fruit damage on number/weight} = \frac{\text{Number/weight of damaged fruits}}{\text{Total number/weight of fruits}} \times 100$$

The weekly meteorological data during the period of experiment were collected from Agro meteorological Observatory, Agricultural Research Institute, Rajendranagar, Hyderabad, Telangana. Weekly mean Maximum temperature, minimum temperature, mean relative humidity and total weekly rainfall were used to work out the association of weather parameters on infestation of the pest on okra. In order to study the influence of key abiotic factors on pest incidence, simple correlations were worked out between the pest incidence and meteorological factors.


## OBSERVATIONS AND ANALYSIS

Population of dynamics of okra shoot and fruit borer was conducted at department of Entomology, College of Agriculture, PJTSAU, Telanagana. The results indicated that *Earias vittella* (Fab) infestation on okra was commenced in 35<sup>th</sup> standard week (3<sup>rd</sup> week of August) that continuously increased till the last picking in the 46<sup>th</sup> standard week (3<sup>rd</sup> week of November) (Table 1 and Fig. 1). It was observed that the infestation of okra shoot and fruit borer was started in the 4<sup>th</sup> week of September with an average larval population 0.5 larvae per 10 plants in *Kharif* 2015. This infestation continuously increased and reached to the peak 6.4 larvae per 10 plants in 40<sup>th</sup> standard week at maximum temperature 31.4°C (Fig. 2), relative humidity 75.5% (Fig. 3), number of rainy days 4.9 (Fig. 4) and thereafter, decreased gradually upto 44<sup>th</sup> week standard week with average population of 6.1 larvae per 10 plants (Table 1 and Fig. 1). Similarly results was recorded by Dangi (2004) 10.3 larvae per 10 plants in first week of October. Siddartha *et al.* (2017) also recorded average larval population of *Earias* spp 1.34±0.18 larvae per plant. Similarly Sharma *et al.* (2010) noticed 0.3 and 0.5 larvae per 10 plants during *Kharif* 2005 and 2006, respectively.


The shoot infestation was commenced on 3<sup>rd</sup> week of August (35<sup>th</sup> standard week) with 5.1 per cent shoot damage. This per cent shoot damage was continuously increased upto 40<sup>th</sup> standard week (1<sup>st</sup> week of October) and maximum shoot damage was recorded 30.63 per cent in 40<sup>th</sup> standard week at maximum temperature 31.4°C (Fig. 2), relative humidity 75.5% (Fig. 3), number of rainy days 4.9 (Fig. 4) and gradually decreased till the

end of crop upto 46<sup>th</sup> standard week with average shoot infestation 19.14 per cent (Table 1 and Fig. 1). Similar results reported by Siddarth *et al.* (2017) with mean shoot


damage  $4.61 \pm 0.11$  per cent and peak infestation was recorded in 45<sup>th</sup> standard week 91.6 per cent (Sharma *et al.*, 2010). The fruit infestation on number was




**Fig. 1:** Seasonal incidence of okra shoot and fruit borer (*Earias vittella*, Fabricius) in different standard weeks during Kharif-2015



**Fig. 3:** Population dynamics of okra shoot and fruit borer against prevailing relative humidity in different standard weeks (from standard week 34 to 46, respectively)



**Fig. 2:** Population dynamics of okra shoot and fruit borer against prevailing temperatures in different standard weeks (from standard week 34 to 46, respectively)



**Fig. 4:** Population dynamics of okra shoot and fruit borer against prevailing rain fall (mm) in different standard weeks (from standard week 34 to 46, respectively)

**Table 1 :** Seasonal incidence of okra shoot and fruit borer (*Earias vittella*, Fabricius) during Kharif-2015

Standard week	Maximum temperature (°C)	Minimum temperature (°C)	Mean RH (%)	No. of rain fall (mm)	No. of larvae/10 plants	Per cent shoot infestation	Per cent fruit damage on number basis	Per cent fruit damage on weight basis
34	30.4	22.43	79.35	4.29	-	-	-	-
35	31.5	22.71	77.57	7.06	0.5	5.1	-	-
36	33.4	22.9	73.75	4.4	1.5	11.61	10.35	8.07
37	28.4	21.9	90.8	13.1	2	14.59	25.42	13.56
38	30.4	22.2	75.3	6.2	1.3	13.5	30.16	16.31
39	31.9	22.3	73.5	0.3	4.2	25.9	32.64	21.53
40	31.4	21.1	75.7	4.9	6.4	30.63	42.56	23.46
41	33.4	19.6	62.9	0	6.3	29.25	46.76	25.89
42	32.4	19.1	66.85	0	5.2	23.16	58.32	28.42
43	32.4	18.1	66.45	0	4.4	22.87	56.21	30.16
44	31.4	20.7	71.3	2.6	5.2	20.42	50.75	31.26
45	31.4	17.4	82.1	0	6.1	21.32	49.17	33.42
46	30.8	15.8	69	0	4.1	19.12	48.12	31.21

commenced on 4<sup>th</sup> week of August (36<sup>th</sup> standard week) and mean fruit damage was recorded 10.35 per cent. The mean fruit damage on number was continuously increased upto 42<sup>nd</sup> standard week and maximum damage was recorded 58.32 per cent in the 42<sup>nd</sup> standard week at maximum temperature 32.4°C (Fig. 2) and relative humidity 66.85% (Fig. 3 and Table 1). Sharma *et al.* (2010) reported fruit damage on number was started in first week of August and recorded maximum damage 54.3 per cent in the 3<sup>rd</sup> week of October (42<sup>nd</sup> standard week). The fruit infestation on weight was started on 4<sup>th</sup> week of August (36<sup>th</sup> standard week) and mean fruit damage on weight was recorded 8.07 per cent. Fruit infestation on weight was continuously increased till the end of crop growth and maximum damage on weight was recorded in the 46<sup>th</sup> standard week (33.26%) at maximum temperature 31.4°C (Fig. 2) and relative humidity 82.1% (Fig. 3 and Table 1). Similar results reported by Sharma *et al.* (2010) that maximum damage noticed on weight basis was 54.7 per cent when the crop was 18 weeks old in 42<sup>nd</sup> standard week. Dangi (2004) also reported maximum fruit damage 72.20 per cent on weight basis.

It is evident from Table 2 that mean larval population was non significantly negatively correlated with maximum temperature ( $r=-0.310668$ ), significantly negatively correlated with minimum temperature ( $r=-0.57949$ ), non-significantly negatively correlated with mean relative humidity ( $r=-0.53234$ ) and significantly negatively correlated with rain fall ( $r=-0.64478$ ). Similar results were reported by Sharma *et al.* (2010) in terms of larval population showed significant negative correlation with mean temperature ( $-r=0.678$ ) and mean relative humidity ( $r=-0.241$ ), but non-significant correlation with rain fall ( $r=-0.241$ ). Siddartha *et al.* (2017) also noticed maximum temperature showed negative correlation ( $r=-0.51$ ) with larval population. Mandal *et al.* (2006) also reported maximum temperature had negative impact on larval population. Incidence of shoot damage was

significantly negatively correlated with maximum ( $r=-0.29381$ ) and minimum temperature ( $r=-0.39122$ ), whereas non significantly negatively correlated with relative humidity ( $r=-0.617$ ) and rain fall ( $r=-0.56084$ ). Similar results reported by Siddartha *et al.* (2017) in terms of shoot infestation, showed that significant negative correlation with maximum temperature ( $r=-0.21$ ), minimum temperature ( $r=-0.08$ ), relative humidity ( $r=-0.45$ ) and rainfall ( $r=-0.05$ ).

Incidence of fruit damage on number basis was significantly correlated with maximum temperature, non-significantly correlated with minimum temperature, relative humidity and rain fall (where,  $r=-0.153141$ ,  $r=-0.74543$ ,  $r=-0.57037$  and  $r=-0.61888$ , respectively). Incidence of fruit damage on weight basis significantly negatively correlated with, maximum temperature, non-significantly correlated with minimum temperature, relative humidity and rainfall ( $r=-0.13657$ ,  $r=-0.81850$ ,  $r=-0.64345$  and  $r=-0.69557$ , respectively). Similar results were reported by Mandal *et al.* (2006) in terms of fruit damage on weight basis, results noticed that maximum temperature had negatively correlated with fruit damage on weight basis ( $r=-0.496$ ). Siddartha *et al.* (2017) also reported that the fruit damage on weight basis was negatively correlated maximum temperature ( $r=-0.35$ ), relative humidity ( $r=-0.04$ ) and rain fall ( $r=-0.02$ ).

## Conclusion :

The incidence of okra shoot and fruit borer was commenced in the 35<sup>th</sup> standard week, peak incidence were recorded in terms of shoot infestation in 40<sup>th</sup> standard week, fruit infestation on number basis in 43<sup>rd</sup> standard week and fruit infestation on weight basis recorded in 45<sup>th</sup> standard week. Maximum larval population was noticed in 40<sup>th</sup> standard week. Larval population, fruit damage on number and weight basis were significantly negatively correlated with maximum temperature and non-significantly negatively correlated with minimum temperature, relative humidity and rain

**Table 2: Correlation co-efficient (r) of pest population and damage of *Earias vittella* (Fabricius) against prevailing weather parameters**

Weather parameters	No. of larvae per 10 plants	Per cent shoot damage	Per cent fruit damage on number basis	Fruit damage on weight basis
Maximum temperature	-0.31066*	-0.29381*	-0.153141*	-0.13657*
Minimum temperature	-0.57949**	-0.39122*	-0.74543	-0.81850
Mean relative humidity	-0.53234**	-0.617**	-0.57037**	-0.64345**
No. of rain fall	-0.64478**	-0.56084**	-0.61888**	-0.69557

\* and \*\* indicate significance of values at  $P=0.05$  and  $0.01$ , respectively

fall. Incidence of shoot damage was significantly correlated with maximum temperature and minimum temperature, and non-significantly negatively correlated with rain fall and relative humidity.

### Acknowledgement:

The researcher's team is sincerely thankful to the Professor Jayashankar Telangana State Agricultural University (PJTSAU) for providing research facilities. We also thank Department of Entomology for providing financial assistance through university during research.

### Authors' affiliations :

**C. NARENDRA REDDY**, Department of Entomology, College of Agriculture, , Professor Jayashankar Telangana State Agricultural University, Rajendranagar, HYDERABAD (TELANGANA) INDIA

**D. ANITHA KUMARI**, Vegetable Research Station (SKLTSHU), ARI, Rajendranagar, HYDERABAD (TELANGANA) INDIA

**D. SRINIVASA CHARY**, Department of Statistics and Mathematics, College of Agriculture, , Professor Jayashankar Telangana State Agricultural University, Rajendranagar, HYDERABAD (TELANGANA) INDIA

### REFERENCES

- Aziz, M.A.**, Hassan, M. and Ali, A. (2011). Impact of abiotic factors on incidence of fruit and shoot damage of spotted bollworms, *Earias* spp. On okra (*Ablemoschus esculentus* L.). *Pakistan J. Zool.*, **43**: 863-868.
- Dabhi, M.V.**, Koshiya, D.J. and Korat, D.M. (2013). Effect of abiotic factors on population aphid and damage by shoot and fruit borer in okra during summer and Kharif season at Anand in Gujarat. *J. Agrometeorology*, **15**(1):71-74.
- Dangi, P.R.** (2004). Incidence of *Earias vittella* (Fabricius) and its management in okra, [*Ablemoschus esculentus* (L.) Moench.] M.Sc. Thesis, Maharana Pratap University of Agriculture and Technology, Udaipur, RAJASTHAN (INDIA).
- Gautham, H.K.**, Singh, N.N. and Rai, A.B. (2014). Screening of okra against shoot and fruit borer *Earias vittella* (Fab.). *Indian J. Agric. Res.*, **48** (1) :72-75.
- Kedar, S.C.**, Kumarange, K.M., Bhujbal, D.S. and Thodsare, N.H. (2014). Insects pests of okra and their management. *Popular Kethi.*, **2**: 112-119.
- Mandal, S.K.**, Sah, S.B. and Guptha, S.C. (2006). Efficacy and economics of biopesticides and insecticides combinations against okra pests. *Internat. J. Agric. Sci.*, **2** (2) : 377-380.
- Nair, K.K.**, Ananthakrishnan, A.N. and David, D.V. (1984). *General and applied entomology*. Tata McGraw Hill Publishing Co. Ltd., New Delhi, India. 589.
- Nath, L.**, Prasad, C.S., Tiwari, G.N. and Kumar, A. (2011). Impact of weather parameters on major insect pests of okra prevailing in western Uttar Pradesh. *Vegetos.*, **24**:152-156.
- Pareek, B.L.**, Kumawat, R.L. and Patni, S.K. (2001). Effect of abiotic factors on the incidence of okra insect pests in semi-arid conditions. Proceedings of the National Conference on Plant Protection-New Horizons in the Millennium, pp: 1-8.
- Sharma, R.P.**, Swaminathan, R. and Bhati, K.K. (2010). Seasonal Incidence of fruit and shoot borer of okra along with climatic factors in Udaipur Region of India. *Asian J. Agric. Res.*, **4**(4): 232-236.
- Siddhartha, D.**, Kotikal, Y.K., Venkateshalu and Sanjiv, D. (2017). Population dynamics of fruit and shoot borers of okra in relation to climatic conditions of northern dry zone of Karnataka. *Global J. Bio-Sci. & Biotechnol.*, **6** (2): 184-190.
- Yadav, J.B.**, Singh, R.S. and Tripathi, R.A. (2007). Effect of weather parameters on incidence of pest complex of okra. *Ann. Plant Protec. Sci.*, **15**(7): 469-539.
- Yadvendu, T.C.** (2001). Evaluation of newer insecticides against insect pests of okra [*Ablemoschus esculentus* (L.) Moench]. M.Sc. Thesis, Maharana Pratap University of Agriculture and Technology, Udaipur, RAJASTHAN (INDIA).

★ ★ ★ ★ ★ 12<sup>th</sup> Year of Excellence ★ ★ ★ ★ ★