

RESEARCH ARTICLE

Seasonal variations of zooplankton distribution with relation to water quality parameters off Godavari Estuary, Andhra Pradesh

■ N. Veerabhadra Rao, N. Prasanthi, S. Bhargavi and T. V. Ramana

SUMMARY

Zooplankton abundance and diversity of zooplankton with relation to physico-chemical parameters in five stations at off Godavari estuary, Bay of Bengal during October 2010 to May 2011. A total of 29 samples were collected from the five stations, whereby 19 zooplankton groups belonging to six phylum were identified. Among the groups, Copepoda was the most dominant and abundant group which contributed 54.17%-72.73% of the total zooplankton population. Zooplankton holds a key position in the food web as it was directly related to the consumption of organic energy produced by phytoplanktonic photosynthesis and then by transforming it to the higher trophic levels of heterotrophs such as fish. This disappearance may be due to the fact that some species occur in spores, under favourable conditions spore germinate and appear as zooplankton. Plankton diversity and physico-chemical parameters of water are important criteria for evaluating the suitability of water for culture practices. Therefore, structure of different fish food organisms assumes greater significance to fisheries management.

Key Words : Water quality parameters, Zooplankton groups, Godavari estuary

How to cite this article : Veerabhadra Rao, N., Prasanthi, N., Bhargavi, S. and Ramana, T. V. (2022). Seasonal variations of zooplankton distribution with relation to water quality parameters off Godavari Estuary, Andhra Pradesh. *Internat. J. Plant Sci.*, 17 (2): 151-156, DOI: 10.15740/HAS/IJPS/17.2/151-156, Copyright@ 2022:Hind Agri-Horticultural Society.

Article chronicle : Received : 24.03.2022; Revised : 21.04.2022; Accepted : 23.05.2022

Zooplankton plays a vital role in aquatic food webs because they are important food for fish and invertebrate predators and they graze heavily on

MEMBERS OF THE RESEARCH FORUM

Author to be contacted :

N. Veerabhadra Rao, Fisheries Research Station, S.V. Veterinary University, Undi (A.P.) India
Email : narasimhaau@gmail.com

Address of the Co-authors:

N. Prasanthi and S. Bhargavi, Department of Environmental Studies, Pragati Engg. and Technology, Surampalem (A.P.) India

T.V. Ramana, Fisheries Research Station, S.V. Veterinary University, Kakinada (A.P.) India

algae, bacteria, protozoa and other invertebrates. They are too small to be important as food for most fish. They might be important in diets of some larval fish. Plankton occupies a significant and exceptional position in the biotic world for their essential role in the aquatic ecosystem. Though they are very small or tiny their absence might show the way the entire life processes in the aquatic ecosystem specially the animal life to a halt. Phytoplankton play a vital role in synthesizing light energy into food while the zooplankton are consumers of phytoplankton and these zooplankton are subsequently being eaten by other animals in their trophic

interrelationships. Zooplankton plays an indispensable role in the aquatic ecosystem. Zooplanktons are the primary consumers in the trophic level which directly or indirectly comprises major protein source of all fishes. The function of zooplankton in upwelling systems is receiving growing attention primarily because of its function in phytoplankton graze, carbon cycling and nutrient recycling. In coastal waters, zooplankton exhibit uneven distribution at scales of 10–100 km. This spatial variability reflects the complex physico-chemical and biological processes related with flow zones (Verheye *et al.*, 1992). Though significant information is available on the interaction of zooplankton with upwelling/plumes from temperate waters (Bradford *et al.*, 1993; Baduini 1997; Vargas and González, 2004; Foster and Battaerd, 1985 and James and Wilkinson, 1988), studies dealing with tropical systems continue to remain fragmented (Hitchcock *et al.*, 2002; Lo *et al.*, 2004 and Boyd and Smith, 1983). They participate an important role in the conservation of energy from primary producer (phytoplankton) to upper trophic levels. The zooplankton incidence and distribution influence pelagic fishery potentials. Thus, they are the initial prey for most fish larvae as well as for many plankton-eating adult fishes. In aquaculture sector, zooplankton is good food source for cultured fish especially fry, fingerlings and juveniles. Indeed, zooplankton is used as one of the bioindicators for accessing aquatic ecosystem health. The zooplankton are additional varied as compared to phytoplankton, their variability in any aquatic ecosystem is prejudiced mainly by roughness, diurnal vertical passage and seasons.

MATERIAL AND METHODS


The study was conducted two conjugative seasons *i.e.* from October 2010 to May 2011. The sampling stations at Godavari Estuary at 10 to 21 km distance from Yanam at 20, 30, 50, 75 and 100 m isobaths. The latitude and longitude of stations 1-5 of 16° 45' 55" N and 82° 28' 44" E. Zooplankton samples were collected using a Bongo net hauled horizontally behind the boat at 2-3 knots. The collected concentrated zooplankton samples were preserved by adding 5% formaldehyde for identification and counting (ind m⁻³) (Wickstead, 1965 and ICES, 2000). Water samples collected with a Niskin sampler from sub surface area. Water temperature was measured using a mercury-filled thermometer (Brannan 0.1°C). Salinity (ppt), (Winkler) dissolved oxygen (ppm), and inorganic nutrients (µM) nitrate and nitrite-N (Azo dye method), phosphate-P (ascorbic acid) and silicate-Si (silico-molybdic acid) were analyzed using methods in Parsons *et al.* (1984).

RESULTS AND DISCUSSION

The hydrographical parameters of zooplankton communities together form a inclusive environment and there is interaction between the zooplankton and phytoplankton. These communications are straight or indirectly subjected to the complex influences (Basawarajeshwari, *et al.*, 2015). Zooplankton was represented by Copepods, Cladocera and Gastropoda. Among the plankton Copepods was dominated and followed by Sagitta and Pteropod. In the phylum Arthropoda recorded 11 species and 3 species represented in phylum Mollusca. The second-largest

Table 1 : Hydrographical parameters at Godavari estuary during October 2010- May 2011

	Chl a	DO	Salinity	SST	Tr.	pH	NO ₃ ⁻	NO ₂ ⁻	NH ₄ ⁺	PO ₄	SiO ₄
Oct.	3.4-12.4 (6.6±3.8)	220.5-225.1 (222.6±1.8)	0.9-1.3 (1.1±0.2)	28.9-29.9 (29.5±0.4)	0.4-1.0 (0.6±0.3)	7.6-8.0 (7.8±0.2)	1.3-3.6 (2.1±1.0)	0.3-2.7 (1.05±1.06)	5.5-6.1 (5.7±0.2)	0.4-0.8 (0.6±0.1)	167.4-235.1 (207.5±27.4)
Nov.	11.3-13.5 (12.6±0.9)	260.7-288.2 (271.8±11.4)	0.9-6.3 (3.3±2.1)	29.4-29.8 (29.6±0.2)	0.6-0.9 (0.7±0.1)	8.1-8.3 (8.2±0.1)	19.1-30.4 (22.6±4.9)	0.2-0.4 (0.3±0.1)	3.8-6.7 (5.2±1.2)	0.2-0.3 (0.2±0.1)	241.8-362.9 (295.7±54.5)
Jan.11	11.2-13.8 (12.4±1.0)	207.0-303.5 (238.2±41.0)	1.8-23.9 (16.4±9.2)	27.3-27.9 (27.5±0.3)	0.7-1.2 (0.9±0.2)	8.0-8.5 (8.1±0.2)	5.5-6.2 (5.8±0.3)	0.4-1.0 (0.7±0.2)	3.0-4.8 (3.6±0.8)	0.28-0.34 (0.31±0.02)	47.7-207.3 (104.7±65.9)
Mar.09	3.6-6.5 (5.0±1.3)	144.4-214.1 (175.5±27.5)	15.0-34.5 (29.2±8.8)	30.0-30.3 (30.1±0.1)	1.2-2.1 (1.6±0.4)	8.0-8.2 (8.1±0.1)	4.3-7.7 (6.3±1.4)	0.4-0.6 (0.5±0.1)	2.5-3.3 (2.9±0.4)	0.4-0.9 (0.6±0.2)	7.5-77.5 (25.2±32.3)
Apr.	4.8-7.4 (6.3±1.0)	159.4-209.2 (177.4±21.2)	13.9-29.1 (24.6±6.7)	31.5-32.0 (31.7±0.2)	0.7-1.1 (0.9±0.1)	7.9-8.1 (8.0±0.04)	2.2-3.6 (3.0±0.5)	0.4-1.2 (1.0±0.4)	2.6-5.2 (4.3±1.1)	0.6-1.2 (0.8±0.3)	17.7-86.0 (36.7±30.5)
May	4.6-12.2 (7.7±3.1)	179.3-194.2 (188.6±6.1)	1.7-22.1 (15.3±8.5)	31.0-32.5 (31.8±0.6)	0.6-1.3 (0.9±0.3)	7.9-8.4 (8.2±0.2)	2.7-4.1 (3.4±0.5)	0.2-1.2 (0.8±0.4)	1.0-3.9 (2.3±1.1)	0.4-0.7 (0.5±0.1)	14.2-222.2 (72.2±92.8)



phylum of invertebrate is Mollusca after the Arthropoda. During the study period quantitative and qualitative variations of zooplankton was observed (Table 1). Highest Zooplankton groups observed in the pre-monsoon periods whereas lowest number in the post monsoon period (Fig.1). Similar results observed by (Mukherjee, 2011). In the ecosystem, zooplankton plays a main role as they consume the phytoplankton and form a major food source

for tertiary producers.

The zooplankton assemblage inhabiting brackish water, mostly comprises representative of Arthropoda and Mollusca groups. The zooplanktons frequently respond straight away to environmental changes because most of the species have short production times. Zooplankton considered as the basic principle natural fish food for young and some adults of organisms, which hold

Table 2 : Zooplankton abundance (Mean, ind m⁻³) at off Godavari estuary during October 2010- May 2011

Phylum	Name	Post-monsoon Oct., Nov. and Jan	Pre-monsoon March, Apr. and May	
Annelida	Polychaete larvae	0.1	0.3	
Arthropoda	Copepod	132.5	248.5	
	Amohipod	0.1	0.4	
	Cladocera	5.1	0.0	
	Euphasid	0.0	0.8	
	Lucifer	0.3	0.8	
	Megalopa larve	0.0	0.1	
	Mysid	0.4	0.0	
	Phillosoma	0.4	0.0	
	Zoa larvae	0.1	1.6	
	Mysis	0.0	1.1	
	Ostracoda	2.7	0.3	
	Mollusca	Gastropoda	3.3	4.4
		Pteropod	1.0	7.1
Bivalves		0.0	1.0	
Echinodermata				
Chaetognatha	Sagitta	2.1	8.8	
Chordata	Oikoplura	0.6	3.5	
	Fish eggs	1.4	4.3	
	Fish larvae	0.1	0.3	

fish production (El-Serafy *et al.*, 2009). Epifanio and Garvine (2001) studied by the variation of their spatial distribution, based on different factors. The higher population density of the zooplankton is during the pre-monsoon period while low population density observed in the post monsoon. Among the plankton copepod is dominated. The zooplankton populations dominated by copepods in the dry season are observed by Egborge (1981). The high population density in the pre-monsoon period may be as a result of abundant food sources from the runoff. The increase of primary production (phytoplankton) is accompanied by increase in zooplankton abundance reported by Rocha *et al.* (1999). Muylaert *et al.* (2003) observed that the zooplankton abundance frequently reach their peak during the dry season in the estuarine ecosystem. Besides food source, low predation rate by fish during wet season caused by plankton growing activity, could support by high density of zooplankton (Ikpi *et al.*, 2013). Higher zooplankton population density in summer might be due to the temperature acceleration in the Godavari estuary (Ikpi *et al.* (2013) reported that the seasonal variation in zooplankton condition could larger be due to the copepoda which normally constitute major food items of larger zooplankton. The air temperature ranged from 25.8°C - 32.5°C and highest temperature recorded in the month of May, 2011 (Fig.2). The one of the important environment factor is temperature, since it is influencing the distribution of plankton. The India in influenced by two monsoon *i.e.* south west monsoon and northwest monsoons.




Fig. 2 : Temperature (Mean) at off Godavari estuary during post-monsoon and pre-monsoon 2011

The lowest value of reading observed during summer (October and November, 2010). The low species diversity was observed post monsoon season which could

be attributed to heavy freshwater influx and low salinity (Godhantaraman, 1994). The turbidity during this season may also responsible for lower values.

Dissolved oxygen was observed in between 44.4 to 303.5 µM and lowest recorded in month of March, 2011. Lower dissolved oxygen values occurred during dry season could maybe due to higher temperatures during summer months. Dissolved oxygen distribution provides a good index of productivity and quality of surroundings. Consumption of high oxygen indicative of higher photosynthetic efficiency and plankton production. Rajgopal (2010) reported that the abundance of phytoplankton which enriched water with high dissolved oxygen during photosynthesis. Dissolved oxygen is a sole of the physico-chemical parameters of the water which need to keep the organisms alive and health of the water body of ecosystem (Madhusudhana and Krishna, 2013).

The pH value ranged between 7.6-8.5 lowest recorded in the month of October whereas highest recorded in the month of January (North East monsoon). Indication of low level of water and high photosynthesis resulting in high production of free CO₂ during the equilibrium towards high value of pH (Siddamalayya and Pratima, 2008). In the present, study months of January and November goes to above 8.5 pH which indicates that the water is a highly production of zooplankton population. In the environment inorganic nitrogen occurs in a range of oxidation status as Nitrate (NO₃) and Nitrite (NO₂), the ammonium ion (NH₄⁺) and molecular Nitrogen (N₂). It is under goes biological and non-biological transmission in environment as part of nitrogen cycle. In the present study Nitrite in between 0.1-2.7 µM; Nitrate in-between 1.3 to 30.4 µM and Ammonia goes to 1.0 to 15.3 µM in the study period.




Fig. 3 : Seasonal observations of nutrients (NH₄, NO₂, NO₃ and PO₄) during post-monsoon and pre-monsoon 2011

Changes in water quality of water body have major effect on structure of zooplankton assemblages that can potentially affect the functioning of ecosystem (Sousa *et al.* (2008). Seasonal distribution of the population structure of zooplankton in connection with physico-chemical parameters Sarkar and Chaudhary (1999). Hence, Zooplankton communities of numerous water bodies have been used as indicators for the status of the ecosystem (Jeppensen *et al.*, 1999; Ramchandra and Solanki, 2007) and related with the concentration of total nitrogen, total phosphorus, algal biomass and the density and size of individuals (Giselle and Bruce, 2007). The inconsistency observed in the movement of zooplankton is due to abiotic parameters. In the aquatic ecosystem plankton play a critical role not only in converting plant food to animal food but also serves as source of food for their organisms (Rajashekhar *et al.*, 2010). In the present study observed that a total of 16 zooplankton species were recorded comprising of 9 rotifera, 3 cladocera and 4 copepods. Copepoda The present preliminary study conducted that the various zooplankton composition. Copepoda constitute higher species abundance, the overall diversity index shows eutrophic nature. Further, detailed investigation through regular monthly sampling with more quantitative analysis to conform the exact status of water body is required which would help to conserve the zooplankton diversity and water quality.

REFERENCES

- Baduini, C.L. (1997). Spatial and temporal patterns of zooplankton biomass in Monterey Bay, California, during the 1991–1993 El Nino and an assessment of the sampling design. *CalCOFI Report.*, **38** : 193–198.
- Boyd, C.M. and Smith, S.L. (1983). Plankton, upwelling and coastally trapped waves of Peru. *Deep-Sea Res.*, **30** : 723–742.
- Bradford, J.M., Murdoch, R.C. and Chapman, B.E. (1993). Composition of macrozooplankton assemblages associated with the formation and decay of pulses within an upwelling plume in greater Cook Strait, New Zealand. *N Z J Mar Freshwater Res.*, **27**:1–22.
- El-Serafy, S.S., Mageed, A.A. and EL-Enany, H.R. (2009). Impact of food, water on the distribution of zooplankton in the main channel of Lake Nasser, Egypt. *J. Egypt. Acad. Soc. Environ. Delop.*, **10** : 121-141.
- Egborge, A.B.M. (1981). The composition, seasonal variation and the distribution of zooplankton in LakeAsejire, Nigeria. *Revue de Zoologie Africaine*, **95** : 136-144.
- Epifanio, C.E. and Garvine, R.W. (2001). Larval transport on the Atlantic continental shelf of North America. *Estu. Coast. Shelf. Sci.*, **52** : 51-77.
- Foster, B.A. and Battaerd, W.R. (1985). Distribution of zooplankton in a coastal upwelling in New Zealand. *N. Z. J. Mar. Fresh Water Res.*, **19** : 213–226.
- Giselle, V.T. and Bruce, R.F. (2007). Relationships among nitrogen and total phosphorus, algal biomass and zooplankton density in the central Amazonia lakes. *Hydrobiologia*, **595** : 177-195.
- Godhantaraman, N. (1994). Species composition and abundance of tintinnids and copepods in the Pichavaram mangroves (South India). *Ciencias Marinas*, **20** : 371-391.
- Hitchcock, G.L., Lane, P., Smith, S., Luo, J. and Ortner, P.B. (2002). Zooplankton spatial distributions in coastal waters of the northern Arabian Sea, August 1995. *Deep-Sea Res.*, **49** (2) : 2403–2423.
- Ikpi, G.U., Offem, B.O. and Okey, I.B. (2013). Plankton distribution and diversity in tropical earthen ponds. *Env. Nat. Resou. Res.*, **3** (3): 45-51.
- James, M.R. and Wilkinson, V.H. (1988) Biomass, carbon ingestion, and ammonia excretion by zooplankton associated with an upwelling plume in western Cook Strait, New Zealand. *N.Z.J. Mar. Fresh Water Res.*, **22** : 249–257.
- Jeppensen, E., Jensen, J.P., Sondergaard, M. and Lauridsen, T.L. (1999). Trophic dynamics in turbid and Clearwater lakes with special emphasis on the role of zooplankton for water clarity. *Hydrobiologia*, **408/409** : 217-231.
- Lo, W., Chung, C. and Shih, C. (2004) Seasonal distribution of copepods in Tapong Bay, Southwestern Taiwan. *Zool Stud.*, **43** : 464–474.
- Madhusudhana, R.K. and Krishna, P.V. (2013). Seasonal variations in hydrographic status of Interu mangrove swamp of River Krishna estuarine region, Andhra Pradesh, India. *Bioinfo Aquatic Ecosystem*, **2** (1) : 43 -50.
- Mukherjee, P. (2011). Stastical analysis of biodiversity of zooplankton population in a filthy Trapa-cum-Fish cultured pond of central India. *Int. J. Zool. Res.*, **1**(2): 24-29.
- Muylaert, K.S., Declerck, V., Geenens, J.V., Wichelen, H., Deegans, J., Vandekerkhove, K.V. and Vyverman, W. (2003). Zooplankton, phytoplankton and the

- microbial food web in two turbid and two clear shallow lakes in Belgium. *Aquat. Ecol.*, **37** : 137-150.
- Rajagopal, T., Thangamani, A. and Archunan, G. (2010). Comparison of physico-chemical parameters and phytoplankton species diversity of two perennial ponds to Sattar area, Tamil Nadu. *J. Env. Biol.*, **31**: 787 - 794.
- Ramchandra, T.V. and Solanki, M. (2007). *Ecological assesment of lentic water bodies of Bangalore*. Envis Technical Report: 25. Indian Institute of Science, Bangalore.
- Rocha, O., Matsumura-Tundisi, Y.T., Espindola, E.L.G, Roche, K.F. and Rietzler, A.C. (1999). Ecological theory applied to reservoir zooplankton, In: Tundisi JG, Straskraba M (eds) *Theoretical reservoir ecology and its applications*. International Institute of Ecology, Brazilian Academy of Sciences. Backhugs Publishers, Leiden, Holland, pp. 29-51.
- Sarkar, S.K. and Chaudhary, B. (1999). *Role of some environmental factors on the fluctuations of plankton in a lentic pond at Calcutta*. Limnological research in India. Daya Publishing House. pp. 108-130.
- Siddamalayya, N. and Pratima, M. (2008). Impact of domestic sewage on freshwater body. *J. Env. Bio.*, **29**: 303- 308.
- Sousa, W., Jose, L., Attayde, Elinez Dasilva Rocha and Eneida Maria and Eskinazi Santanna (2008). The response of zooplankton assemblages to variations in the water quality of four man-made lakes in semi-arid northeastern Brazil. *J. Plankton Research*, **30** (6) : 699- 708.
- Vargas, C.A. and González, H.E. (2004). Plankton community structure and carbon cycling in a coastal upwelling system. II. Microheterotrophic pathway. *Aquat. Microb. Ecol.*, **34** : 165–180.
- Verhey, H.M., Hutchings, L., Huggett, J.A. and Painting, S.J. (1992). Mesozooplankton dynamics in the Benguela ecosystem, with emphasis on the herbivorous copepods. *South African J. Mar. Sci.*, **12** : 561–584.
- Wickstead, J.H. (1965). *An introduction to the study of Tropical Plankton*. Hutchinson and Co., London

17th
Year
★★★★★ of Excellence ★★★★★