Visit us: www.researchjournal.co.in

RESEARCH PAPER

■ ISSN: 0973-130X

Direct seeding of rice with drum seeder is made easy rice cultivation in Visakhapatnam district of North Coastal Zone of Andhra Pradesh

D. Uma Maheswara Rao, P. B. Pradeep Kumar **and** K. Tejeswara Rao*
District Agricutral Advisory and Transfer of Technology Centre, Kondempudi, Visakhapatnam (A.P.) India
(Email: tejaseniorscientist@gmail.com)

Abstract : Paddy is major predominant crop during *Kharif* in Visakhapatnam district of Andhra Pradesh, cultivated in an area of 1,17,608 ha, out of total cropped area of 1,92,638 ha with productivity of 2664 kg/ha. Farmers grow crop by adopting traditional method of paddy cultivation, use more seed rate, close spacing, late transplanting with over aged seedlings common phenomenon due to erratic rainfall climate change. Scarcity of labour and escalation in labor wages, reduction in labor efficiency are leading to low net returns. In this context DAATT Centre, Visakhapatnam district of ANGRAU, in collaboration with K.V.K., Kondempudi and Department of Agriculture, Visakhapatnam has introduced 'Drum Seeder technology which is very easy and less labour involved. Drum seeder technology is boon to farmers to save money and time and crop comes to harvest 7-10 days earlier than normal transplanted paddy. DAATT Centre, Visakhapatnam has organized On-Farm Demonstrations (OFDs) in farmer fields in two seasons *Kharif*, 2019 and *Rabi*, 2019-20. Drum seeder technology in paddy recorded 12.49% yield over normal transplanting method of paddy cultivation during *Kharif* season. The results from the study showed that the farmers realized the 58.19% increase in net income due to increased grain yield by 12.49% with reduction of cost of cultivation by 9.35%, , it could be attributed to reduction in manual labour of 12 man labour and 25 women labour per ha and also increase in yield attributes and yield.

Key Words: Paddy drum seeder, Conoweeder, OFT, Yield and yield attributes, B:C ratio

View Point Article: Uma Maheswara Rao, D., Pradeep Kumar, P. B. and Tejeswara Rao, K. (2022). Direct seeding of rice with drum seeder is made easy rice cultivation in Visakhapatnam district of North Coastal Zone of Andhra Pradesh. *Internat. J. agric. Sci.*, **18** (2): 661-665, **DOI:10.15740/HAS/IJAS/18.2/661-665.** Copyright@ 2022: Hind Agri-Horticultural Society.

Article History: Received: 05.03.2022; Revised: 11.04.2022; Accepted: 13.05.2022

Introduction

Rice area has been decreasing in state like Andhra Pradesh, although overall productivity is increasing, there is a decrease in compound growth rate in rice productivity at national level (Krishnaiah, 1999). There is no scope for expansion of area for rice cultivation. Rice yields are plateau in the irrigated ecosystem and the rainfed

system with low productivity of 2.5 to 3.5t ha⁻¹, it has become imperative to increase rice production per unit area per unit time to feed the teaming millions in the new millennium. India has to produce 135-145 million tones by 2020 A.D. to feed the additional 350million people. To do so, the productivity should be raised to 3.2t ha⁻¹ a by 2020AD from the present level of 2.05t ha⁻¹ (The Hindu Survey of Indian Agriculture, 2006).

^{*}Author for correspondence:

Visakhapatnam district is the one of the rice growing districts in Andhra Pradesh. Farmers grow rice in 1.17 lakh ha during Kharif and 2000 ha during Rabi. Cost of cultivation is rising year by year due to many reasons i.e. social reasons, situational factors and input cost. At present cost of cultivation per hectare is between Rs.34,250/- to Rs. 38,500/-. This is mainly due to escalation of labour wages and scarcity of labour in villages during agricultural season, labour requirement is very intense at the time of transplanting season and increase in price of fertilizers.

To overcome this, wet paddy seeding with drum seeder can reduce the labour requirement during transplanting season, technology is very simple and can be adopted by the farmers easily, cost of cultivation can be reduced and receive quality crop harvest at 7-10 days earlier than normal transplanted field. Paddy drum seeder technology holds special significance in the present day production system with regard to saving labour component by 30-50% and increase the productivity by 20-30% (Annual Reports of Directorate of Rice Research, 2003, Rajendranagar, Hyderabad). Paddy drum seeder technology, a new way of cultivation is gaining more attention of the farmers in Visakhapatnam district. But the paddy drum seeder technology has its own implications to adopt such as right choice of field, irrigation facilities, varieties, land preparation, weed management and machinery available.

The prime concern of any programme related to agriculture is to enhance productivity and with reduced cost. In order to reduce the cost of cultivation DAATT Centre, Visakhapatnam took initiatives in collaboration with KVK, Kondempudi to promote direct seeding of paddy with drum seeder in the district the following objectives.

Objectives:


- To test the feasibility of paddy drum seeder technology in the Visakhapatnam district through On Farm Trials (OFTs), method demonstrations
- To record the yield in paddy drum seeder technology in comparison with normal method to convince the farmer.
- To analyze the economics of paddy cultivation in Visakahpatnam district.

MATERIAL AND METHODS

Scientists in KVK and DAATT Centre,

Visakhapatnam district of ANGRAU in collaboration with Department of Agriculture of the district has introduced paddy drum seeder technology with comparing normal method of cultivation through organizing On-Farm Trials (OFTs) during Kharif, 2019 and Rabi, 2019-20 in 8 locations.

Flow chart of handed over of paddy drum seeder technology to the farmers:



Details and advantages of direct paddy seeder technology:

Direct Paddy Drum Seeder (Fig. A) is small equipment for sowing germinated paddy seed directly in wetland field is fabricated and it is used for demonstration. There is no need for transplantation. It is a manually pulled implement. It covers 8 rows of 20 cm row-to-row spacing at a time. It is made up of plastic materials. Cono weeder (Fig. B) is used for intercultivaton to incorporate weed in inbetween rows apart of 20cm.




Fig. A: Drum seeder



Fig. B : Conoweeder

Salient features:

- Labour cost is reduced drastically
- Cost of cultivation is reduced because, cost on nursery raising, nursery pulling and transplanting can be saved
 - Uniformity in seed sowing and Plant population
 - Continuous drilling of seeds is eliminated.
- Reduction in seed rate and thinning cost. Crop matures 7-10 days earlier than the transplanted paddy Light in weight and easy to handle
 - An area of 1 hectare per day can be shown
- Saving in seed requirements: 12-18 kg per acre is sufficient depending on variety

Farmer fields are selected to conduct On Farm Trials (OFTs) with proper drainage facility and regulation of water. The varieties cultivated in paddy drum seeder technology were in Kharif with MTU-1001, in Rabi with MTU-1010.

Since planting of crop in both drum seeder technology and normal transplanting method data pertaining to crop stand with number of tillers per hill, number of tillers per square meter, incidence of pests and diseases if any at regular intervals followed by yield contributing parameters like effective tillers, panicles per hill and square meter, length and texture of panicle, number of grains per panicle and 1000 grain weight are recorded. Yield per 5x5m² was collected and calculated per hectare area.

Means of yield attributes, yield and cost of cultivation were arrived for yield in both drum seeder technology and normal transplanting methods. Percentage yield increase over normal method was calculated and comparative analysis of cost benefit ratio per hectare was arrived and presented in the tables.

RESULTS AND DISCUSSION

The On-Farm Demonstrations on drum seeder technology is conducted for two seasons both in *Kharif*, 2019 and Rabi, 2019-20 in innovative farmer fields and yield attributes and yield are depicted in following tables.

Crop duration:

The duration of the crop (Table 1) is recorded, it was observed that there is reduction of duration to reach the harvesting of rice crop was observed. The reduction of crop duration 9 days was observed when compared to normal transplantation method of rice cultivation. The results are corroborates with the Chandrasekhara Rao et al. (2013).

Yield attributes:

All yield attributes, during all the years were recorded better in drum seeder technology than normal method (Table 1). During all the seasons of three years of demonstrations and in all the locations with different varieties recorded the number of productive tillers per/ Sq.mt and number of grains per panicle are 229 and 192 are more than normal practice 201 and 176 (MSSRF, 2002), respectively.

Yield:

Grain yield (Table 1) increase was achieved to a tune of 11.39% in drum seeder method (6245 kg ha⁻¹) over normal method of cultivation (5604 kg ha⁻¹). The use of drum seeder is superior to broad casting method of raising the rice crop. Higher yield in drum seeder

Table 1: Mean data on yield and yield attributes of on-farm demonstrations on paddy drum seeder technology conducted during Kharif-2019 and Rabi, 2019-20

Season	No.of locations	Duration		No.of productive tillers/Sq.mt		No.of grains/panicle		Yield Kg/ha		Percentage increase over	
		Demo	Check	Demo	Check	Demo	Check	Demo	Check	check	
Kharif- 2019	3	124	134	227	190	213	198	6580	5680	15.85	
Rabi- 2019-2020	5	113	121	230	212	171	153	5910	5528	6.92	
Average		119	128	229	201	192	176	6245	5604	11.39	

Table 2: Mean of comparative analysis of cost of cultivation including labour per hectare of paddy recorded during *Kharif-*2019 and *Rabi*, 2019-20

		Paddy drum sæder technology					Conventional transplantation method				
Sr. No	Practice	Labour cost		Input cost		Total	Labour cost		Input cost		Total
	T ractice	No	Cost	Input	Cost Rs.	cost in Rs./ha	No.of labour	Cost Rs/ha	Input	Cost Rs.	cost in Rs./ha
1.	Nursery management	0	0	0	0	0	2 M	600	FYM	500	1100
							preparation				
2.	Seed quantity and cost	0	0	40kg/ha	1200	1200	0	0	75kg/ha	2250	2250
				@Rs 30					@R30		
3.	Land preparation of	10 M	6600	0	0	6600	8M and	5600	0	0	5600
	main field	Pudlling					puddling				
4.	Nursery	0	0	0	0	0	15M	4950	0	0	4950
	pulling and spreading						3 W				
5.	Transplanting/sowing	5M 3W	2000	0	0	2000	25W	3750	0		3750
6.	Manures and	8M	2400	DAP-125kg	8000	10400	8M	2400	DAP-125	8000	10400
	fertilizers management /ha			urea 150 kg					kg		
				Mop :85 kg					urea 150		
									kg		
									Mop :85		
									kg		
7.	Weeding and herbicide	3M	4000	Herbicides	2500	6500	1M	3300	Herbicide	1000	4300
	inter-cultivation	20W					20W				
8.	Plant protection	2M	1000	PP	3500	4500	4M	1800	PP	4500	6300
		2W		chemicals			4W		chemicals		
9.	Irrigation management	8 M	2400	0	0	2400	10M	3000	0	0	3000
10.	Harv esting	27 W	5400	0	0	5400	25 W	5000	0	0	5000
11.	Threshing, winnowing and	16M	8000	0	0	8000	16M	8000	0	0	8000
	bagging	16W					16W				
	Total	52 M	31800		15200	470 00	64M	38400		13500	51850
		68 W					93W				

M-Male F-Female

Table 3:	Table 3: Economics of the paddy drum seeder technology vs normal transplantation method recorded during Kharif-2019 and Rabi, 2019-20								
Sr. No.	Particulars	Paddy drum seeder technology method	Transplantation method	Difference					
1.	Grain yield kg/ha	6580	5680	900					
2.	Straw yield kg/ha	7370	6362	1008					
3.	Grain value (Rs. 14/kg)	92120	79520	12600					
4.	Straw value (Rs.0.5/kg)	3685	3180	504					
5.	Gross income Rs./ha	95805	82700	13104					
6.	Total cost of cultivation Rs/ha	47000	51850	-4850					
7.	Net income Rs./ha	48805	30851	17954					
8.	C:B ratio	2.03	1.59	0.44					

technology is contributed by more number of productive tillers by supported by profuse root system resulted in more number of panicles.

Labour involvement in rice cultivation:

There is reduction in laobur utilization (Table 2) is observed in drum seeder technology i.e., 12 man labour and 25 women labour when compared to normal method of transplanting method.

Economics:

Additional grain yield and straw yield (Table 3) of 900 kg ha⁻¹ and 1,008 kg ha⁻¹ recorded in drum seed technology compared with normal practice of transplantation, this could be due to uniform plant population, good tillering capacity. Additional net income of Rs.15,404 ha⁻¹ received in drum seeder technology with reduction of cost of cultivation of Rs.2,300 ha⁻¹ realized over normal transplanting. The net income increased by Rs. 17,954 per hectare in drum seeder technology. It was mainly due to the reduction in cost of sowing operation and transplanting operations. It was observed that the cost-benefit ration was higher in direct sowing (2.03) which is significantly higher than in conventional method (1.59).

Conclusion:

The results from the study showed that the farmers

realized the 58.19% increase in net income due to increased grain yield by 12.49% with reduction of cost of cultivation by 9.35%, it could be attributed to reduction in manual labour of 15 man labour and 27 women labour and also increase in yield attributes and yield (Paladugu et al., 2004).

REFERENCES

Annual reports of Indian Institute of Rice Research, 2003, Rajendranagar, Hyderabad.

Chandrasekhararao, C., Jintendra, S. and Murthy, T.G.K. (2013). Resource optimization in rice through direct seeding by drum seeder. Internat. J. Agric. & Food Sci. Technol., 4: 239-246.

Krishnaiah, K. (1999). Rice production in India-CREMNET status and future concerns. Paper presented at 2nd CREMNET Workshop-cum-Group Meeting, 24-27, August 1999, SWMRI, Thanjavur, India.

MSSRF (2002). The system of rice intensification-SRI, In: Updates on SRI activities/Progress around the world, CIIFAD Report, April, 2004. In: http://ciifad.cornell.edu/sri

Paladugu, S., Thati, S., Lekhi, M., Yadla, S., Poli, R. and Alapati, S. (2004). Studies on varietal performance under SRI and non-SRI. World Rice Research Conference, Nov (5-7), 2004 HELD AT Tsukuba International Congress Centers (Epochas, Tsukuba), Tsukuba, Ibaraki, Japan Section-26: pp. 553.

The Hindu Survey of Indian Agriculture (2006). pp.5-54.

