

Influence of cucumber mosaic virus on growth, moisture and dry matter content of chilli (*Capsicum annum* L.)

HENA, K.D. DWIVEDI, N.A. ANSARI AND J.P. TEWARI

Accepted : August, 2008

See end of the article for authors' affiliations

Correspondence to:

J.P. TEWARI

Department of Botany,
M.L.K. (P.G.) College,
BALRAMPUR (U.P.)
INDIA

ABSTRACT

A study was made on the effect of two chilli strains (severe and mild) of cucumber mosaic virus infections on the growth, moisture and dry matter content in chilli. The infection affected the growth of the plants adversely. More dry matter and less moisture contents were found in diseased plants as compared to healthy plants. The effect was more pronounced with severe strain.

Key words : Metabolic activity, CMV, Chilli.

Chilli (*Capsicum annum* L.), also called red pepper is an important cash crop in India and grown for its pungent fruits which are commonly used both green and ripe. Chilli mosaic disease of chilli caused by cucumber mosaic virus (Anjaneyulu and Appararao, 1967; Pandey *et al.*, 2004). Most of the investigation, done in the past on this disease were confined to the characterization of the virus (Mishra, 1963; Rao *et al.*, 1970) and so far no study has been made on the physiology of infected plants to understand the effect of virus infection on the metabolism of host. In the present study, two locally collected chilli strains (mild and severe) were selected to see their effect on growth, moisture and dry matter content in popular "pusa jwala" variety of chilli.

MATERIALS AND METHODS

Observation were made after 10, 20, 30, 40 and 50 days of inoculation. Three lots of 25 chilli plants were taken, first and second lots were inoculated with mild and severe strains of the virus, respectively. While the third lot was kept as healthy (control).

Seedlings were 15 days old at the time of inoculation. The growth was measured in centimeter. The moisture content of root, stem and leaves of healthy and diseased chilli plants were determined. Root, stem and leaves from five healthy and five diseased plants were collected separately in polythene bags and weighed to determine the fresh weight. These were then transferred to an oven and dried at 65°C till a constant weight was obtained. The differences between fresh and dry weight was taken as the moisture content. The moisture content thus obtained has been expressed as per cent moisture on fresh weight basis.

Average dry weight of root, stem and leaves from

five plant were taken per treatment at each interval and evaluated as per cent dry matter content on fresh weight basis.

All experiments were carried out in an insect proof chamber the data were analysed statistically using the methods given by Chandel (2004).

RESULTS AND DISCUSSION

The perusal Table 1 show that both the strains reduce the growth of chilli plants, the severe strains induces more adverse effect than the mild one. When statistically analysed these data are significant at 5 per cent level. The perusal of data (Table 2 and 3) show that the infection had influenced the dry matter content and moisture level

Table 1 : Effect of severe and mild strain of cucumber mosaic virus on the growth of chilli plants

Days after Inoculation	Healthy plant	Plant height in (cm.) (Average)	
		Severe strain	Mild strain
10	7.0	4.0	5.5
20	10.0	7.0	8.0
30	15.0	9.0	11.0
40	21.5	12.0	16.0
50	33.0	17.0	21.5
Average	17.3	9.8	12.4
S.E.±	4.15	1.99	2.57
C.D. (P=0.05)	8.84	4.24	5.47

of the host plant. Root, stem and leaf samples taken from infected plants had less moisture content and more dry matter than their comparable healthy counterparts on fresh weight basis but the moisture content whether taken from healthy or infected samples increased gradually with

Table 2 : Moisture content (per cent) of plant parts of chilli infected with two strains of cucumber mosaic virus

Days after inoculation	Root			Stem			Leaf		
	Healthy	Infected		Healthy	Infected		Healthy	Infected	
		Severe	Mild		Severe	Mild		Severe	Mild
10	62.3	58.3	60.5	62.5	59.0	61.2	63.0	61.0	62.0
20	56.9	62.3	63.2	65.0	62.5	63.4	66.0	62.0	62.5
30	69.8	65.2	67.0	69.5	66.0	67.8	69.5	64.5	65.0
40	72.7	67.8	68.6	71.4	66.8	70.0	73.8	65.2	69.0
50	75.0	70.5	71.8	77.8	73.2	74.5	75.0	70.2	72.0
Average	67.34	64.82	66.22	69.24	65.5	67.38	69.46	64.58	66.1
S.E. \pm	3.02	1.94	1.78	2.38	2.14	2.11	2.03	1.43	1.72
C.D. (P=0.05)	6.43	4.13	3.79	5.07	4.51	4.49	4.32	3.04	3.66

Table 3 : Dry matter content (Per cent of fresh weight) of chilli infected with two strains of cucumber mosaic virus

Days after inoculation	Root			Stem			Leaf		
	Healthy	Infected		Healthy	Infected		Healthy	Infected	
		Severe	Mild		Severe	Mild		Severe	Mild
10	39.5	41.8	41.0	39.4	41.5	40.5	37.0	39.0	37.8
20	35.3	39.7	38.7	37.0	29.7	38.0	33.6	38.0	37.2
30	31.8	36.2	35.0	32.1	36.2	39.0	31.6	34.5	35.0
40	29.2	33.3	32.8	30.5	35.2	34.0	26.2	34.3	31.0
50	26.5	30.5	29.0	24.5	28.5	27.5	25.0	28.2	27.5
Average	32.46	36.3	35.3	32.7	34.22	34.4	30.68	34.8	33.7
S.E. \pm	2.04	1.84	1.90	2.33	2.11	2.04	2.02	1.61	1.75
C.D. (P=0.05)	4.34	3.92	4.85	4.96	4.49	4.34	4.30	3.43	3.72

the age of the plant, while the dry matter content indicated a reverse trend calculated on fresh weight basis. All these data were found to be significant at 5 per cent level when analyzed statistically.

There are various reports on the reduction of growth in plants due to virus infection (Caldwell, 1934; Jeyarajan and Ramakrishnan, 1961; Srivastava *et al.*, 2005; Dubey, 1972; Srivastava, 1971, 1978 and Suteri, 1974).

In this study the reduction in growth of chilli plants has been recorded as a result of virus infection. No precise cause can be given as to what exactly contributed to this stunting but there are reports available which show disturbed auxin metabolism in virus infected plants (Grieve, 1936; Smith *et al.*, 1968).

Many workers have reported the effect of virus infection on dry matter and moisture content of plants infected with different viruses (Campbell, 1925; Dubey, 1972; Srivastava, 1971, 1978 and Suteri, 1974).

In the present study it was observed that root, stem and leaves samples taken from infected plants had less moisture content invariably (Table 2) and more dry matter content (Table 3). Moisture content of root, stem and leaf increased continuously with age of the plant in both healthy and infected samples. (Grieve, 1936; Smith *et al.*, 1968; Campbell, 1925; Caldwell, 1934; Srivastava, 1971,

1978 and Srivastava *et al.*, 2005) The changes in the dry matter and moisture content was pronounced in plants infected by severe strain virus as compared to mild strain of CMV.

Authors' affiliations:

HENA, K.D. DWIVEDI AND N.A. ANSARI, Plant Pathology Laboratory, Department of Botany, M.L.K. (P.G.) College, BALRAMPUR (U.P.) INDIA

REFERENCES

- Anjaneyulu, A. and Appa Rao, A. (1967)**. Natural occurrence of cucumber mosaic virus on chilli in India. *Indian Phytopath.*, **20** : 380-381.
- Caldwell, J. (1934)**. The physiology of virus disease in plants VI. Some effects of mosaic on the metabolism of the tomato. *Ann. Appl. Biol.*, **21** : 206-224.
- Campbell, E.G. (1925)**. Potato root as affecting the carbohydrate, water and nitrogen content of the host. *Phytopathology*, **15** : 427-430.
- Chandel, S.R.S. (2004)**. *A Hand Book of Agricultural Statistics*. Achal prakashan mandir Kanpur, India.
- Dubey, L.N. (1972)**. Some studies on metabolism and histology of virus infected chilli (*Capsicum annum* L.) Ph.D. Thesis, Gorakhpur University, Gorakhpur.

- Greive, B.J. (1936).** Spotted wilt virus and the hormones heteroauxin. *Nature*, **138** : 129.
- Jeyarajan, R. and Ramakrishnan, K. (1961).** Studies on a virus disease of chilli (*Capsicum* sp.) *South Indian Hort.*, **9** : 1-12.
- Mishra, M.D. (1963).** Host range and physical properties of a virus causing mosaic symptoms and necrosis on chillies (*Capsicum annum* L.). *Indian J. Microbiol.*, **3** : 77-84.
- Pandey, A., Ansari, N.A., Tewari, J.P. and Srivastava, G.P.(2004).** Screening of chilli varieties against mosaic disease *J.Liv-world*, **11** : 24-26.
- Rao, K.N. Apparao, A. and Reddy, D.V.R. (1970).** A ring spot strain of potato virus x on chilli (*Capsicum annum*). *Indian Phytopath*, **23** : 69-73.
- Smith, S.H., Mecal, S.R. and Harris, J.H. (1968).** Auxin transport in curly top virus infected tomato *Phytopathology*, **58** : 1669-1670.
- Srivastava, G.P. (1971).** Studies on the metabolic activities of virus infected sugarcane Ph.D. Thesis. Gorakhpur University Gorakhpur.
- Srivastava, S.B. (1978).** Studies on two mosaic disease of Brinjal (*Solanum melongena* L) Ph.D. Thesis, Gorakhpur university Gorakhpur.
- Srivastava, A.K., Shukla, A., Srivastava, G.P., Ansari, N.A., Mishra, V.A. and Tewari, J.P. (2005).** Influence of a mosaic virus on chlorophyll content of chilli (*Capsicum annum* L.) *Vegetos*, **18** : 161-162.
- Suteri, B.D. (1974).** Studies on mosaic of soyabean and metabolic changes in infected plants. Ph.D. Thesis, Gorakhpur University Gorakhpur.
