

Evaluation of different fungicides and biopesticides against Sclerotinia blight of brinjal (*Solanum melongena* L.)

RAMESH SINGH, P.C. SINGH, NARENDRA SINGH AND ALKA

Accepted : September, 2008

See end of the article for authors' affiliations

Correspondence to:

NARENDRA SINGH
Oil-seed section, C.S.A.
University of Agriculture
and Technology,
KANPUR (U.P.) INDIA

ABSTRACT

Sclerotinia blight caused by *Sclerotinia sclerotiorum* is an important disease of brinjal (*Solanum melongena* L.) in eastern U.P. Efficacy of fungicides, plant extracts and biopesticide were tested *in vitro* and *in vivo*. Bavistin (0.1%), Vitavax (0.1%), and Topsin-M (0.15%) proved to be the most effective in inhibiting the growth of pathogen *in vitro* and controlling the diseases in the field. Biopesticide, Nimbidine was also proved effective, but slightly less effective than systemic fungicide except Chlorothalonil. Nimbidine being a safe eco-friendly and economical bioproduct which can be used in the management of the disease.

Key words : Brinjal , *Solanum melongena* L. Sclerotinia blight, *Sclerotinia sclerotiorum*, Management, Fungicides and biopesticide.

Brinjal or egg plant (*Solanum melongena* L.) is one of the most common, popular and principal annual vegetable crops grown in all the three seasons under irrigated conditions. Brinjal was observed to suffer from Sclerotinia blight caused by *Sclerotinia sclerotiorum* (L.) de Bary, a most important and severe disease in eastern part of U.P. Not much work appears to have been done on this disease except the report of its occurrence. Recently, the disease has assumed the serious proportion damaging the crop to the tune at 20-40 per cent. Therefore, it was felt necessary to explore the possibility for the management of this disease through the use of fungicides, biopesticide and plant extracts.

MATERIALS AND METHODS

The efficacy of fungicides, biopesticide and plant extracts against the pathogen *in vitro* was tested by poison food technique as suggested by Schmitz (1930) using PDA medium. Bavistin, Vitavax, Blitox-50, Chlorothalonil, Kitazin, Ridomil, Zineb, Captafol, Indofil M-45, Captan, Topsin-M, one antibiotic Streptocycline (0.05%), Nimbidine, Garlic and Sadabahar extract (0.5%) were used (Table 1).

Extracts of the botanicals, Sadabahar (*Vinica rogeia*) and garlic (*Allium sativum*) were prepared by crushing their leaves (100g each) in 100ml of sterilized distilled water. The extracts were then filtered through a muslin cloth and centrifuged for 30 min at 5000 rpm. The extracts were sterilized by passing them through a Millipore filter (0.22 μ m pore size) using a swimmy filter adapter. The materials were dried at room temperature (25 \pm 2^o C) for 6 hours to remove the traces of water.

Subsequently 0.5% concentration of the extract of each botanical was used for bio-assay test by food poison technique. The radial growth of *S. sclerotiorum* in three replications were recorded separately and their averages were taken. The per cent inhibition over control was calculated by the formula (Bliss, 1934) as given below:

$$\text{Per cent inhibition over control} = \frac{C-T}{C} \times 100$$

where, C = growth of fungus in control

T = growth of fungus in treatment

In order to find out a suitable control of the disease, efficacy of fungicides, host extract and biopesticides were assessed in field trial during *Kharif* season 2003-2004 and 2004-2005. A highly sick field with known history of Sclerotinia blight of brinjal was selected. The brinjal variety, Black Beauty was transplanted in 4x4 m plot size in Randomized Block Design with 4 replications. The five fungicides *viz.* Vitavax and Bavistin(0.1%), Vapam (0.2%), Topsin-M(0.15%), Chlorothalonil (.02%) and one biopesticide, Nimbidine (.02%) were used as spray (Table 2). The first spray was done at the onset of disease followed by two more sprays at 10 days intervals. Observations were recorded at 15 days after the final spray. The control plots were sprayed with water. For recording the disease incidence, forty randomly selected plants per plot were examined and the disease incidence in percentage was transformed into angles and analysed statistically. Yield was estimated on plot basis without considering the border rows in Q/ha.

RESULTS AND DISCUSSION

The results presented in (Table 1) indicate that all the fungicides, plant extracts and bio-pesticide were significantly superior over control in inhibiting the growth of the pathogen *in vitro*. Vitavax, Bavistin, Blitox-50 and Topsin-M were the most effective fungicides and they completely inhibited the growth of pathogen. Nimbidine and Chlorothionil were found to be the next best in inhibiting the growth of pathogen. These were statistically different showing 97.32 and 95.00 per cent inhibition over control, respectively. Captan was the least effective fungicide. Sadabahar and garlic extracts inhibited 83.87 and 82.26 per cent growth of pathogen and appeared statistically at par to each other. The radial growth of the pathogen in the case of rest of fungicides varied from 6.95 to 48.00

mm that showed their ineffectiveness. The findings of Roy and Saikia (1976) and Singh and Gangopadhyay (1984) also confirmed the efficacy of Vitavax in checking the growth of *S. sclerotiorum* under laboratory conditions. Moore and Atkins (1977) observed fungicidal properties of the garlic extract against some fungal species.

The results of field test with five fungicides and one biopesticide, Nimbidine (Table 2) indicate their effectiveness in managing the disease. Spraying of Bavistin (0.1%) and Vitavax (0.1%) at the intervals of 15 days was more effective in minimizing the disease intensity and increasing the yield and proved statistically at par. Highest yield of 236.0 Q./ha was obtained with Bavistin followed by Vitavax (235.0 Q/ha. The next effective fungicide was Tapsin-M (0.15%) which showed

Table 1 : Effect of fungicides, plant extracts and biopesticide on colony growth of *Sclerotinia sclerotiorum* on P.D.A.

| Sr. No. | Fungicide / biopesticide | Conc. w/v | Av. colony diameter after 5 day (mm) | Per cent inhibition over control |
|---------|--------------------------|-----------|--------------------------------------|----------------------------------|
| 1. | Vitavax | 0.10 | 0.00 | 100 |
| 2. | Bavistin | 0.10 | 0.00 | 100 |
| 3. | Blitox-50 | 0.20 | 0.00 | 100 |
| 4. | Topsin-M | 0.15 | 0.00 | 100 |
| 5. | Nimbidine | 0.50 | 2.30 | 97.32 |
| 6. | Chlorotholonil | 0.20 | 4.25 | 95.00 |
| 7. | Kitazin | 0.20 | 6.95 | 91.96 |
| 8. | Ridiomil | 0.20 | 8.50 | 90.10 |
| 9. | Zineb | 0.20 | 9.12 | 89.93 |
| 10. | Captafol | 0.20 | 12.15 | 85.87 |
| 11. | Sadabahar extract | 0.50 | 14.30 | 83.87 |
| 12. | Garlic extract | 0.50 | 15.25 | 82.26 |
| 13. | Streptocycline | 0.05 | 21.00 | 75.38 |
| 14. | Indofil M-45 | 0.20 | 28.00 | 67.44 |
| 15. | Captan | 0.20 | 48.00 | 44.18 |
| 16. | Control | - | 86.00 | - |
| | C.D. (P=0.05) | | 9.98 | |

Table2: Efficacy of fungicides and biopesticide against *Sclerotinia* blight of brinjal under field condition

| Sr. No. | Product | Dose % | Disease incidence % | | | Yield Q/ ha | | |
|---------|----------------|--------|---------------------|------------------|------------------|-------------|------|-------|
| | | | 2004 | 2005 | Mean | 2004 | 2005 | Mean |
| 1. | Vitavax | 0.10 | 9.50 (17.95) | 8.50 (16.95) | 9.00 (17.45) | 232 | 238 | 235.0 |
| 2. | Bavistin | 0.10 | 9.00 (17.45) | 10.00 (18.40) | 9.50 (17.95) | 237 | 235 | 236.0 |
| 3. | Topsin-M | 0.15 | 11.25 (19.59) | 12.20 (20.44) | 11.75 (20.04) | 233 | 230 | 231.5 |
| 4. | Blitox-50 | 0.15 | 12.30 (20.53) | 13.50 (21.55) | 12.90 (21.04) | 230 | 227 | 228.5 |
| 5. | Nimbidine | 0.50 | 14.25 (22.17) | 13.80 (21.80) | 14.02 (21.99) | 225 | 226 | 225.5 |
| 6. | Chlorotholonil | 0.20 | 16.56 (23.23) | 17.25 (24.53) | 16.90 (24.27) | 220 | 218 | 219.0 |
| 7. | Control | -- | 30.20 (33.33) | 31.80 (34.32) | 31.00 (33.83) | 185 | 180 | 182.5 |
| | C.D.(P=0.05) | | (2.53) | (2.10) | (2.25) | 40.3 | 42.0 | 41.2 |

Figures in parenthesis are angular transformed value.

11.75 per cent disease and 231.5 Q/ha yield. Chlorothalonil (0.2%) proved to be the least effective one. The biopesticide, Nimbidine proved numerically better than Chlorothalonil but was at par statistically showing mean disease intensity of 14.02 and 225.5Q/ha yield.

The performance of systemic fungicides was better than non-systemic fungicides and it could be possible to control the disease through the spraying of fungicide on brinjal. Observations of Pegg (1972) on control of Sclerotinia stem rot (*S. sclerotiorum*) of Frenchbean by spraying of Benomyl support the present finding regarding foliar sprays. Similarly, Singh *et al.* (1994) suggested that Sclerotinia rot of mustard could be managed by the spraying of Vitavax, Benomyl and Topsin-M. Singh *et al.* (2003) observed the efficacy of Bavistin, Vitavax and Topsin-M under field condition against *S. sclerotiorum* in *Trachyspermum ammi* (Ajowan) and was found to be effective in managing the disease.

Authors' affiliations:

RAMESH SINGH, Department of Plant Pathology, T.D. (P.G.) College, JAUNPUR (U.P.) INDIA

P.C. SINGH, Department of Plant Pathology, T.D.(P.G.) College, JAUNPUR (U.P.) INDIA

ALKA, Department of Botany, D.G. College, KANPUR (U.P.) INDIA

REFERENCES

- Bliss, C. I. (1934).** The method of probits. *Science*, **79**: 38.
- Moore, G.S. and Atkins, R.D. (1977).** The fungicidal and fungistatic effect of an aqueous garlic extract on medically important yeast like fungi. *Mycologia*, **69**:341-348.
- Pegg, K.G. (1972).** Control of Sclerotinia stem rot in bean with Benomyl. *Australian J. Exp. Agric. & Animal Husb.*, **12**: 81-82.
- Roy, K.K. and Saikia, U.N. (1976).** White blight of mustard and its control. *Indian J. Agric. Sci.*, **46**: 214-277.
- Schmitz, H. (1930).** Suggested toximetric method for wood preservative. *Indust. Fungin Chemistry (Fed.)*, **4** : 361-633.
- Singh, Ramesh, Singh, S.B. and Palat, Ram (2003).** Management of Sclerotinia stem rot of Ajowan through fungicides and biopesticides. *Ann. Pl. Protec. Sci.*, **11 (1)** : 164-165.
- Singh, R., Tripathi, N.N., Kaushik, C.D. and Singh, R. (1994).** Management of Sclerotinia rot of Indian mustard by fungicides. *Crop Res. Hisar*, **7**: 276-281.
- Singh, U.P. and Gangopadhyay, G.S.(1984).** Control of white rot of seed cauliflower. *Pesticides*, **18**: 23-24.
