

Management of Rhizoctonia Root Rot of Cauliflower Through IDM Practices

M.R. DABBAS, D.P. SINGH AND J.R. YADAV

International Journal of Plant Protection, Vol. 2 No. 1 : 128-130 (April to September, 2009)

See end of the article for authors' affiliations

Correspondence to :
M.R. DABBAS
Department of
Vegetable Science, C.S.
Azad University of
Agriculture and
Technology, KANPUR
(U.P.) INDIA

SUMMARY

Rhizoctonia root rot due to *Rhizoctonia solani* causes much damage to the crop in curd production of cauliflower. To manage the disease through IDM practice, an experiment was conducted with six treatments of chemicals and biocontrol agents for seed treatment, seedling treatment, nursery treatment and soil treatment with four replications. The seed treatment with carbendazim @ 2g/kg seed, raising seedling in solarized beds, crop raising in green manure field + neem cake 25kg/ha with soil treatment by *Trichoderma viride* @ 2 kg/ha gave the lowest disease intensity of 11.27% with maximum curd yield 299.58q/ha and highest per cent disease control (51.00) over control treatment.

Cauliflower (*Brassica oleracea* L. var. *botrytis*) is an important and popular vegetable in India. India is the largest producer of cauliflower in the world and apart from India, the other major producers of cauliflower in the world are China, France, Italy, U.K, U.S.A., Spain, Poland, Germany and Pakistan. It is a major vegetable crop grown mainly in states like Bihar, Uttar Pradesh, Orissa, West Bengal, Assam, Haryana and Maharashtra.

Severe losses to the cauliflower are caused by different fungal pathogens such as *Pythium*, *Peronospora parasitica*, *Alternaria brassicae*, *Sclerotinia sclerotiorum*. But *Rhizoctonia solani*, is responsible for reduction and uncertain yield of cauliflower. The pathogen is soil borne in nature. Most of the cauliflower varieties are highly susceptible to *Rhizoctonia* root rot and to minimize the infection of the fungus, the present investigation was laid out.

MATERIALS AND METHODS

The experiment was laid out at the research farm of Department of Vegetable Science, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur. The experiment was conducted in Randomized Block Design (RBD) with six treatments along with four replications. The cauliflower variety Snowball-16 was taken for transplanting to know the control measures for disease. Crop was planted after thorough mixing of organic manure 150q/ha, phosphatic fertilizer (50kg P₂O₅/ha), potassic fertilizers (50 kg MOP/ha) and half nitrogenous fertilizer (62.5 kg/ha) in the soil. Remaining dose of nitrogen was applied

as broad cast at 30 days and 60 days after transplanting. The soil of experimental plot was sandy loam in nature, well drained with low CN ratio. The experiment was conducted in sick plot in both the years (2006-07 and 2007-08). Seed, seedling and soil treatment by chemicals and biopesticides were done as described by Lifschitz *et al.* (1985). Solarization of beds was done as described by Pullman *et al.* (1981). Two foliar sprays of chemicals and bioagent was done at 45 and 60 days after transplanting. Observations on disease intensity was recorded in both the crop seasons. The treatments viz., seed treatment by carboxin @ 2g/kg of seed+2 need based sprays of Flusilazole (0.1%); seed treatment by carbendazim (0.25%) + seedling dip in carbendazim solution (0.1%)+2 need based sprays of validamycin (0.1%), seed treatment by carbendazim (0.1%) + removal of infected seedling +2 sprays of cow dung (5%) + cow urine (5%), seed treatment by carbendazim (0.25%)+ raising of seedling in solarized beds+crop raising in green manure + neem cake 25 kg/ha applied in plot + soil treatment by *Trichoderma viride* 2 kg/ha and seed treatment by *Trichoderma viride* (0.5%)+ soil treatment by *Trichoderma viride*, 10g/kg FYM/ m² area+ 2 sprays of cow urine (5%) at 15 days interval from 7 days after transplanting were used.

RESULTS AND DISCUSSION

Disease intensity :

Perusal of results depicted in Table 1, revealed that significantly (P<0.05) lower disease intensity was recorded in seed

Key words :

Flusilazole,
Validamycin,
Trichoderma viride,
Cauliflower,
Rhizoctonia root rot,
Rhizoctonia solani

Accepted :
March, 2009

Table 1 : Effect of treatments on disease intensity and yield of cauliflower

Sr. No.	Treatments	Disease intensity (%)			Yield q/ha			% Disease control
		2006-07	2007-08	Mean	2006-07	2007-08	Mean	
T ₁	Seed treatment by carboxin @ 2g/kg seed +2 need based sprays of flusilazole (0.1%)	21.44	19.50	20.47	256.00	259.85	257.93	11.00
T ₂	Seed treatment by carbendazim (0.25%) +seedling dip in carbendazim solution (0.1%) + 2 need based sprays of validamycin (0.1%)	18.63	16.95	17.79	268.50	279.22	273.86	22.65
T ₃	Seed treatment by carbendazim (0.1%) + removal as infected seedling +2 sprays of dunk (5%) + cow urine (5%)	15.93	15.84	15.89	260.00	281.15	270.58	30.91
T ₄	Seed treatment by carbendazim (0.25%)+raising seedling in solarized beds +crop raising in G.M.+neem cake 25kg/ha+soil treatment by <i>T.viride</i> 2 kg/ha	11.88	10.65	11.27	294.00	305.15	299.58	51.00
T ₅	Seed treatment by <i>T. viride</i> (0.5%)+soil treatment by <i>T. viride</i> 10g/kg F.Y.M./m ² area+2 sprays of cow urine (5%) at 15 days interval.	13.46	12.50	12.98	280.50	299.50	290.00	43.57
T ₆	Control	23.79	22.20	23.00	220.00	248.20	234.10	
C.D. (P=0.05)		1.51	1.87	21.45	20.21			
CV%		4.09	4.17	5.41	6.24			

treatment by carbendazim (0.25%)+ raising of seedling in solarized beds+ crop raising in green manure +neemcake 25kg/ha +soil treatment by *Trichoderma viride* 2kg/ha (11.27%) as compared to seed treatment by *Trichoderma viride* (0.5%) +soil treatment by *Trichoderma viride* 10g/kg FYM/m² area+2 sprays of cow urine (5%) at 15 days interval after transplanting (12.98%); seed treatment by carbendazim (0.1%) + removal of infected seedlings+ 2 sprays of cow dung (5%)+ 5% cow urine (15.89%); seed treatment by carbendazim (0.25%)+ seedling dip in carbendazim solution (0.1%) + 2 need sprays of 0.1% validamycin (17.79%) and seed treatment by carboxin @ 2g/kg seed+2 need based sprays of 0.1% flusilazole (20.47%). The disease intensity recorded in present investigation is in conformity of findings reported by Iqbal and Khalid (2002), Saikia and Gandhi (2003), Harender Raj (2004), Sharma and Dureja (2004) and Abbasi *et al.* (2003).

Curd yield :

The average curd yield of cauliflower was 299.50, 290.00, 273.86, 270.58 and 257.93 q/ha in T₄, T₅, T₂, T₃, T₁, respectively. The statistical analysis revealed that significantly (P<0.05) higher curd yield of cauliflower was recorded in seed treatment by carbendazim (0.25%) +raising of seedling in solarized beds+ crop raising in green manure+neem cake 25kg/ha+ soil treatment by *Trichoderma viride* 2kg/ha (T₄) because of low disease intensity of *Rhizoctonia* root rot. The finding of present investigation is similar to Gaikwad and Nimbalkar (2003)

and Harender Raj (2004).

CONCLUSION

On the basis of above results it may be concluded that seed treatment by carbendazim (0.25%) +raising of seedlings in solarized beds+ crop raising in green manure+ neem cake 25kg/ha +soil treatment by *Trichoderma viride* 2kg/ha may be recommended for management of *Rhizoctonia* root rot disease of cauliflower.

Authors' affiliations:

J.P. SINGH, Oilseed Section, C.S. Azad University of Agriculture and Technology, KANPUR (U.P.) INDIA

J.R. YADAV, Department of Vegetable Science, C.S. Azad University of Agriculture and Technology, KANPUR (U.P.) INDIA

REFERENCES

- Aabbasi, I.H., Wakil, A.A. and Satour, M.M. (2003). Studies on the bio agent, *Trichoderma* in Egypt. *Egyptian J. Phytopathol.*, **31** (1/2): 59-73.
- Gaikwad, A.P. and Nimbalkar, C.A. (2003). Management of collar and root rot (*Rhizoctonia solani*) of bell paper with bioagent *Trichoderma* spp. and fungicides. *J. Maharashtra agric. Univ.*, **28** (3), 270-273.
- Harender Raj (2004). Effect of solarization of farmyard manure-amended soil for management of damping off caused by *Pythium ultimum*, *Rhizoctonia solani*, *Sclerotium rolfsii* in vegetable crops in nurseries. *Indian J. agric. Sci.*, **74** (8) : 425-429.

Iqbal, M.A. and Khalid, Iftikhar (2002). *In vitro* evaluation of various antagonists against mycelial growth of *Rhizoctonia solani* and effect of seed treatment with antagonists on the control of root rot of cauliflower (*Brassica oleracea* L.). *Pakistan J. Phytopathol.*, **14** (1):11-14.

Lifschitz, R., Lifschitz, S. and Baker, R. (1985). Disease incidence of *Rhizoctonia* pre-emergence damping off by use of integrated chemical and biological control. *Plant Dis.*, **69**:431-434.

Pullman, G.S., Devay, J.E., Garber, R. and Weinhold, A.R. (1981). Soil sterilization: Effect of *Verticillium* wilt of cotton and soil borne population of *Verticillium dahliae*, *Pythium* spp., *Rhizoctonia solani* and *Thielaviopsis brassicola*. *Phytopathol.*, **71**: 954-959.

Saikia, M.K. and Gandhi, S.K. (2003). Comparative activities of three antagonistic fungi against cauliflower stem rot pathogen, *Rhizoctonia solani*. *J. Mycol. Pl. Pathol.*, **33** (1) : 138-140.

Sharma, Pratibha and Dureja, Prem (2004). Evaluation of *Trichoderma harzianum* and *T. viride* isolates at BCA pathogen crop interface. *J. MycolPl. Pathol.* , **34** (1) : 47-55.
