

Screening of fungi isolated from poultry farm soil for keratinolytic activity

■ K.D. MINI, MINI K. PAUL¹ AND JYOTHIS MATHEW¹

Member of the Research Forum

Associate Author :

¹Sree Sankara College, Kalady,
ERNAKULAM (KERALA) INDIA

Author for correspondence :

K.D. MINI
Sree Sankara College, Kalady,
ERNAKULAM (KERALA) INDIA

Email: minianup@yahoo.co.in

Abstract : Poultry farm soil samples collected from different localities of Ernakulam and Thrissur districts of Kerala were screened for the keratinolytic fungi. During the course of study 8 different fungi were isolated and identified. *Aspergillus*, *Chrysosporium*, *Microsporum*, *Trichophyton* and *Penicillium* were the fungi isolated and were grown in wheat bran substrate. Feather keratin powder was added to the substrate to enhance the enzyme production. They were found utilizing keratin substrate releasing keratinase enzyme into the medium. These enzymes were assayed for their activity. Some cultural conditions were tested to attain maximum keratinase production. Maximum enzyme production was reached on the 4th day of incubation of the culture at 37°C and pH 8.5.

Key words : Keratinolytic fungus, Keratin, Keratinase, Bioremediation, Soil fungi, Enzyme assay, Keratinophilic fungus

How to cite this paper : Mini, K.D., Paul, Mini K. and Mathew, Jyothis (2012). Screening of fungi isolated from poultry farm soil for keratinolytic activity, *Vet. Sci. Res. J.*, 3(1 & 2) : 1 - 3.

Paper history : Received : 06.06.2012; Revised : 25.07.2012; Accepted : 20.08.2012

INTRODUCTION

The selected fungi can degrade the hair, nail, hoof, wool etc. present in the soil. This capacity of the fungus can be exploited for bioremediation.

Fungi are an important component of the soil microbiota more in abundance than bacteria, their population depends upon soil depth and nutrient conditions. The soil samples were collected from different sites of Ernakulam and Trichur districts. The soil collection mainly focussed the proximities of poultry farms, dumping sites of animal hair, hoof, nail etc. Different soils have specific fungus flora, but the majority of species found in them are cosmopolitan (Ainsworth and Sussman, 1968). Fungi present in the soil include keratinophilic (keratin loving) and some keratinolytic (keratin digesting) strains. Many of them are potential pathogen to both human and animals. Soils that are rich in keratinous materials are most conducive for the growth and occurrence of keratinophilic fungi (Moallaei and Zaini, 2006). Keratin is a major component of feathers. Among the microbes that cycle keratin protein in nature, keratinophilic fungi are very common

and the most diverse. If keratinolytic fungi were not there to cycle this highly stable protein (keratin), we can imagine the quantity of keratin that would have accumulated on earth, since a vast quantity of keratin is shed by the vertebrates. Indian soils contain many more keratinophilic fungi than those presently recorded, and there is need for further taxonomic and ecological studies of this interesting group of organisms (Sharma and Rajak, 2003). The potential use of keratinases have different applications where keratins should be hydrolysed, such as the leather and detergent industries, textiles, waste bioconversion, medicine, and cosmetics for drug delivery through nails and degradation of keratinized skin. Fungi also display lipolytic activity and remove petroleum hydrocarbons from the medium during degradation of proteins.

A distinctive feature of keratin is its relatively high sulphur content due to the presence of sulphur containing amino acids viz., cysteine, cystiene and methionine. Thus, the disulfide bonds are considered to be responsible for the stability of keratin and its resistance to enzymatic degradation (Kunert, 1973). Keratinolytic mycoflora love to grow and even reproduce on keratin materials such as skin, hair, nail, fur,

feather, horn, hoof, beak etc. They utilize keratin as carbon source (Cooke, 1980). Keratinophilic fungi are important ecologically and present in the environment with variable distribution patterns and cause human and animal mycoses. This study reports the prevalence of keratinolytic fungi from poultry farm soil collected from different locations of Ernakulam and Trichur districts in Kerala.

RESEARCH METHODOLOGY

Sterilised feathers, keratin, keratin salt medium, soil, distilled water, petridishes and Sabouraud dextrose agar (SDA), lactophenol cotton blue etc. were needed for the study.

Collection of soil:

Soil was collected in sealed containers using sterile spoon from the poultry farm premises where keratin and hence keratinolytic fungi were present.

Isolation of keratinolytic fungus:

10g of the soil was mixed with 100ml. of distilled water and the dilutions were plated on keratin agar medium (g/250ml) containing keratin-2.5, MgSO₄- 0.25, KH₂PO₄- 0.115, K₂HPO₄ - 0.25 and agar - 5. Streptomycin 1% was mixed with the medium. Plates were incubated at 37°C for 5 days. The plates in which clear zones were seen indicated the fungi with keratinolytic activity. They were carefully isolated and stored in SDA (Sabouraud dextrose agar) medium (g/250ml) peptone - 2.5, dextrose - 10 and agar - 5.

Identification of fungus:

The isolated fungi were stained with lactophenol cotton blue and observed it under the microscope. By noting the morphology of the fungus identification was done. For the selection of the fungi with high keratinolytic activity, already identified fungi were cultured in feather keratin substrate.

Preparation of feather keratin substrate:

Fairly large amount of chicken feather was collected from the poultry farm and washed well with chloroform-methanol (1:1, v/v), and finally with distilled water. It was then dried in sunlight and sized into 1cm length. Sterilization was done by tyndallization at 100°C for 20 min on five successive days. It was then powdered in sterile condition.

Preparation of medium for fungal culture:

The medium was prepared for inoculating the fungus as follows(g/l): wheat bran - 10, KH₂PO₄ - 0.46, K₂HPO₄ - 1, MgSO₄-0.5, at a pH 8.5. Erlenmeyer flasks containing 20 ml of sterilized medium supplemented with 10g of sterilized feather as a keratin source was incubated at 37°C and 120 rpm for 4-5 days. Flasks containing the medium with a disc of agar without the fungus served as control. For each species, one test flask

and one control set were maintained. For inoculation, spore suspensions 2x10⁸ of 5 day-old inoculum was used.

Extraction of enzyme:

Phosphate buffer having pH 8.5 was used for the extraction of culture filtrate. After 4-5 days of incubation, 100ml. of PO₄ buffer was added to each flask. The flasks were rotated in the shaking incubator at 200rpm for 1 hr. at 37°C. After 1 hr. the ingredients were filtered through whatman's No. 1 filter paper and the filtrate was used for enzyme activity.

Enzyme activity:

Keratinase was assayed by Lowry *et al.* method. One unit of enzyme activity is defined as the amount of enzyme liberating one microgram of tyrosine per minute per ml. under the defined conditions. The activity of the enzyme is expressed in units/gds (gram dry weight).

RESULTS AND DISCUSSION

Out of 60 soil samples collected from the poultry farm premises, 51 samples were found positive in fungal growth. A total of 5 genera and 8 species were isolated. In the study some of the soil samples gave single species and some yielded mixed growth of two or more species of fungi. *Aspergillus* species was isolated frequently. Table 1 reveals the per cent occurrence of fungi isolated from the soil.

Fungi isolated and % of occurrence		
Fungus	% of occurrence	Enzyme activity U/gds
<i>Spergillus niger</i>	25	406
<i>A.flavus</i>	22	560
<i>A.fumigatus</i>	12	425
<i>A.nidulans</i>	8	324
<i>Chrysosporium keratinophylum</i>	12	523
<i>Microsporium gypseum</i>	10	308
<i>Trychophyton mentagrophytes</i>	8	297
<i>Penicillium spp.</i>	3	

Among the seven species of fungi better activity was recorded in *A.flavus* (560U/gds) followed by *Chrysosporium keratinophylum* (523U/gds) *A. fumigatus* (425U/gds), *A.niger* (406U/gds) *A.nidulans* (324U/gds) *Microsporium gypseum* (308U/gds) and *Trychophyton mentagrophytes* (297U/gds). The extracellular enzymes secreted by these fungi are responsible for the degradation of keratin in nature. Change in alkalinity may be due to the high alkaline nature of the medium due to cysteine, keratinase and protein. Evidences of keratinolysis lie on the ability of fungi to release soluble sulphur containing aminoacid and polypeptides into the medium. Santos *et al.* (1996) have investigated that *A.fumigatus* was useful for

the microbial conversion of keratinous waste and *A. flavus* has been selected as a prospective producer of a keratinolytic enzyme (Gradišar *et al.*, 2000).

Conclusion:

Large number of microbes is present in the environment which is capable of recycling keratin protein. Among them keratinolytic fungi are most diverse and so common. The capacity of these fungi can be exploited for degrading the keratinous wastes and thus an effective means of bioremediation.

LITERATURE CITED

- Abu, P., Hilda, A. and Gopinath, S.C. (2004). Keratinophilic fungi of poultry farm and feather dumping soil in Tamil Nadu, India. *Mycopathologia*, **158**(3): 303-909.
- Ainsworth, G.C. and Sussman, A.S. (1968). *The Fungi an advanced treatise* Vol. III the Fungal Population, pp. 426-496
- Bodey, G.P. and Vartivarian, S. (1989). Aspergillosis. *Eur. J. Clin. Microbiol. Infect. Dis.*, **8** : 413-437.
- Cooke, R.C. (1980). *Fungi Man and Environment*. Longman Group Ltd.London pp 17-19.
- Fillipello, M.V. (2000). Keratinophilic fungi: Their role in nature and degradation of keratinic substrates. In: *Biology of dermatophytes and Other keratinophilic fungi*, Kushawaha RKS, Guarro J, editors. *Revista Iberoamericana de Micologia*. Spain, pp. 77-85.
- Gradišar, H., Kern, S. and Friedrich, J. (2000). Keratinase of *Doratomyces microsprus*. *Appl. Microbiol. Biotechnol.*, **53**: 196-200.
- Jesenka, Z., Durkovsky, J., Rosinsky, I., Polak, M., Zamboova, E. and Baca, B. (1992). Filamentous micromycetes in otitis. *Cesk Epidemiol. Mikrobiol. Imunol.*, **41**: 337-341.
- Kunert, J. (1973). Keratin decomposition by dermatophytes. I.Sulfite production as a possible way of substrate denaturation.*Z. Allg. Mikrobiol.*, **13** : 489-498.
- Moallaei, H. and Zaini, F. (2006). Isolation of keratinophilic fungi from soil samples of forests and farm yards. *Iranian J. Publ. Health*, **35**(4): 62-69.
- Santos, R. M. D. B., Firmino, A. A. P., de Sá, C. M. and Felix, C. R. (1996). Keratinolytic activity of *Aspergillus fumigatus* Fresenius. *Curr. Microbiol.*, **33** : 364-370.
- Sarquis, M.I.M and Oliveira, P.C. (1990). Diversity of microfungi in the sandy soil of Panema Beach, Rio de Janeiro, Brasil.*J. Basic Microbiol.*, **36** : 51-58.
- Sharma, R. and Rajak, R.C. (2003). Keratinophilic fungi: Nature's keratin degrading machines! Their isolation, identification, and ecological role. *Resonance*, pp. 28-40.
