

## Evaluation of various fungicides and bio-organics against foot rot disease of finger millet caused by *Sclerotium rolfsii* SAAC

■ V.G. MUNDE<sup>1</sup>, M.P. DIWAKAR<sup>1</sup>, B.B. THOMBRE<sup>1</sup> AND UTPAL DEY

### AUTHORS' INFO

#### Associated Co-author :

<sup>1</sup>Department of Plant Pathology,  
 College of Agriculture, Dr.  
 Babasaheb Sawant Konkan Krishi  
 Vidyapeeth, Dapoli, RATNAGIRI  
 (M.S.) INDIA

#### Author for correspondence :

#### UTPAL DEY

Department of Plant Pathology,  
 College of Agriculture, Dr.  
 Babasaheb Sawant Konkan Krishi  
 Vidyapeeth, Dapoli, RATNAGIRI  
 (M.S.) INDIA  
 Email : utpaldey86@gmail.com

**ABSTRACT :** The studies were undertaken for the management of foot rot disease of finger millet (nagli or ragi). Results of poisoned food technique revealed that the fungicides viz., Tilt (0.05%), Hexathir (0.2%), Score (0.05%) and Contaf plus (0.05%) were best as they totally inhibited mycelial growth. These treatments were followed by Bavistin (44.44%), Dithane M-45 (40.77%), Blue copper (38.22%) and Captaf (34.11%). In bio-organics, maximum per cent reduction was achieved due to Chetna (54.88%). As compared to this, Purna (24.11%), Reviver (21.55%) and Amogh (21.11%) gave less per cent reduction of test fungus over control.

**Key Words :** Foot Rot, *Sclerotium rolfsii*, Nagli, Fungicide, Bio-organics, Finger millet

**How to cite this paper :** Munde, V.G., Diwakar, M.P., Thombre, B.B. and Dey, Utpal (2012). Evaluation of various fungicides and bio-organics against foot rot disease of finger millet caused by *Sclerotium rolfsii* SAAC, *Adv. Res. J. Crop Improv.*, 3 (2) : 167-168.

**Paper History :** Received : 13.08.2012; Accepted : 14.11.2012

**F**inger millet [*Eleusine coracana* (L.) Gaertn] is an important millet crop grown in India. Foot rot of nagli or ragi caused by *Sclerotium rolfsii* Sacc. has been found quite serious and wide spread. It is most destructive disease of the crop as it infects the crop in seedlings stage resulting in complete failure of the crop. Under suitable condition, it causes 40 to 50 per cent loss (Basta and Tamang, 1983). Taking in to account the economic importance of the disease, the investigations were made on the disease and its causal agent.

The efficacy of various fungicides (hexathir, bavistin, blue copper, dithane M-45, tilt, score, captaf and contaf plus) and bio organics (chetna, purna, reviver and amogh) on growth and sporulation of the isolates of the pathogen was studied using poisoned food and inhibition zone techniques. A technique in which nutrient medium was poisoned with fungitoxicant and bio-organic toxicant on which the test fungus grew. It was followed here for *Sclerotium rolfsii*. For this, all the glassware used in the study were sterilized prior to use. PDA was used as a medium. PDA was prepared and sterilized at 15 lbs per square inch for 15 minutes in an autoclave and then it was distributed in 250 ml capacity conical flasks. Required quantity of each fungicide and bio-organic was added in each flask, so as to get the desired concentrations. The fungicides and bio-organics were thoroughly mixed in the medium and poured in sterilized Petri plates. Concentrations of each

fungicide and bio-organic were replicated thrice. The culture of the test fungus, *Sclerotium rolfsii* was grown on PDA for 4 – 5 days until the Petri plates were fully covered but did not form the sclerotia.

The fungal discs of 0.5 cm diameter were cut with the help of sterilized cork borer and transferred aseptically under the condition of Laminar Flow Bench at the centre of each Petri plate, containing poisoned PDA medium. The suitable control of PDA without fungicide and bio-organic inoculated with fungal disc served as control. The plates were incubated at 28 ± 1°C in incubator and observations for colony diameter and sclerotial formation were recorded until whole of the plate in control treatment was fully covered with mycelial growth.

Per cent inhibition of growth was calculated by the following formula (Horsfall, 1956):

$$X = \frac{Y - Z}{Y} \times 100$$

where,

X = Per cent inhibition

Y = Growth of fungus in control (cm)

Z = Growth of fungus in treatment (cm)

Data obtained on per cent inhibition was subjected to statistical analysis.

The total eight fungicides and four bio-organics were

**Table 1 : *In vitro* effects of various fungicides and bio-organics on colony diameter of *Sclerotium rolfii* Sacc**

Tr. No.	Fungicide and bio-organic	Concentration (%)	Mean colony diameter (cm)*	Per cent inhibition over control
T <sub>1</sub>	Carbendazim (Bavistin)	0.20	5.00	44.44
T <sub>2</sub>	Mancozeb (Dithane M-45)	0.25	5.33	40.77
T <sub>3</sub>	Copper oxychloride (Blue copper)	0.25	5.56	38.22
T <sub>4</sub>	Amogh	0.30	7.11	21.11
T <sub>5</sub>	Propiconazole	0.05	0.00	100.00
T <sub>6</sub>	Thiram (Hexathir)	0.20	0.00	100.00
T <sub>7</sub>	Difencozole	0.05	0.00	100.00
T <sub>8</sub>	Hexaconazole	0.05	0.00	100.00
T <sub>9</sub>	Captan (Captaf)	0.05	5.93	34.11
T <sub>10</sub>	Purna	0.30	6.83	24.11
T <sub>11</sub>	Reviver	0.30	7.06	21.55
T <sub>12</sub>	Chetana	0.30	4.06	54.88
T <sub>13</sub>	Control (No fungicide and bioorganic)	--	9.00	
	S.E. ±		0.17	
	C.D. (P=0.05)		0.50	

\* Mean of three replications

screened against *Sclerotium rolfii* by PFT. The data in Table 1, revealed that all the fungicides and bio-organics screened against *S. rolfii* by PFT were found significantly superior in reducing the growth of test fungus when measured as colony diameter. Total inhibition (100%) of *S. rolfii* was achieved due to tilt (0.05%), hexathir (0.2%), score (0.05%) and contaf plus (0.05%) being equally effective and statistically at par. These were followed by bavistin (0.2%), dithane M-45 (0.25%), blue copper (0.25%) and captaf (0.05%) resulting in 44.44, 40.77, 38.22 and 34.11 per cent reduction of test fungus over control, respectively. All the treatments T<sub>5</sub>, T<sub>6</sub>, T<sub>7</sub>, T<sub>8</sub> and T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub> were statistically at par. In case of bio-organics maximum per cent reduction was achieved due to Chetna (54.88%). As compared to this, purna, reviver and amogh gave less per cent reduction of test fungus over control as 24.11, 21.55 and 21.11 per cent, respectively. All the fungicides and bio-organics tested were found significantly effective in controlling the fungus when compared to control. Similar results were obtained by

Bharadwaj and Agarwala (1986) who reported that thiram (3000 ppm) was best to control seedling blight of apple caused *S. rolfii*. Also, Nagmote (2000) observed that the captan (0.25%), thiram (0.25%) and mancozeb (0.25%) were found effective against *S. rolfii* whereas carbendazim and copper oxychloride were found least effective. Johnson and Subramanyam (2000) also reported that complete inhibition of radial growth of *S. rolfii* was achieved by hexaconazole, penconazole and propiconazole, tricyclazole and mancozeb, resulting in 80 and 53 per cent inhibition, respectively. No inhibition was recorded with carbendazim and copper oxychloride.

Results obtained in present study are also in agreement with Prabhu and Hiremath (2003) who screened both systemic and non-systemic fungicides against the pathogen *S. rolfii* and found that hexaconazole and propiconazole were highly effective among the systemic and thiram and mancozeb showed the minimum per cent inhibition of mycelial growth of the fungus among the non-systemics.

## LITERATURE CITED

- Batsa, B.K.** and Tamang, D.B. (1983). Preliminary report on the study of millet diseases in Nepal. In : Maize and finger millet, 10<sup>th</sup> summer workshop 23-28 Jan. 1983, Rampur, Chitwan.
- Bharadwaj, L.N.** and Agarwala, R.K. (1986). Seedling blight of apple and its control. In : Advances in research on temperate fruits. Proc. of the National Symposium on temperate fruits. pp. 397-401.
- Horsfall, J.G.** (1956). *Principles of fungicidal action*. Cronical Botanica (U.S.A.) Publ., 207pp.
- Johnson, M.** and Subramanyam, K. (2000). *In vitro* efficacy of fungicides against stem rot pathogen (*Sclerotium rolfii*) of groundnut. *Ann. Pl. Protec. Sci.*, **8**(2): 255-257.
- Nagmote, A.M.** (2000). Virulence viability, host range and control of *Sclerotium rolfii*. M.Sc. Thesis, Dr. Panjarao Deshmukh Krishii Vidyapeeth, Akola, M.S. (INDIA).
- Prabhu, H.U.** and P.C. Hiremath 2003: Bioefficacy of fungicides against collar rot of cotton caused by *S. rolfii* Sacc. *Karnataka J. Agric. Sci.*, **16**(4) : 576-579.

\*\*\*\*\*