

Viability of probiotics in flavoured yoghurts made with different starter culture during storage

■ MADHU, M. SHIVA PRAKASH AND ANITA KOCHHAR

SUMMARY : Six different yoghurts were prepared with mango pulp and pineapple essence and sugar combination with *Lactobacillus bulgaricus*, *Streptococcus thermophilus*, *Lactobacillus acidophilus*, *Lactobacillus sporogens*, *Bifido bifidum*, *Bifido longum* and *Bifido infantis* as starter culture. Three types of yoghurts were developed under each flavours, with different combination of probiotics and termed as A₁, B₁, C₁, A₂, B₂ and C₂. A₁, B₁, C₁ were mango yoghurts and A₂, B₂ and C₂ were pineapple yoghurts. Statistically, the significant difference were found in viable counts from 7th day to 14th day (P<0.05) in all yoghurts except in A₂ and C₂. The difference of viable counts from 0 day to 7th day was also found to be significant (P<0.05) in A₂ and C₂. No significant difference was found from 14th day to 30th day in all yoghurts as 90- 96 per cent viability loss of probiotics was observed by 14th day in all yoghurts. Pineapple yoghurt with probiotic blends of *Lactobacillus bulgaricus*, *Streptococcus thermophilus* and *Lactobacillus sporogens i.e.*, B₂ had higher viability, among the different types of yoghurts developed.

KEY WORDS : Analysis of variance (ANOVA), Complete randomized design (CRD), A- *Lactobacillus bulgaricus*, *Streptococcus thermophilus*, B- *Lactobacillus bulgaricus*, *Streptococcus thermophilus*, *Lactobacillus sporogens*, C- *Lactobacillus bulgaricus*, *Streptococcus thermophilus*, *Lactobacillus*, *acidophilus*, *Bifido bifidum*, *Bifidum longum*, *Bifido infantis*

How to cite this paper : Madhu, Prakash, M. Shiva and Kochhar, Anita (2012). Viability of probiotics in flavoured yoghurts made with different starter culture during storage. *Internat. J. Proc. & Post Harvest Technol.*, 3 (2) : 286-290.

Research chronicle : Received : 21.06.2012; **Revised :** 12.10.2012; **Accepted :** 22.11.2012

Yoghurt is considered by nutritionists to be a very nutritious and healthy food compared to milk. Probiotic cultures are live bacteria which help in better absorption of nutrients. They play an important role in reduction of serum cholesterol, alleviation of lactose intolerance, reduction of diarrhea, prevention and suppression of colon cancer, stimulation of the immune system etc. Yoghurt is prepared by fermenting milk with starter cultures containing different types of probiotics, normally *streptococcus thermophillus* and *lactobacillus bulgaricus*. Use of different probiotic blends in combination have several health benefits. Lactic Acid Bacteria (LAB) and bifidobacteria are the

most common types of microbes used as probiotics. Probiotics are commonly consumed as part of fermented foods with specially added active live cultures, such as in yoghurt, soy yoghurt, or as dietary supplements. The introduction of yoghurt with added probiotics could play a significant role in national health care programs especially in developing countries, where diarrhoea and gastrointestinal problems are common. The increase in the per capita annual consumption of yoghurt in the majority of the countries has been attributed to both the ever-increasing availability of fruit or flavoured yoghurt, and to the diversity of presentations of the product. Although the main choice of any probiotic microbial strain to be used as a starter culture or a blend with a starter culture is based on the health aspects beneficial to humans (Gardiner *et al.* 2002). The aim of the study was to investigate the changes in microbiological properties in mango and pineapple yoghurt made with different probiotic cultures during storage.

MEMBERS OF THE RESEARCH FORUM

Author for Correspondence :

MADHU, Department of Foods and Nutrition, Post-Graduate and Research Centre, Acharya N.G. Ranga Agricultural University, Rajendra Nagar, HYDERABAD (A.P.) INDIA

Coopted Authors:

M. SHIVA PRAKASH, Department of Microbiology and Immunology, National Institute of Nutrition (ICMR) HYDERABAD (A.P.) INDIA


ANITA KOCHHAR, Department of Food and Nutrition, College of Home Science, Punjab Agricultural University, LUDHIANA (PUNJAB) INDIA

EXPERIMENTAL METHODS

The raw material *viz.*, double toned milk, powdered sugar,

milk powder, pineapple essence, food color, pasturised mango pulp, and plastic sterile cups used for preparation of yoghurts were purchased from the local market.

The probiotic stock cultures required for the study *i.e.*, *Lactobacillus bulgaricus*, *Streptococcus thermophilus*, *Lactobacillus acidophilus*, *Lactobacillus sporogens*, *Bifido bifidum*, *Bifido longum*, and *Bifido infantis* in powder form were obtained from National Institute of Nutrition, Hyderabad. The two different flavoured yoghurt *i.e.*, mango and pineapple was prepared by using standard technique as described below in the form of flow chart:



The prepared products were tested for the viability of probiotics within three hours of preparation as well as in stored samples after 7, 14, 21 and 30 days. The microbiological method *i.e.*, Spread plate method (Clark, 1993) used for bacterial count on MRS plate is described below.

One gram of sample was transferred into a test tube containing 9 ml of saline. It was mixed well by shaking for 2 min. and 1 ml from this was taken and diluted further through a series of test tubes, containing 9 ml of sterile normal saline blank solution by an appropriate decimal dilution method. Recommended final dilution was upto 10^8 - 10^9 . Then, from the final diluted tube take 100 μ l and pour into the MRS agar medium petridishes and spread into medium by clockwise and anticlockwise via spread plate technique. The plates were incubated at 40°C for 24- 48 hours and the number of colonies in each plates was counted in colony counter machine. These isolated colonies were sub cultured and subjected for gram staining for identification of species.

Lactobacilli and Bifidobacteria appeared as large, white colonies in or on the surface of the medium. The average number of colonies, multiplied by the dilution number (dilution factor) and divided by quantity of sample on weight basis and expressed as colony forming units/gram.

$$\text{Number of colony forming units (Cfus) per g of the sample} = \frac{\text{Mean number of Cfu's} \times \text{Dilution}}{\text{Quantity of sample on weight basis}}$$

The results obtained from 10 replications of all organoleptic qualities scores and 2 replications on viable counts of log cfu/g values were analysed by analysis of variance (ANOVA), using Complete Randomized Design (CRD) and Tukey HSD Test for Post-ANOVA Pair-Wise Comparisons. Analysis of variance were applied for testing the significance between the flavours, period of storage, different products and viability of products.

EXPERIMENTAL FINDINGS AND ANALYSIS

The viability of probiotic bacteria in yoghurt depends on the strains used, the interaction between species present, the production of hydrogen peroxide due to bacterial metabolism and the final acidity of the product.

The percentage loss of probiotic bacteria at different periods of storage at 4 °C are presented in Table 1. On the day of preparation and after storing the sample for a period 7th, 14th, 21st and 30th days, the average percentage of viability loss in mango flavoured yoghurts A₁, B₁ and C₁ was 29.9 per cent, 26.53 per cent and 24 per cent at 7th day and 99.99 per cent, 99.7 per cent and 99.9 per cent at 30th day respectively. In the case of pineapple flavour yoghurts A₂, B₂ and C₂ it was 26 per cent, 27 per cent and 29 per cent on 7th day and 98.4 per cent, 99.4 per cent and 99.95 per cent on 30th day, respectively. C₁ and C₂ had high viability loss *i.e.*, 93.56 and 92.5 per cent on 14th days compared to others. The loss of probiotic in all yoghurt could be due to

competition for survival of probiotics as, *Lb. bulgaricus* affect the survival of *Lb. acidophilus* and *bifidobacteria* due to the acid and hydrogen peroxide produced during fermentation stage. Due to its proteolytic nature, *Lb. bulgaricus* grows rapidly and produces acid quickly; whilst appearing to liberate the essential amino acids such as valine, glycine and histidine, that are required to support the growth of *bifidobacteria*. *S. thermophilus* does not inhibit the growth of probiotic organisms. These observation were in conformity with that of Mortazavian *et al.* (2007) who reported the decrease in viability of *L.acidophilus* and *B.lactis* at 2°C, 5°C and 8°C of refrigerated storage of 20 days.

Table 1 : Viability loss (%) of probiotics in different period of storage

| Flavour | yoghurts | Storage period | | | |
|-----------|----------------|----------------|--------|--------|--------|
| | | 7 | 14 | 21 | 30 |
| Mango | A ₁ | 29.9 % | 96.06% | 99.05% | 99.99% |
| | B ₁ | 26.53% | 95% | 99.02% | 99.7% |
| | C ₁ | 24% | 93.56% | 97.75% | 99.9% |
| | A ₂ | 26% | 95% | 98.6% | 98.4% |
| Pineapple | B ₂ | 27% | 90.8% | 99.16% | 99.4% |
| | C ₂ | 29% | 92.5% | 94.5% | 99.95% |

The log cfu /g values at 10⁹ of the probiotic strains in developed yoghurt at different periods of storage are presented in Table 2 and Fig. 1, 2 and 3 which shows the variation in the microbial cell count during the refrigerated storage period at 7-day intervals. During storage, the viable probiotic microorganism in A₁ was 1.733 ± 0.347 × 10⁹ on 0 day reduced to 1.466 ± 0.297 × 10⁹ on 7th day of storage. From 0 day to 7th day no significant difference found. But it reduced to 0.277 ± 0.038 × 10⁹ on 14th day of storage. The analysis of variance in difference of viable counts from 7th day to 14th day were found to be significant (P<0.05). No significant difference was found from 14th day to 30th days as viability loss was 90- 96 per cent by 14th day itself only in different yoghurts.. Luther *et al.* (1931) stated that high counts of viable organisms were obtained at lower or moderate acidities and greater destruction occurred in more acid cultures. According to Dave and Shah (1996) hydrogen

peroxide produced by *L. delbruecki* ssp. *Bulgaricus* bacteria is the most important viability-reducing factor during refrigerated storage. Similar trends were observed in B₁ and C₁.

Likewise, during storage of pineapple yoghurts, the probiotic counts declined as shown in Table 2 and Fig. 4, 5 and 6. During storage, the viable probiotic microorganism in A₂ which was 1.733 ± 0.347 × 10⁹ on 0 day, reduced to 1.210 ± 0.294 × 10⁹ on 7th day. In C₂ reduction was from 2.362 ± 0.187 × 10⁹ to 1.354 ± 0.255 × 10⁹ on 7th day of storage. It further reduced to



Fig 1: Viability of probiotics in mango A₁ yoghurt during different periods of storage



Fig 2: Viability of probiotics in mango B₁ yoghurt during different periods of storage



Fig 3: Viability of probiotics in mango C₁ yoghurt during different periods of storage

Table 2: Mean value of viability of probiotics count in the mango and pineapple yoghurt from 0 days to 4 weeks at 7 day intervals (log cfu/g)

| Flavours | yoghurts | 0 day | 7 d | 14 d | 21 d | 30 d |
|-----------|----------------|---|--|--|---|---|
| Mango | A ₁ | 1.733 ± 0.347 _a (5×10 ⁹) | 1.4661 ± 0.297 _a (3.5×10 ⁹) | 0.277±0.038 _b (3.2×10 ⁸) | 0.082 ± 0.009 _b (8.5×10 ⁷) | 0.046± 0.021 _b (4.8×10 ⁷) |
| | B ₁ | 2.291±0.151 _a (9×10 ⁹) | 2.035±0.195 _a (6.8×10 ⁹) | 0.366±0.104 _b (4.5×10 ⁸) | 0.084± 0.007 _b (8.9×10 ⁷) | 0.026± 0.005 _b (2.0×10 ⁷) |
| | C ₁ | 2.362 ± 0.187 _a (9.8×10 ⁹) | 2.067±0.275 _a (7.2×10 ⁹) | 0.509±0.138 _b (6.8×10 ⁸) | 0.195±0.082 _b (2.2×10 ⁷) | 0.009 ± 0.002 _b (9.0×10 ⁶) |
| | A ₂ | 1.733 ± 0.347 _a (5×10 ⁹) | 1.210±0.294 _{ab} (2.5×10 ⁹) | 0.266±0.084 _b (3.0×10 ⁸) | 0.067±0.019 _b (7.0×10 ⁷) | 0.063± 0.005 _b (6.5×10 ⁷) |
| Pineapple | B ₂ | 2.291±0.151 _a (9×10 ⁹) | 1.781±0.394 _a (5.4×10 ⁹) | 0.599±0.011 _b (8.2×10 ⁸) | 0.072± 0.009 _b (7.5×10 ⁷) | 0.047±0.024 _b (4.2×10 ⁷) |
| | C ₂ | 2.362 ± 0.187 _a (9.8×10 ⁹) | 1.354±0.255 _b (3×10 ⁹) | 0.539±0.087 _{bc} (7.2×10 ⁸) | 0.042±0.012 _c (4.3×10 ⁷) | 0.005±0.000 _c (5.0×10 ⁶) |

Means with different subscripts in rows were significantly different P<0.05

The values in tables are calculated by logarithmic method ,all the values are expressed in 10⁹ Values in parentheses are original value




Fig 4: Viability of probiotics in pineapple A₂ yoghurt during different periods of storage



Fig 5: Viability of probiotics in pineapple B₂ yoghurt during different periods of storage



Fig 6: Viability of probiotics in pineapple C₂ yoghurt during different periods of storage

$0.266 \pm 0.084 \times 10^9$ in A₂, $0.539 \pm 0.087 \times 10^9$ in C₂ on 14th day of storage. The analysis of variance in difference of viable counts from 0 day to 7th day and 7th day to 14th day were found to significant ($P < 0.05$). Whereas, in B₂ containing *Lactobacillus bulgaricus*, *Streptococcus thermophilus* and *Lactobacillus sporogenes* there was no significant difference from 0 day to 7th day, but significant decrease was found from 7th day to 14th day. No significant difference was found from 14th day to 30th day as viability loss was 90-96 per cent by 14th day itself only in different yoghurts. These observations indicated continuous declining of probiotics in all the developed yoghurt samples during storage. Tamime *et al.* (2005) suggested that probiotic organisms should be present in a food to a minimum concentration of 10^6 cfu g⁻¹, or the daily intake should be about 10^9 cfu g⁻¹. Such high numbers have been suggested to compensate for possible losses in the numbers of the probiotic organisms during passage through the stomach and intestine. In the present study, the level of 10^9 cfu g⁻¹ was observed at the time of production *i.e.*, on 0 day in all types of yoghurts.

The viability of probiotic organism in C₁ (7.2×10^9) was

more in comparison to other types on 7th day of storage. However, the flavour of pineapple with B₂ blends of probiotics *i.e.*, 8.2×10^8 significantly increased the viability when compared to A₁, B₁, C₁, A₂ and C₂ at 14th day. It could be due to *L. sporogenes* which have longer viability than others. In support of these finding a stability study, done by Losada and Olleros.(2002) reported that *L. sporogenes* were highly viable during storage.

Payot *et al.* (1999) reported that spores of *B. coagulans* / *L. sporogenes* are ellipsoidal bodies located at one of the cellular poles, resistant to heat and adversal environmental conditions, and able to germinate in presence of diluted HCl or NaOH solutions. The longevity of yoghurt was significantly more by fermenting the yoghurt by B₂ blends of probiotics than with other types of probiotic blends. The reduction of probiotics counts observed on 7th day of storage in this study might be due to the incorporation of pineapple essence or mango pulp.

The lowest viability was observed in A₂ containing *L. bulgaricus* and *S. thermophilus* on 14th day of storage. It could be due to *S. thermophilus* and *L. bulgaricus* might not be grown properly at 4° C. Vinderola *et al.* (2000) reported that after 7 days of storage, the counts of *L. bulgaricus* showed a sharp decline (6–10-fold) from the initial counts and finally decreased by 97–99 per cent in all samples at the end of storage. They also reported that *S. thermophilus* counts were higher by at least 1 log order than those for *L. delbrueckii ssp. bulgaricus* in yoghurts containing probiotic bacteria. Kim *et al.* (1993), Medina and Jordano (1994), Lim *et al.* (1995) and Dave and Shah (1997b) also reported almost similar results. However, in the present study viability could not be carried out on individual species of probiotics, but viability was tested on a combination of organism. Mortazavian *et al.* (2007) reported that at higher temperatures 5 and 8° C compared to 2° C the *L. Delbrueckii ssp. Bulgaricus* bacteria grow faster, therefore, higher amounts of lactic acid and hydrogen peroxide could be produced. Chandan and Shahani (1993), suggested that yoghurt must have at least 10^8 cfu/g probiotic counts at the time of preparation. In the present study the viability counts were 10^8 upto 14 day which satisfies the above suggestion .

Bakirci *et al.* (2008) observed that addition of fruit and sugar in yoghurt led to an increase in the viable counts of probiotic organisms. They also reported that the total viable counts of *L. acidophilus* and *Bifidobacterium spp.* decreased slightly for longer period of storage, but remained at sufficient levels ($> 6 \log$ cfu/g) up to 14th days. Con *et al.* (1996) reported that the addition of fruit flavours to yogurt did not show significant effect on the viability of total bacteria.

Overall, it was observed that in all the yoghurt developed, the probiotics used were viable incorporated upto 7th day and after that declined from 14th day. These levels of live organisms present upto 7th day were sufficient to have beneficial effect *i.e.*, to have probiotic effect. From the finding of the present

study, the yoghurt B₂ containing *Lactobacillus bulgaricus*, *Streptococcus thermophilus* and *Lactobacillus sporogenes* can be recommended as a is a good combination of probiotic for higher viability, specially with addition of pineapple essence added yoghurt

Conclusion :

The demand for fruit flavoured yoghurts is increasing in the recent years. Hence, there is a great scope to develop and popularize fruit yoghurts in India. There are very few research studies on the flavoured yoghurt with different combination of probiotic blends.

During storage, the viable probiotic microorganisms in A₁ were 1.733×10^9 on 0 day which reduced to 1.466×10^9 on 7th day of storage. From 0 day to 7th day no significant difference was found. But it reduced to 0.277×10^9 on 14th day of storage. The difference in viable counts from 7th day to 14th day were found to be significant ($P < 0.05$). After that significant difference was not found from 14th day to 30th day. Similar trends were observed in B₁ and C₁. The viability of probiotic organisms in C₁ $2.067 \times 10^9 (7.2 \times 10^9)$ was more in comparison to other types

on 7th day of storage.

The viability of probiotics in all yoghurts was assessed using spread plate method at 0, 7th day, 14th day, 21st day and 30th day. The viability of probiotic organisms declined from 0 day to 30th day and in C₁ the viability of probiotics was higher *i.e.*, $2.067 \times 10^9 (7.2 \times 10^9)$ in comparison to other types on 7th day of storage and the viability of B₂ blends of probiotics $0.599 \times 10^9 (8.2 \times 10^8)$ was higher on 14th day, compared to other yoghurts. Statistically, the significant difference were found in viable counts from 7th day to 14th day ($P < 0.05$) in all yoghurts except in A₂ and C₂. The difference of viable counts from 0 day to 7th day was also found to be significant ($P < 0.05$) in A₂ and C₂. No significant difference was found from 14th day to 30th day in all yoghurts as 90- 96 per cent viability loss of probiotics was observed by 14th day in all yoghurt. Pineapple yoghurt with probiotic blends of *Lactobacillus bulgaricus*, *Streptococcus thermophilus* and *Lactobacillus sporogens i.e.*, B₂ had higher viability, among the different types of yoghurt developed. The probiotic based yoghurt developed can be used as food supplement for human consumption, as these products have commercial value. These can be supplied as nutraceuticals or therapeutical supplements.

LITERATURE CITED

- Bakirci, I. and Kavaz, A. (2008).** An investigation of some properties of banana yogurts made with commercial ABT-2 starter culture during storage. *Internat. J. Dairy Technol.*, **61**(3):270-276.
- Con, A.H.C., Akmakci, S.M.C., Aglar, A. and Gokalp, H.Y. (1996).** Effects of different fruits and storage periods on microbiological qualities of fruit-flavoured yoghurt produced in Turkey. *J. Food Protec.*, **59**(4): 402-406.
- Chandan, R.C. and Shahani, K.M. (1993).** *Yoghurt: In dairy science and technology 2*. Product Manufacturing, ed. YH Hui, PVCH Publishers. I -/56. NEW YORK, U.S.A.
- Dave, R.J. and Shah, N.P. (1997b).** Viability of yoghurt and probiotics bacteria in yoghurt made from commercial starter cultures. *Internat. Dairy J.*, **7**:31-34.
- Gardiner, G.E., Bouchier, P., O'Sullivan, E., Kelly, J., Collins, J.K., Fitzgerald, G., Ross, R.P. and Stanton, C. (2002).** A spray-dried culture of cheese manufacture. *Internat. Dairy J.*, **12**:749-756.
- Kim, E.R., Lee, K.W., Park, Y.H and Kwak, H.S. (1993).** The study of lactic acid bacteria in yoghurt during delivery and storage. *Korean J. Dairy Sci.*, **14**: 260-263.
- Lim, K.S., Huh, C.S., Baek, Y.J. and Kim, H.U. (1995).** A selective enumeration medium for bifidobacteria in fermented dairy products. *J. Dairy Sci.*, **78**:2108-2112.
- Losada, M.A. and Olleros, T. (2002).** Towards a healthier diet for the colon: the influence of FOS's and Lactobacilli on intestinal health. *Nutr. Res.*, **22**: 71-84.
- Medina, L.M. and Jordano, R. (1994).** Survival of constitutive microflora in commercially fermented milk containing bifidobacteria during refrigerated storage. *J. Food Protec.*, **56**: 731-733.
- Mortazavian, A.M., Ehsani, M.R., Mousavi, S.M., Rezaei, K., Sohrabvandi, S. and Reinheimer, J.A. (2007).** Effect of refrigerated storage temperature on the viability of probiotic micro-organisms in yogurt. *Internat. J. Dairy Technol.*, **60**(2): 123-127.
- Payot, T., Chemaly, Z., Fick, M. (1999).** Lactic acid production by *Bacillus coagulans*-kinetic studies and optimization of culture medium for batch and continuous fermentations. *Enzyme & Microbial Technol.*, **24**: 191-199.
- Tamime, A.Y. (2005).** *Probiotic Dairy Products*, Black well Publishing, Oxford. 39-72.
- Vinderola, C.G., Bailo, N. and Reinheime, J.A. (2000).** Survival of probiotic microflora in Argentinian yoghurts during refrigerated storage. *Food Res. Internat.*, **33**(2) : 97-102.

