

Attraction of male moth catches of boll worms toward sex pheromone traps in Bt. cotton

J.K. PATEL, M.V. VEKARIA, I.S. PATEL AND P.S. PATEL

International Journal of Plant Protection (October, 2010), Vol. 3 No. 2 : 332-334

See end of the article for authors' affiliations

Correspondence to :

I.S. PATEL

Department of
Entomology, C.P.
College of Agriculture,
S.D. Agricultural
University,
SARDARKRUSHI-
NAGAR (GUJARAT)
INDIA

SUMMARY

The results revealed that the attraction of male moths of *Helioverpa* to pheromone traps commenced during the first week of September. Maximum attraction of male moth was trapped during the second week of October (13/trap). The attraction of spotted boll worm started in the beginning of the last week of September to first week of February. It was maximum during between second week of October to first week of November. The attraction of male moths of pink boll worm started in the beginning of the fourth week of November and continued till the first week of March. The maximum trapped moths were in the last week of January (19/traps/week) and maximum moths were 112/traps. The attraction of *Spodoptera* started during the second week of August. The maximum attraction of male moths was in the second week of September (38 moths/trap/week).

Key words : Bt. cotton, Cotton boll worms, Sex pheromone

In cotton, boll worms are the most destructive pests which cause severe loss in terms of quality and quantity of cotton. Some times, serious outbreak of the boll worm has been found to cause loss up to 40 to 50 per cent in yield and even total failure of the crop. For the management of cotton boll worm, sex pheromone trap is one of the components of integrated pest management. Before developing any spray schedule programme, study on attraction of male moths toward sex pheromone traps in cotton crop is very essential. Attempt was therefore made on cotton during *Kharif*, 2007-08.

MATERIALS AND METHODS

Bt. cotton crop was sown in the 1st week of July during *Kharif* 2008-09 in plots of 800 m². Four sex pheromone traps were installed in the centre of 200 sq.m. cropped area and a single rubber septa viz., *Helicoverpa armigera* for helilure, *Earias vittella* for ervitlure, *Pectinophora gossypiella* for gossyplure and *Spodoptera* for spodolure was placed separately in each pheromone trap. The rubber septae changed at 15th days intervals and traps were positioned just above the canopy height of cotton crop. Male moth catches in each trap of each boll worm were counted daily and mean catches per trap per week of each boll

worm were computed.

RESULTS AND DISCUSSION

The results obtained from the present investigation have been discussed under following sub heads :

American boll worm:

The attraction of male moths of *Helioverpa* to pheromone traps was commenced during the first week of September 2008-09. Maximum attraction of male moth was 13 in during the second week of October. It was gradually declined in the second week of December. Throughout the crop season, 82 male moths /trap were trapped in sex pheromone trap. The present findings are similar to the results of Gupta *et al.* (1996) who reported that the American boll worm started in the end of July and higher during the middle of November to middle of January when cotton crop had maximum fruiting bodies. Pawar *et al.* (1984) also observed more moth catches of *H. armigera* during August-September and November-December.

Spotted boll worm:

The attraction of male moths of spotted boll worm (*E. vittella*) started in the beginning of the last week of September to first week of

Accepted :
September, 2010

Table 1 : Monitoring of boll worm pests in *Bt* cotton through pheromone traps

Standard week	Date	Moth catches			
		<i>H. armigera</i>	<i>E. vittella</i>	<i>P. gossypiella</i>	<i>S. litura</i>
29	20-07-08	0	0	0	0
30	27-07-08	0	0	0	0
31	03/08/08	0	0	0	0
32	10/08/08	0	0	0	1
33	17-08-08	0	0	0	3
34	24-08-08	0	0	0	8
35	31-08-08	0	0	0	27
36	07-90-8	2	0	0	38
37	14-09-08	6	0	0	28
38	21-09-08	8	0	0	21
39	28-09-08	11	1	0	14
40	05-10-08	9	2	0	15
41	12-10-08	13	13	0	9
42	19-10-08	7	35	0	7
43	26-10-08	5	39	0	4
44	02-11-08	4	36	0	1
45	09-11-08	7	45	0	0
46	16-11-08	2	14	0	0
47	23-11-08	4	9	0	0
48	30-11-08	1	7	0	0
49	07-12-08	2	5	0	0
50	14-12-08	1	4	0	0
51	21-12-08	0	2	0	0
52	28-12-08	0	3	8	0
1	04-01-09	0	2	12	0
2	11-01-09	0	3	14	0
3	18-01-09	0	6	16	0
4	25-01-09	0	4	19	0
5	01-02-09	0	1	17	0
6	08-02-09	0	0	9	0
7	15-02-09	0	0	3	0
8	22-02-09	0	0	1	0
9	01-03-09	0	0	0	0
Total :		82	231	112	147

February during 2008-09. It ranged from 1 to 45 /trap/ week. The moth catches of spotted boll worm observed maximum between second week of October to first week of November. Thereafter, emergence of spotted boll worm gradually reduced but continued in small number till the first week of February. Throughout the crop season, 231 male moths/trap were recorded. Similar findings were also reported by Gupta *et al.* (1996) and Bharpoda *et al.* (2000).

Pink boll worm:

The attraction of male moths of pink boll worm (*P.*

gossypiella) started in the beginning of the fourth week of November and continued till the first week of March during 2008-09. The maximum trapped moths were in the last week of January (19 /traps/week). Thereafter, emergence of pink boll worm gradually reduced but continued in small number till the fourth week of February. During the season, pink boll worm moths were trapped 112/traps. The present findings match with the Natarajan and Dhara (2003) who reported that the first moth catches was in the second week of November and high catches in the month of January and the peak level was in the

second week of January.

***Spodoptera litura*:**

The attraction of male moths started during the second week of August. The maximum attraction of male moths was recorded in the second week of September (38 moths/trap/week). Attraction gradually declined after second week of October. Throughout the season, attraction of male moths were (147 /traps). Similar findings were also reported by Bharpoda *et al.* (2000).

Authors' affiliations:

M.V. VEKARIA, J.K. PATEL AND P.S. PATEL,
Department of Entomology, C.P. College of Agriculture,
S.D. Agricultural University, SARDARKRUSHI-
NAGAR (GUJARAT) INDIA

REFERENCES

- Bharpoda, T.M., Patel, H.P., Patel, Usha P., Patel, G.P., Patel, J.J. and Patel, J.R. (2000).** Integrated pest management (IPM) in cotton H-6 cultivated in middle Gujarat. *Indian J. Ent.*, **62** (4) : 327-331.
- Gupta, M.P., Gupta, D.P. and Shrivastav, K.K. (1996).** Population dynamics of cotton boll worms in Madhya Pradesh. *Annl. Entomol.*, **14** (1) : 61-66.
- Natarajan, K. and Dhara, J.B. (2003).** Studies on the population dynamics of cotton pests and their natural enemies in the cotton ecosystem. Central Institute of Cotton Research, Coimbatore. : pp. 60.
- Pawar, C.S., Sithanathan, S., Bhatnagar, V.S., Srivastav, C.P. and Reed, W. (1984).** The development of sex pheromone trapping of *Helicoverpa armigera* at ICRISAT, India. *Trop. : Pest Mgmt.*, **34** (1) : 39-43.
