



## Extraction of organic compounds from different three medicinal plant

GOLOKJA KRISHNA DWIVEDI AND ARVIND KUMAR SINGH

### ABSTRACT

The plants have long served a major source of medicinal compounds. Medicinal plants are value added for the contents and the chemical composition of their active components. Therefore, the demand of plant based therapeutics has increased many folds because they are natural products, being non narcotics, having no side effects and easily available at affordable prices. The goal of the present investigation was to estimate soluble sugar, chlorophyll, protein, lipids, phenols and starch from different plants which is presented in different proportion in plant species. The highest amount of soluble sugar, chlorophyll, protein, lipids, phenols and starch was obtained from *Azadirachta indica* (Neem), *Vinca rosea* (Sadahabad), *Mentha avensis* (Pudina), respectively.

**Key words :** Soluble sugar, Chlorophyll, Protein, Lipids, Phenols, Starch

**How to cite this paper :** Dwivedi, Golokja Krishna and Singh, Arvind Kumar (2012). Extraction of organic compounds from different three medicinal plant, *Ann. Pharm. & Pharm. Sci.*, **3** (1) : 45-46.

**Article chronicle :** Received : 31.03.2012; Revised : 10.04.2012; Accepted : 20.04.2012

Plant play a significant role in providing primary health care services to rural people. They serve as therapeutic agents as well as important raw materials for the manufacture or traditional and modern medicine. Substantial amount of foreign exchange can be earned by exporting medicinal plants to other countries. In this way indigenous medicinal plants play significant role of an economy of a country. Plants, as extracts and in various other forms, are being used for centuries in different traditional system of medicine for the treatment of human ailments, particularly those caused by pathogenic bacteria, fungi, as well as virus. Presence of various compounds and their uses has extensively been emphasized by number of workers (Despande *et al.*, 1980; Dougal, 1981; Collins, 1987). Progress in medicinal plants research has undergone a phenomenal growth during last two decades. World trend towards the utilization of natural plant remedies has created an enormous need for information about the properties and uses

of medicinal plants as antitumor, antianalgesic, insecticides, rotenoids etc. Demand on plant based therapeutics has increased many fold because they are natural products having no side effects and easily available at affordable prices (Govil *et al.*, 2002). The goal of present work is to analyze primary metabolites which are direct concerned with metabolic processes like-respiration, photosynthesis, protein, lipid synthesis.

In the proposed work different species were collected from College of Forestry, Department of Agriculture, T.D.P.G. College, Jaunpur and Department of Botany, Shibli National PG College, Azamgarh and Allahabad Agricultural Institute. Fresh leaves were washed and dried in an incubator at 37°C and made powder. This powder was later used for further analysis such as total soluble sugar and chlorophyll (Sadasivam and Manickam, 1992), protein Lowry *et al.* (1951), total lipids (Bligh and Dyer, 1959), total phenols (Bray and Thorpe, 1954), starch (Dubois *et al.*, 1951).

Highest amount of sugar was observed in the *Azadirachta indica* (Neem) *i.e.* 3.10 while lowest was obtained from *Vinca rosea* (Sadabahar) *i.e.* 0.04. Highest content of chlorophyll was observed in leaf of *Vinca rosea* (Sadabahar) *i.e.* 1.54 per cent, while lowest was obtained from *Mentha avensis* (Pudina) *i.e.* 0.05 per cent. Maximum amount of protein was estimated in *Azadirachta indica* (Neem) *i.e.* 9.54 per cent and lowest was obtained from *Vinca rosea* (Sadabahar) and

### MEMBERS OF THE RESEARCH FORUM

#### Address for correspondence :

**GOLOKJA KRISHNA DWIVEDI**, Department of Botany, T.D.P.G. College, JAUNPUR (U.P.) INDIA  
Email : gkdwivedi9670405164@yahoo.com

#### Coopted authors :

**ARVIND KUMAR SINGH**, Department of Botany, T.D.P.G. College, JAUNPUR (U.P.) INDIA

**Table 1 : Organic compounds from different three medicinal plant**

Name of plant	Amount of constituents					
	Soluble sugar (%)	Chlorophyll (%)	Protein (%)	Lipids (%)	Phenols (%)	Starch (%)
<i>Azadirachta indica</i> (Neem)	3.10	0.60	9.54	7.00	0.05	3.45
<i>Vinca rosea</i> (Sadabahar)	0.04	1.541	0.04	0.04	0.02	0.07
<i>Mentha avensis</i> (Pudina)	0.07	0.05	0.04	0.02	0.07	0.01

*Mentha avensis* (Pudina) *i.e.* 0.04 per cent and lowest amount of lipids was obtained from *Mentha avensis* (Pudina) *i.e.* 0.02 per cent. Maximum amount of phenols was found in *Mentha avensis* (Pudina) *i.e.* 0.07 per cent while lowest was found from *Vinca rosea* (Sadabahar) *i.e.* 0.02 per cent and starch was estimated as 3.45 per cent in *Azadirachta indica* (Neem) and lowest was calculated in *Mentha avensis* (Pudina) *i.e.* 0.01 per cent as show in Table 1.

## REFERENCES

**Bligh, E.G. and Dyer, W.J.** (1959). Biochemical methods of plant physiology. *Canadian Biochem. & Physiol.*, **37** : 911-915.

**Bray, H.G. and Thorpe, W.V.** (1954). Biochemical estimation of phenolic compounds. *Math. Biochem. Anal.*, **1**: 27-52.

**Collins, H.A.** (1987). Determinants of yield of secondary products in plants tissue cultures. *Adv. Bot. Res.*, **13** : 145-87 (Eng.)

**Despande, V.V., Mendulkar, K.N. and Sadre, N.L.** (1980). Antifertility activity of *Azadirachta indica* in mice. *J. Postgrad. Med. (Bombay)*, **26** : 167-170.

**Dobois, M.K. Gillies, J.K. Hamilton, Rebers, P.A. and Smith, F.** (1951). A colorimetric method for the determination of sugar, *Nature*, **168** : 167-171.

**Dougal, D.K.** (1981). Tissue culture and the study of secondary (natural) products. *Biochem. Plants*, **7**:2134.

**Govil, J.N., Singh, V.K. and Singh, G.** (2002) : Recent progress in medicinal plants. *Ethanomedicine & Pharmacog.*, **1** : 37 :51-55.

