Internat. J. Proc. & Post Harvest Technol.

Volume 4 | Issue 2 | December, 2013 | 138-139



Research Note

International Journal of Processing and Post Harvest Technology

Application of fermentation in food processing including its industrial aspects

■ SUBHA GANGULY*

All India Coordinated Research Project on Post Harvest Technology (ICAR) Department of Fish Processing Technology, Faculty of Fishery Sciences, West Bengal University of Animal and Fishery Science, Panchasayar, KOLKATA (W.B.) INDIA (Email : ganguly38@gmail.com)

*Author for Correspondence

Research chronicle : Received : 20.12.2012; Accepted : 30.11.2013

SUMMARY:

Fermentation is brought about by the conversion of sugars into ethanol chemically. The fermentation technology applicable to food processing sector is also popularly known as zymology or zymurgy. Fermentation is an important and popular technique in food processing technology. It is resulted from the chemical reaction resulting from the breakdown of higher carbohydrates to alcohols and organic acids or alcoholic derivatives.

KEY WORDS : Fermentation, Food processing

How to cite this paper : Ganguly, Subha (2013). Application of fermentation in food processing including its industrial aspects. *Internat. J. Proc. & Post Harvest Technol.*, **4**(2):138-139.

Free ermentation is a microbial technique and the reaction to be controlled in favorable and desirable conditions for food safety and quality after fermentation, especially in the production of alcoholic premium quality beverages like beer, wine and cider (Cavalieri *et al.*, 2003; Steinkraus, 1995; Ganguly, 2012a). The same technology is employed in the bred manufacturing industries for leavening activity brought about by the production of carbon dioxide by the microbial or yeast activity. The preservation effect during fermentation is attributed to the production of lactic acid in sour foods such as yoghurt, dry sausages, pickles, sauerkraut and vinegar (extremely diluted acetic acid) (Steinkraus, 1995; Harden and Young, 1906).

The fermentation technology under controlled conditions is an age old practice both in households and industries for food processing and preservation, be it alcoholic beverage products of edible products derived from vegetable, fish and meat sources (Steinkraus, 1995). Louis Pasteur, the renowned French chemist is the world famous and first known zymologist in history, who in 1856 established the pivotal role of yeasts in fermentation. Pasteur originally defined fermentation as "respiration without air" after regular performances of lengthy experimental protocols (Dubos, 1995). After observation of the breakdown of sugars to alcohols by the action of yeast, the pioneer concluded that the entire reaction is driven by the chemical catalytic action of certain forces called ferments inside the yeast cells. It was further observed that the yeast extracts can bring bout fermentation of sugars even also in the absence of viable yeast cells. In 1897, Eduard Buchner of Humboldt University of Berlin, Germany discovered that sugars are fermented in the absence of viable cells also in the fermentation mixture. The yeast cells secrete a chemical component called zymase. For his memorable contributions in research and discovery of cell-free fermentation, in 1907 Buchner was awarded with the prestigious Nobel Prize in Chemistry. In 1906, NAD⁺ was discovered out of studies carried out from ethanol fermentation.

Primary benefits :

Fermentation technology is primarily employed for the

preservation of different food by production of acids and alcohols, biological fortification and enrichment of food items with potential biogenic products like essential amino acids, easily digestible proteins, essential fatty acids and useful vitamins, neutralization of anti-nutritional factors, to diversify and enrich the diet with various aromas, flavours and textures in food substrates and decrease in requirements of further processing techniques like cooking etc (Steinkraus, 1995).

Fermented fish products available worldwide :

Specifically, in the fish processing technological research aspect, Bagoong, Faseekh, Fish sauce, Garum, Hákarl, Jeotgal, Hentak (Sarojnalini and Vishwanath, 1995), Rakfisk, Shrimp paste, Surströmming, Shidal (Ahmed *et al.*, 2012a; 2012b; Muzaddadi and Basu, 2012) and Ngari (Thapa *et al.*, 2004) are the popular fermented fish products worth mentioning.

Public health risks involved from consumption :

There are certain risks and health hazards associated with excess and regular consumption of fermented food products. In Alaska, since 1985, there has been increase in incidences of botulism exceeding the case reported in the Americas. This is mainly caused for the practice of allowing whole fish, fish heads and meat of animals like sea lions, walrus, whale flippers, birds, seal tallow, beaver tails etc. to ferment for prolonged periods before consumption by the resident Eskimos there. During this extended fermentation, if plastic wrappers or containers are used, then *Clostridium botulinum* gets a conducive condition to thrive in the micro-aerophilic condition inside the plastic containers (Ganguly, 2012b).

Alaska has witnessed a steady increase of cases of botulism since 1985. It has more cases of botulism than any other state in the United States of America. This is caused by the traditional Eskimo practice of allowing animal products such as whole fish, fish heads, walrus, sea lion, and whale flippers, beaver tails, seal oil, birds, etc., to ferment for an extended period of time before being consumed. The risk is exacerbated when a plastic container is used for this purpose instead of the old-fashioned, traditional method, a grass-lined hole, as the botulinum bacteria thrive in the anaerobic conditions created by the air-tight enclosure in plastic containers (U.S. federal agency Report).

Acknowledgement :

The authors are thankful to the AICRP on Post Harvest Technology (ICAR), Kolkata centre, WBUAFS.

LITERATURE CITED

- Ahmed, S., Dora, K.C., Sarkar, S., Chowdhury, S. and Ganguly, S. (2012a). Isolation and molecular characterization of *Bacillus* species from *Shidal* a fermented fish product of Assam. *Indian J. Fisheries*, Accepted.
- Ahmed, S., Dora, K.C., Sarkar, S., Chowdhury, S. and Ganguly, S. (2012b). Quality analysis of shidal- a traditional fermented fish product of Assam. *Indian J. Fisheries*, Accepted.
- Cavalieri, D., McGovern, P.E., Hartl, D.L., Mortimer, R. and Polsinelli, M. (2003). Evidence for S. cerevisiae fermentation in ancient wine. J. Molecular Evolution, 57(Suppl 1): S226-232.
- Dubos, J. (1995). Louis Pasteur: Free Lance of Science, Gollancz. In: K.L. Manchester. Louis Pasteur (1822-1895)- chance and the prepared mind. *Trends Biotechnol*, 13(12): 511-515.
- Ganguly, S. (2012a). Food Processing Technology. AV Akademikerverlag GmbH & Co. KG, Saarbrücken, Germany. LAP LAMBERT Academic Publishing (ISBN: 978-3-8383-8653-9).
- Ganguly, S. (2012b). Food Microbiology. AV Akademikerverlag GmbH & Co. KG, Saarbrücken, Germany. LAP LAMBERT Academic Publishing (ISBN: 978-3-8484-8217-7).
- Harden, A. and Young, W.J. (1906). The Alcoholic Ferment of Yeast-Juice. Proceedings of the Royal Society of London. 78(526): 369-375.
- Muzaddadi, A.U. and Basu S. (2012). An accelerated process for fermented fish (*seedal*) production in Northeast region of India. *Indian J. Animal Sci.*, **82**(1): 98–106.
- Sarojnalini, Ch. and Vishwanath, W. (1995). Nutritional characteristics of the two fermented fish products-Hentak and Ngari of Manipur. J. Indian Fisheries Association, 25: 75-85.
- Steinkraus, K.H. (1995). Ed. Handbook of Indigenous Fermented Foods. NY, Marcel Dekker, Inc.
- Thapa, N., Pal, J. and Tamang, J.P. (2004). Microbial diversity in ngari, hentak and tungtap, fermented fish products of North-East India. *World J. Microbiol. Biotechnol*, 20: 599-607.
- U.S. federal agency Report. Why does Alaska have more botulism? Centers for Disease Control and Prevention.

