

Evaluavation of bio-agents against *Curvulara lunata*, the causal agent of grain discolouration in rice

K. SUMANGALA AND M.B. PATIL



International Journal of Plant Protection, Vol. 4 No. 1 (April, 2011) : 193-195

See end of the article for authors' affiliations

Correspondence to :
K. SUMANGALA
Department of Plant
Pathology, College of
Agriculture, University
of Agricultural
Sciences, RAICHUR
(KARNATAKA)
INDIA
Email : sumakoulagi@
gmail.com

SUMMARY

In the present research studies, the antagonistic microorganisms were tested for their efficacy in inhibition of growth of *C. lunata*. Many fungi have been isolated from discolored grain, in which *Curvularia lunata* (Wakker) Boedign was found dominant pathogen (35.30%) of grain discolouration of rice in rice growing tracts of Tungabhadra Project Area and Upper Krishna Project Area of Karnataka state in India comprising of Raichur, Koppal and Gulbarga districts during 2007. Among the different bio-agents, *Bacillus subtilis* (97.77%) followed by fungal bio-agent *Trichoderma viride* (96.44%) and *T. harzianum* (93.50%) were found to be effective. Seed treatment with *T. viride* was found to inhibit *C. lunata* association with grain discolouration with 90.05 per cent germination and 1170.00 vigour index followed by *Bacillus subtilis* with 87.99% germination and 989.11 vigour index.

Sumangala, K. and Patil, M.B. (2011). Evaluavation of bio-agents against *Curvulara lunata*, a causal agent of grain discolouration in rice. *Internat. J. Pl. Protec.*, 4(1): 193-195.

Key words :
Rice, Grain
discolouration,
*Curvularia
lunata*, Bio-
agents

Rice (*Oryzae sativa* L.) is most widely grown cereal crop of the world. Majority of the fungi, viz., *Curvularia lunata*, *Alternaria alternata*, *Fusarium moniliformae*, and *Heliminthosporium oryzae*, responsible for causing grain discolouration have been reported to be seed borne in nature (Ou, 1985; Mew *et al.*, 1988 and Singh, 1993). Grain discolouration now has assumed to be great importance in recent years because of the changes in cropping practices; intensive system like increased fertilizer application and yearly more rice seasons. But not too many resistant varieties combined with good yield characters are available for cultivation.

In the present investigation, an attempt has been made to identify the best bio-agent for the inhibition of mycelial growth of *C. lunata* and so also to get the higher seed germination and seedling vigour through seed treatment.

MATERIALS AND METHODS

The present investigation on grain discolouration in rice was carried out during

2006-2007 at the Department of Plant Pathology, College of Agriculture, Raichur, University of Agricultural Sciences Dharwad. Raichur is situated in North Eastern Dry Zone (Zone 2) of Karnataka State at 16°12' N latitude/ 77°21' E longitude with an altitude of 389.37 m above mean sea level.

Collection of the samples:

Discoloured rice grains were collected from different rice growing areas of Raichur, Gulbarga, Koppal and Dharwad districts. The collected samples were packed in cloth bags and stored at room temperature (25 ± 2° C) for further investigation. Fungi were isolated and identified based on the morphological characteristics.

Dual culture test:

Bioagents obtained indigenously and from Tamil Nadu Agricultural University (TNAU), Coimbatore were tested for their efficacy under *in vitro* using dual culture technique as well as seed treatment.

Bioagents were evaluated for their

Received :
December, 2010
Accepted :
February, 2011

efficacy through dual culture technique. The bioagents and the test fungus were inoculated side by side on a single Petridish containing solidified PDA medium. Three replications were maintained for each treatment with one control by maintaining only pathogen and bio-agent, separately. They were incubated for eight days. The diameter of the colony of both bioagents and the pathogen was measured in two directions and average was recorded. Per cent inhibition of growth of the test fungus was calculated by using the formula of Vincent (1927).

Seed treatment with biocontrol agent:

The efficacy of biocontrol agents were studied, The seeds of paddy were soaked in culture of fungal and bacterial bio-agent for 4 hrs followed by 30 minute shade drying and the seeds were inoculated with spore suspensions of (1×10^5 spores/ ml) *C. lunata*. The seeds treated with the fungus alone was (control inoculated) and distilled water (control uninoculated) were maintained for comparison. Forty seeds were sown separately in pots having sterilised soil for each treatment. The seed germination and seedling vigour were calculated 21 days after sowing, as per the method of Abdulbaki and Anderson (1976).

Seedling vigour was measured by calculating vigour index. The vigour index was calculated by using the following formula given by Abdul-Baki and Anderson (1976):

$$\text{Seedling vigor} = (\text{Mean shoot length} + \text{Mean root length}) \times \text{germination}(\%)$$

RESULTS AND DISCUSSION

The frequency of mycoflora has to be detected to get better control measure for them. It is evident from earlier reports and present investigation that, the major fungi isolated from the discoloured rice samples from North Eastern Karnataka were *C. lunata* (35.30 %) followed by *A.alternata* (20.60 %), *F.moniliforme* (18.20 %) with other organisms including, *H.oryzae*, *P.grisea*, *Aspergillus*, *Penicillium*, *Rhizopus* spp. and other unidentified organisms (Table 1). Similar results were reported by Babo and Lokesh (1996).

Efficacy of bacterial and fungal bio-agents was studied under *in vitro* condition and the results on inhibition of mycelial growth of *C. lunata* were recorded and presented in Table 2 and Fig. 1. There was a significant difference among the bio-agents tested. Among the fungal bio agents, *T. viride* (96.44%) was

Table 1 : Mycoflora of discoloured grains of rice detected by Potato dextrose agar plate method

Sr. No.	Fungi	Per cent incidence	
		Unsterilized	Sterilized
1.	<i>Curvularia lunata</i>	35.30	8.53
2.	<i>Alternaria alternata</i>	20.60	--
3.	<i>Fusarium moniliforme</i>	18.20	10.50
4.	<i>Aspergillus</i> sp.	7.20	--
5.	<i>Penicillium</i> sp.	6.50	--
6.	<i>Helminthosporium oryzae</i>	5.10	--
7.	<i>Pyricularia grisea</i>	--	1.20
8.	<i>Rhizopus</i> sp.	1.50	--

Table 2 : Efficacy of bio-agents in inhibiting the growth of *C. lunata*

Sr. No.	Bioagent	Per cent inhibition
1.	<i>Bacillus subtilis</i> (E)	40.99 (39.21)**
2.	<i>Bacillus subtilis</i> (PGPR)	43.00 (41.02)
3.	<i>Bacillus subtilis</i> (I)	97.77 (81.25)
4.	<i>Pseudomonas fluorescens</i> (E)	41.99 (40.38)
5.	<i>Pseudomonas fluorescens</i> (I)	47.44 (44.10)
6.	<i>Trichoderma viride</i> (I)	96.44 (72.59)
7.	<i>Trichoderma harzianum</i> (I)	93.50 (75.23)
S.E.±		0.92
C.D. (P=0.01)		2.84

** Arcsine values




Fig. 1 : Evaluavation of bio-agents against *Curvularia lunata*

found to be significant in inhibiting the mycelial growth of *C. lunata* followed by *T.harzianum* 93.50. Among the bacterial bio-agents, *Bacillus subtilis* (I) (97.77%) was recorded maximum inhibition which was found superior among all the bio agents followed by *P. fluorescens* (I) (47.44%) and *Bacillus subtilis* (PGPR)

Table 3 : Effect of seed treatment with bioagents on seed germination and vigour index

Sr. No.	Treatments	Germination (%)	Mean root length (cm)	Mean shoot length (cm)	Vigour index
1.	<i>Bacillus subtilis</i> (E)	70.66 (57.22)**	3.3	4.3	538.00
2.	<i>Bacillus subtilis</i> (PGPR)	73.00 (59.75)	3.4	4.7	639.26
3.	<i>Bacillus subtilis</i> (I)	87.99 (70.65)	4.9	5.2	989.11
4.	<i>Pseudomonas fluorescens</i> (E)	74.66 (59.75)	3.5	4.6	632.12
5.	<i>Pseudomonas fluorescens</i> (I)	43.23 (41.10)	3.2	4.4	411.07
6.	<i>Trichoderma viride</i> (I)	90.05 (86.45)	5.1	6.5	1170.00
7.	<i>Trichoderma harzianum</i> (I)	85.00 (67.25)	4.5	6.1	816.49
8.	Control inoculated	26.00 (31.71)	3.5	4.5	208.00
9.	Control (Un inoculated)	87.10 (68.92)	3.0	3.6	512.90
S.E.±		2.20	0.08	0.07	109.61
C.D. (P=0.01)		8.82	0.25	0.22	322.18

** Arcsine values

(43.00%) and least per cent inhibition was recorded in *P. fluorescens* (E) (41.99%) which was at par with *B. subtilis*(E) (40.99). Similar studies were carried out by Saleem and Kandasamy (2002) and Sarhan and Shibly (2003).

Effect of seed treatment with bioagents:

The efficacy of different bio-agents was tested through seed treatment against *C. lunata* and the results of the experiment are presented in Table 3.

The results indicated that the bio agent *T. viride* (I) gave good germination per cent of 90.05 and vigour index (1170.00) followed by *B. subtilis* (I) recording germination (87.99%) and (989.11) vigour index, respectively and least germination (43.23) and with vigour index (411.70) was recorded in *P. fluorescens* (I) which is in agreement with Das *et al.* (1998).

Authors' affiliations:

M.B. PATIL, Department of Plant Pathology, College of Agriculture, University of Agricultural Sciences, RAICHUR (KARNATAKA) INDIA

REFERENCES

Adbulbaki, A.A. and Anderson, J.D. (1976). Vigour determination in soybean seed by multiple criteria. *Crop Sci.*, **13**: 630-633.

Babo, H.N.R. and Lokesh, S. (1996). Seed mycoflora of some paddy (*Oryzae sativa*) varieties in Karnataka (India). *Pl. Dis. Res.*, **11**: 49-59.

Das, B.C., Khalruzzaman, A.S.N. and Boro, C.C. (1998). Biological seed treatment for management of sheath blight of rice. *J. Mycol & Pl. Pathol.*, **28**: 45-47.

Mew, T.W., Bridge, J., Hibino, H., Bonman J.M. and Merca, S. D. (1988). Rice pathogen of quarantine importance. In: *Rice seed Health*. Proceedings of the International Workshop on rice health. IRRI, pp.101-115

Ou, S. H. (1985). *Rice Disease*. CAB International Mycological Institute, Kew, Surrey, UK, p. 380.

Singh, R. A. (1993). Management of seedborne disease of rice. *Indian Farmers Digest*, **24** (3-4): 22-24.

Saleem, B. and Kandasamy (2002). Antagonism of *Bacillus* species (Strain BC 21) towards *Curvularia lunata*. *Curr. Sci.*, **82** (12): 1456-1457.

Sarhan, A. and Shibly, M.K.A. (2003). Biological control of pathogenic fungi associated with rice seeds. *Arabian J. Pl. Prot.*, **21**(2): 102-108

Vincent, J ,M (1927). Distribution of fungal hypae in the presence of certain inhibitors. *Nature*, **159**: 850.
