

Management of *Macrophomina phaseolina* through non-systemic fungicides

A.M. MORADIA



International Journal of Plant Protection, Vol. 4 No. 1 (April, 2011) : 239-240

Correspondence to :

A.M. MORADIA
Dry Farming Research
Station, Junagadh
Agricultural University,
BHAVANAGAR
(GUJARAT) INDIA
Email: ammoradia@
gmail.com

SUMMARY

To study the efficacy of different fungicides under *in vitro* conditions the poisoned food technique was employed. Eight non-systemic fungicides were tested at 500, 1000, 2000 and 3000 ppm concentrations. The most effective fungicide was found to be the copper oxychloride where cent per cent inhibition of growth was recorded at 500 ppm concentration with toxicity index of 400. Next in order of superiority were boric acid and sodium metabisulphide which also inhibited cent per cent growth of the test fungus but at 2000 ppm concentration.

Moradia, A.M. (2011). Management of *Macrophomina phaseolina* in groundnut through systemic non-fungicides. *Internat. J. Pl. Protec.*, 4(1): 239-240.

Key words :

Management,
Macrophomina
phaseolina, Non-
systemic
fungicides, *In*
vitro

The poisoned food technique has been employed for different non-systemic fungicides under *in vitro* conditions by various scientists and reviewed here their conclusion. Mathukia (1982) reported that 92 to 100 per cent inhibition was observed in ceresan, captan, brassicol and benlate at 500, 750 and 1000 ppm. Vitavax, busan, thiram, cuman and thiophanate methyl (topsin) also inhibited the growth but only at 1000 ppm concentration. Prashanthi *et al.* (2000) stated that chlorothalonil was the most effective at 3000 ppm, where 82.45 % inhibition of mycelial growth was recorded. Hence, the present study was aimed for studying the efficacy of non-systemic fungicides for management of dry root rot (groundnut isolate) under *in vitro*.

To study the efficacy of different fungicides under *in vitro* conditions, the poisoned food technique was employed. Eight non-systemic fungicides were tested at 500, 1000, 2000 and 3000 ppm concentrations. Measured quantities of these fungicides were incorporated in Potato dextrose agar medium separately in flasks and then shaken well to given uniform dispersal of the fungicides. The medium containing different fungicidal concentrations were then poured into Petri plates under aseptic conditions. After cooling

and solidifying the medium in Petri plates it was inoculated with 4mm disc of actively growing culture of *M. phaseolina* (groundnut isolate) under aseptic conditions and then incubated at $30 \pm 1^\circ\text{C}$ temperature for 8 days. Observations on colony diameter was recorded in each of the fungicidal concentrations and the per cent inhibition of growth of *M. phaseolina* in each treatment was calculated by using the equation formula of Vincent (1927).

Perusal of data in Table 1 clearly indicates that all the fungicides were capable of inhibiting the growth of the fungus at various concentrations. The most effective fungicide was found to be the copper oxychloride where cent per cent inhibition of growth was recorded at 500 ppm concentration with toxicity index of 400. Next in superiority were boric acid and sodium metabisulphide which also inhibited cent per cent growth of the test fungus but at 2000 ppm concentration. Although the treatment of mancozeb and thiram were also found effective and provided cent per cent inhibition at highest *i.e.* 3000 ppm concentration. Dinocap and chlorothalonil were found mediocre in their performance for inhibiting the growth of test fungus, where average inhibition of growth

Received :
January, 2011
Accepted :
March, 2011

Table 1 : Per cent inhibition of growth of *M. phaseolina* at different concentrations of non-systemic fungicides under *in vitro* conditions

Non-systemic fungicide	Concentration (ppm)*				Mean	Toxicity index**
	500	1000	2000	3000		
Mancozeb (50% WP)	27.26	65.22	89.85	100.00	70.58	282.33
Chlorothalonil (Cavach 75% WP)	77.17	79.41	81.30	84.26	80.54	322.14
Cuprex (Copper oxy. 50% WP)	100.00	100.00	100.00	100.00	100.00	400.00
Boric acid	87.75	95.45	100.00	100.00	95.80	383.20
Sodium metaby.	11.37	26.71	100.00	100.00	59.52	238.08
Sulphur (Suphax 50% WP)	0.00	0.00	3.93	16.08	05.00	20.00
Thiram (Thiride 72% WP)	70.70	71.96	79.74	100.00	80.60	322.40
Dinocap (Karathan 48% EC)	51.22	58.78	80.85	82.41	68.32	273.26
Control	0.00	0.00	0.00	0.00	0.00	0.00
	Between fungicide			Within fungicide		
S.E. ±	0.66			1.08		
C.D. (P=0.05)	1.97			NS		
C V %	0.98			1.07		

* = Average of three replications,

** = Maximum toxicity index 300

NS = Non-significant

Copper oxy. = Copper oxychloride,

Sodium metaby. = Sodium metabysulphide

was 82.41 and 84.26 per cent respectively. Sulphur proved to be the least effective. Total inhibition of growth of *M. phaseolina* with thiram (Mathukia, 1982) and chlorothalonil (Prashanthi *et al.*, 2000) at 3000 ppm concentration has already been reported.

Prashanthi, S.K., Kulkarni, Shrikant, Sangam, V.S. and Kulkarni, M.S. (2000). Chemical control of *Rhizoctonia bataticola* (Taub.) Butler, the causal agent of root rot of safflower. *Pl. Disease Res.*, **15** (2) :186-190.

Vincent, J.M. 1927. Distortion of fungal hyphae in presence of certain inhibitors. *Nature*, **150** : 850.

REFERENCES

Mathukia, R.G. (1982). Investigations on *Macrophomina phaseolina* (Tassi.) Goid. causing root rot and leaf blight of groundnut (*Arachis hypogaea* L.). M.Sc. (Ag.) Thesis, Gujarat Agricultural University, Sardarkrishinagar, Gujarat (India).
